

Minimally invasive mitral valve surgery is associated with equivalent cost and shorter hospital stay when compared with traditional sternotomy

Pavan Atluri, MD, Robert L. Stetson, MHA, George Hung, BA, Ann C. Gaffey, MD, Wilson Y. Szeto, MD, Michael A. Acker, MD, and W. Clark Hargrove, MD


ABSTRACT

Objective: Mitral valve surgery is increasingly performed through minimally invasive approaches. There are limited data regarding the cost of minimally invasive mitral valve surgery. Moreover, there are no data on the specific costs associated with mitral valve surgery. We undertook this study to compare the costs (total and subcomponent) of minimally invasive mitral valve surgery relative to traditional sternotomy.

Methods: All isolated mitral valve repairs performed in our health system from March 2012 through September 2013 were analyzed. To ensure like sets of patients, only those patients who underwent isolated mitral valve repairs with preoperative Society of Thoracic Surgeons scores of less than 4 were included in this study. A total of 159 patients were identified (sternotomy, 68; mini, 91). Total incurred direct cost was obtained from hospital financial records.

Results: Analysis demonstrated no difference in total cost (operative and postoperative) of mitral valve repair between mini and sternotomy ($\$25,515 \pm \7598 vs $\$26,049 \pm \$11,737$; $P = .74$). Operative costs were higher for the mini cohort, whereas postoperative costs were significantly lower. Postoperative intensive care unit and total hospital stays were both significantly shorter for the mini cohort. There were no differences in postoperative complications or survival between groups.

Conclusions: Minimally invasive mitral valve surgery can be performed with overall equivalent cost and shorter hospital stay relative to traditional sternotomy. There is greater operative cost associated with minimally invasive mitral valve surgery that is offset by shorter intensive care unit and hospital stays. (*J Thorac Cardiovasc Surg* 2016;151:385-8)



Cost associated with standard sternotomy and minimally invasive mitral valve repair.

Central Message

Total, operative, and postoperative costs were assessed for mitral valve repair performed by sternotomy and minimally invasive means.

Perspective

Economic concerns should not limit the use of minimally invasive mitral valve surgery. Minimally invasive mitral valve surgery can be performed with excellent outcomes in appropriately selected patients, which should also not limit application of this approach.

See Editorial Commentary page 389.

Minimally invasive mitral valve surgery has become increasingly prevalent. Several studies demonstrating equivalent operative results with excellent long-term

outcomes, namely freedom from recurrent mitral regurgitation, have provided support for this approach to the mitral valve.¹ It has been demonstrated repeatedly that minimally invasive mitral valve repair is associated with longer operative and cardiopulmonary bypass times.² One would expect that with the increased need for reusable and disposable operative equipment (tissue retractors, cannulas, endoaortic balloons), there would

From the Division of Cardiovascular Surgery, Department of Surgery, University of Pennsylvania, Philadelphia, Pa.

Supported by the University of Pennsylvania, Department of Surgery.

Received for publication May 5, 2015; revisions received July 30, 2015; accepted for publication Aug 29, 2015; available ahead of print Sept 30, 2015.

Address for reprints: Pavan Atluri, MD, Director, Minimally Invasive and Robotic Cardiac Surgery, Director, Heart Transplantation and Mechanical Circulatory Support Program, Division of Cardiovascular Surgery, Department of Surgery, University of Pennsylvania, 6 Silverstein Pavilion, 3400 Spruce St, Philadelphia, PA 19104 (E-mail: pavan.atluri@uphs.upenn.edu).

0022-5223/\$36.00

Copyright © 2016 by The American Association for Thoracic Surgery

<http://dx.doi.org/10.1016/j.jtcvs.2015.08.106>

Scanning this QR code will take you to the article title page.



Abbreviations and Acronyms

ICU = intensive care unit
IQR = interquartile range

be greater cost associated with this approach. On the other hand, numerous groups, including ours, have demonstrated shorter intensive care unit (ICU) and hospital stays associated with a minimally invasive technique. Unfortunately, there are minimal data in the literature regarding the economic considerations of minimally invasive mitral valve surgery. We undertook this study to evaluate the health care costs associated with minimally invasive mitral valve surgery relative to a traditional sternotomy approach. We hypothesized that minimally invasive mitral valve surgery could be performed with equivalent cost and outcomes to that performed through a sternotomy.

METHODS

Study Design

To assess the cost associated with either minimally invasive (right minithoracotomy) or sternotomy approaches to mitral valve repair, all patients who underwent isolated mitral valve repair at the University of Pennsylvania from March 2012 through September 2013 were evaluated. To ensure equivalent cost comparison, only isolated mitral valve repairs were taken into account. Patients who underwent concomitant procedures, such as ablation procedures for atrial fibrillation, tricuspid valve repair, and coronary artery bypass grafting, were excluded. Because of the cost and variability of expense associated with mitral valve prosthesis, mitral valve replacements were also excluded. In addition, high-risk patients (Society of Thoracic Surgeons risk score >4) were excluded from this analysis to have "like" subsets. Robotic repairs were also excluded. One hundred fifty-nine patients were identified who fit these criteria (sternotomy, $n = 68$; minimally invasive, $n = 91$).

Complete operative and postoperative costs associated with mitral valve repair were individually gathered from institutional billing records. Only direct hospital costs were used in this analysis; indirect costs and overhead were not incorporated. The cost components were grouped according to major subgroups for each patient and then further averaged according to the approach. Standard preoperative demographic and operative variables were compared between groups with our institutional Society of Thoracic Surgeons database. Postoperative outcomes and major complications (death, wound infection, stroke, reoperation for bleeding, atrial fibrillation, hospital and ICU stays, and survival) were also analyzed. The University of Pennsylvania Institutional Review Board approved these data for use in research, with a waiver of patient consent.

Operative Approach

All patients who underwent a minimally invasive mitral valve operation through a limited right thoracotomy with endoscopic assistance were incorporated into the minimally invasive cohort. Aortic occlusion was achieved with an endoaortic crossclamp (Edwards Lifesciences Corporation, Irvine, Calif) in most cases (81%); in the remainder of the operations, a Chitwood clamp was used. We performed the minimally invasive operation with a limited right anterior thoracotomy and femoral cannulation, as previously described.^{3,4} For the sternotomy subgroup, the

mitral valve was approached through a standard full sternotomy with standard central bicaval cannulation. Standard and similar operative repair techniques were used for both approaches.

In the setting of reoperative surgery in the minimal access cohort, the operation was performed through a similar right anterior thoracotomy. Once cardiopulmonary bypass was established through the femoral vessels, the lung was dissected free of the right lateral pericardium and the pericardium was opened. If necessary, intrapericardial adhesions were dissected free. The endoaortic balloon was used for aortic occlusion. The mitral valve was exposed through the interatrial groove, and the operation was conducted in standard fashion.

Statistical Analysis

Continuous variables are expressed as the median with interquartile range (IQR), and categorical variables are presented as proportions. Costs are expressed as mean \pm SD. Differences between groups were assessed with the Fisher exact test for categorical variables, the independent Student *t* test for normally distributed continuous variables, and the Mann-Whitney *U* test for nonparametrically distributed values. All tests were 2-tailed. The statistical analysis was performed with IBM SPSS Statistics (version 20.0; IBM Corporation, Armonk, NY).

RESULTS

Preoperative Variables

There was no difference in median preoperative Society of Thoracic Surgeons score between the minimally invasive and sternotomy cohorts (0.47; IQR, 0.36-0.95 vs 0.61; IQR, 0.34-1.35; $P = .25$), demonstrating equivalent operative risks. Additionally, there was no difference in age (59 years; IQR, 52-66.5 years vs 57 years; IQR, 47-69 years; $P = .5$), sex (39.6% vs 38.2% female; $P > .999$), or ethnicity (87.9% vs 91.1% white; $P = .8$) between minimally invasive and sternotomy cohorts, respectively. An equivalent percentage of patients in each group had a previous sternotomy (8.6% vs 9.6%; $P = .8$).

Operative, Postoperative, and Total Costs Associated With Mitral Valve Repair

Analysis of total direct cost demonstrated equivalent costs between minimally invasive and sternotomy cohorts ($\$25,515 \pm \7598 vs $\$26,049 \pm \$11,737$; $P = .74$). There was a higher operative cost associated with the minimally invasive cohort than with the sternotomy cohort ($\$17,246 \pm \3823 vs $\$13,786 \pm \2594 ; $P < .0000001$). This was offset by significantly lower postoperative costs for the minimally invasive cohort ($\$8268 \pm \6501 vs $\$12,263 \pm \$10,836$; $P = .008$).

Breakdown of the operative costs revealed the following major subcategories (minimally invasive vs sternotomy): operative supplies, $\$7539 \pm \3577 versus $\$5350 \pm \1950 ($P = .0000002$); operative time, $\$3595 \pm \698 versus $\$3120 \pm \650 ($P = .00002$); perfusion, $\$3071 \pm \619 versus $\$2125 \pm \619 ($P < .000001$); nursing, $\$1685$ versus $\$1636 \pm \287 ($P = .2$); laboratory and blood bank,

Download English Version:

<https://daneshyari.com/en/article/2978958>

Download Persian Version:

<https://daneshyari.com/article/2978958>

[Daneshyari.com](https://daneshyari.com)