



## CASE REPORT

# Renal sympathetic denervation for treatment of resistant hypertension<sup>☆</sup>

P. Araújo Gonçalves<sup>a,\*</sup>, M. Sousa Almeida<sup>a</sup>, P. Branco<sup>b</sup>, A. Gaspar<sup>b</sup>, Helder Dóres<sup>a</sup>,  
Maria Salomé Carvalho<sup>a</sup>, Maria João Andrade<sup>a</sup>, A. Weigert<sup>b</sup>, J. Diogo Barata<sup>b</sup>,  
A. Canha Gomes<sup>c</sup>, L. Raposo<sup>a</sup>, H. Mesquita Gabriel<sup>a</sup>, R. Campante Teles<sup>a</sup>, M. Mendes<sup>a</sup>

<sup>a</sup> Serviço de Cardiologia, Hospital de Santa Cruz, Carnaxide, Portugal

<sup>b</sup> Serviço de Nefrologia, Hospital de Santa Cruz, Carnaxide, Portugal

<sup>c</sup> Serviço de Anestesiologia, Hospital de Santa Cruz, Carnaxide, Portugal

Received 12 September 2011; accepted 5 February 2012

Available online 13 September 2012

### KEYWORDS

Resistant  
hypertension;  
Renal denervation;  
Renal arteries

**Abstract** Hypertension is an important cardiovascular risk factor and although there have been many improvements in pharmacological treatment, a significant percentage of patients are still considered resistant. The authors describe two cases of radiofrequency renal sympathetic denervation that illustrate the feasibility of this new technique for the treatment of resistant hypertension. The procedure consists of the application of radiofrequency energy inside the renal arteries to ablate afferent and efferent sympathetic renal activity, which has been implicated in the pathophysiology of hypertension.

© 2011 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L. All rights reserved.

### PALAVRAS-CHAVE

Hipertensão arterial  
resistente;  
Ablação por  
radiofrequência;  
Artérias renais

### Ablação da atividade simpática renal para tratamento da hipertensão arterial resistente

**Resumo** A hipertensão arterial é um importante fator de risco cardiovascular e apesar dos inúmeros avanços na terapêutica farmacológica, existe ainda uma percentagem importante de doentes que são resistentes. Os autores descrevem 2 casos clínicos de ablação da atividade simpática renal por radiofrequência, que ilustram a exequibilidade de uma nova técnica para o tratamento da hipertensão arterial resistente. O procedimento consiste na aplicação de energia de radiofrequência nas artérias renais, para ablação da atividade simpática renal aferente e eferente, implicada na fisiopatologia da hipertensão arterial.

© 2011 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L. Todos os direitos reservados.

<sup>☆</sup> Please cite this article as: Araújo Gonçalves, P; Ablação da actividade simpática renal para tratamento da hipertensão arterial resistente. Rev Port Cardiol 2012. doi:10.1016/j.repc.2012.02.016.

\* Corresponding author.

E-mail address: [paraujogoncalves@yahoo.co.uk](mailto:paraujogoncalves@yahoo.co.uk) (P. Araújo Gonçalves).

## Introduction


Hypertension is an important cardiovascular risk factor that is implicated in myocardial infarction and stroke, which are major causes of mortality and morbidity in developed countries.<sup>1</sup> Despite its high prevalence, hypertension is far from being controlled, as shown by the PAP study on the prevalence, awareness, treatment and control of hypertension in Portugal.<sup>2</sup> This study, performed in 2003, found an overall hypertension prevalence of 42.1%, and only 46.1% of these patients knew they were hypertensive, 39.0% were taking antihypertensive medication and only 11.2% were controlled.

A technique has recently been developed to treat resistant hypertension by modulating renal sympathetic nerve activity using radiofrequency ablation.<sup>3–5</sup> The aim of the procedure is to reduce afferent and efferent nerve activity of the renal sympathetic plexus (Figure 1), whose role and importance in the pathophysiology of hypertension has been recognized for several decades.<sup>6,7</sup> The efficacy and safety of the technique was first documented in the Symplicity HTN-1 trial,<sup>4</sup> and a recently published 24-month follow-up study reported a fall of 32 mmHg in systolic blood pressure (BP) and 14 mmHg in diastolic BP.<sup>8</sup> The Symplicity HTN-2 randomized trial documented a reduction of 32 mmHg in systolic BP and 12 mmHg in diastolic BP at six months, with no significant changes in the control group, in a population of patients with mean baseline BP of 178/98 mmHg, medicated with an average of five antihypertensive drugs.<sup>5</sup>

This article presents two case reports of treatment of resistant hypertension by radiofrequency ablation of renal sympathetic activity. The procedures were approved by the hospital's ethics committee and performed with the patients' informed consent.

## Case report 1

A 50-year-old black man, a smoker, with no previous cardiovascular events, was being treated in the hypertension clinic. Secondary causes of hypertension had been ruled out. He was medicated with clonidine (0.45 mg/day), irbesartan (300 mg/day), amlodipine (10 mg/day), indapamide



**Figure 1** Application of radiofrequency energy in the renal arteries for ablation of afferent and efferent sympathetic activity.



**Figure 2** Symplicity® catheter and radiofrequency generator for renal artery ablation.

(1.5 mg/day) and bisoprolol (5 mg/day). Renal function was normal, with estimated glomerular filtration rate (eGFR) of 115 mL/min/1.73 m<sup>2</sup>.

His BP at the last consultation prior to the procedure was 173/118 mmHg, and 189/143 mmHg at the time of admission, despite his usual therapy. In a previous exercise test under medication, his BP had risen from 160/100 mmHg at rest to 200/120 mmHg at peak exercise. Echocardiography documented concentric left ventricular hypertrophy, with normal global systolic function and no significant valve abnormalities. Ambulatory BP monitoring (ABPM) showed severe systolic and diastolic hypertension and a nondipper pattern (daytime: 145/104/118/41 mmHg; nighttime: 136/96/111/40 mmHg; mean systolic and diastolic BP, mean BP and pulse pressure, respectively).

Previous abdominal CT angiography showed favorable renal artery anatomy for the procedure.

The procedure was performed with anesthesia support (propofol and remifentanyl) for pain control, and anticoagulation with heparin for a target activated clotting time of >250s. After gaining vascular access via the right femoral artery, abdominal aortography and selective renal artery angiography were performed. Following catheterization, a Symplicity® catheter (Medtronic, USA) was inserted into each renal artery, connected to a radiofrequency generator that is programmed and controlled automatically, independent of the operator (Figure 2). Six applications of radiofrequency energy (8 W) lasting 120s each were performed in the right and left renal arteries (Figures 3 and 4). Fluoroscopy time was 13 min, the quantity of contrast was 70 mL, and total procedure time was 53 min. There were no complications related to the intervention. Renal angiography at the end of the procedure showed normal flow and no signs of dissection; some irregularities in the luminal contour were observed due to edema at the ablation sites, as is common in such procedures (Figure 5).

At the end of the procedure, the access site was closed using an Angio-Seal® device. There were no post-procedural complications and the patient was discharged the following day; he continues to be monitored in the hypertension clinic, beta-blocker therapy having been discontinued. His BP was 150/86 mmHg one month after the procedure, a reduction of 23 mmHg in systolic BP and 32 mmHg in diastolic BP compared to before the intervention and with one less drug class. Renal function showed no significant changes following the procedure, with eGFR of 123 mL/min/1.73 m<sup>2</sup>.

The patient is scheduled to undergo ABPM at three and six months, and renal CT angiography at six months.

## Case report 2

A 59-year-old white woman, obese (body mass index 32), with type 2 diabetes, was being treated in the

Download English Version:

<https://daneshyari.com/en/article/3020607>

Download Persian Version:

<https://daneshyari.com/article/3020607>

[Daneshyari.com](https://daneshyari.com)