

# Sleep Neurobiology and Critical Care Illness



Xavier Drouot, MD, PhD<sup>a,b,c,\*</sup>, Solene Quentin, MD<sup>a,b,c</sup>

## KEYWORDS

- Sleep alterations • Neurobiology • Sleep EEG pattern • Sleep organization
- Circadian rhythms

## KEY POINTS

- Intensive care unit (ICU) patients experience severe sleep alterations, with reductions in several sleep stages, marked sleep fragmentation, low sleep continuity, and circadian rhythm disorganization.
- The numerous sources of these sleep alterations are associated with disruptions of sleep neurobiological processes and sleep dynamics that can alter sleep restorative functions.
- Understanding the neurobiology of sleep in the ICU is a major challenge for future sleep studies in critically ill patients.

## INTRODUCTION

That critical illnesses and environmental factors in intensive care units (ICUs) are associated with sleep disturbances was recognized shortly after the first ICUs were created. Many studies documented objective lack of restorative sleep.<sup>1–4</sup> Sleep alterations in critically ill patients differ from sleep changes observed in ambulatory patients (such as patients with sleep apnea syndrome) in pathophysiology as well as the consequences of sleep loss.

The literature regarding consequences of sleep deprivation on health is growing rapidly but ICU patients are unlikely to avoid the biological and neurobehavioral repercussions of sleep loss. To appreciate all of the phenomena triggered by sleep loss in the ICU, it is important to understand the neurobiology of healthy sleep and the specific neurobiological derangements of sleep in critically ill patients.

---

Funding sources: None.

Conflicts of interest: None.

<sup>a</sup> CHU de Poitiers, Department of Clinical Neurophysiology, Hôpital Jean Bernard, 2 rue de la Milétrie, Poitiers 86000, France; <sup>b</sup> Univ Poitiers, University of Medicine and Pharmacy, 6 rue de la Milétrie, Poitiers 86000, France; <sup>c</sup> INSERM, CIC 1402, Equipe Alive, CHU de Poitiers, Cours Est J. Bernard, Poitiers 86000, France

\* Corresponding author. CHU de Poitiers, Department of Clinical Neurophysiology, Hôpital Jean Bernard, 2 rue de la Milétrie, Poitiers 86000, France.

E-mail address: [xavier.drouot@chu-poitiers.fr](mailto:xavier.drouot@chu-poitiers.fr)

Crit Care Clin 31 (2015) 379–391

<http://dx.doi.org/10.1016/j.ccc.2015.03.001>

[criticalcare.theclinics.com](http://criticalcare.theclinics.com)

0749-0704/15/\$ – see front matter © 2015 Elsevier Inc. All rights reserved.

## NEUROBIOLOGY OF THE NORMAL SLEEP CYCLE

In human beings, sleep is composed of non-rapid eye movement (NREM) sleep, which can be light NREM (stages 1 and 2) or deep sleep (stages 3 and 4). The distinction between wake and NREM sleep is made by visual analysis of a 30-second portion of an electroencephalogram (EEG): during waking, the EEG shows a mix of fast oscillations (>16 Hz) and alpha rhythm (8–12 Hz) of low amplitudes (<10  $\mu\text{V}$ ). During light NREM sleep (stages 1 and 2), the background EEG is characterized by slow theta oscillations (frequency between 4 and 7 Hz) and sleep spindle and K complexes. These latter regularly occur and provide the landmark of stage 2. During deep NREM sleep (also called slow wave sleep; stages 3 and 4), the EEG shows slow waves (0.5–2 Hz) of high amplitude (>75  $\mu\text{V}$ ). Rapid eye movement (REM) sleep is a particular sleep stage in which the EEG shows theta and alpha rhythms. Identification of REM sleep is based on the presence of rapid eye movements identified on electro-oculograms and complete chin muscle atonia. During REM sleep, the brain is highly active, and most dreams and nightmares occur during this stage.

Human sleep is monophasic, and is programmed to occur during nighttime. The sleep-wake cycle is accurately organized and controlled. A sleep deficit elicits a compensatory increase in the intensity and duration of sleep, and excessive sleep reduces sleep propensity. This process could be represented as a sleep-pressure regulation, which would maintain this pressure between an upper and a lower limit. Sleep homeostasis can be represented by the interaction of 2 main physiologic processes. The first process is known as process S, which increases during waking and declines during sleep. Electroencephalographic slow wave activity (SWA) corresponds with an indicator of sleep homeostasis and the level of SWA is determined by the duration of prior sleep and waking. The timing and propensity to fall asleep are also modulated by a circadian process. This second process is driven by the internal clock. This circadian rhythm is sensible to external factors that help to keep the sleep-wake cycle synchronized with night-day alternation. The main external synchronizing factors are light, physical activities, meals, and social interactions.

The quantity of sleep is acutely regulated, and sleep deprivation has many neurobiological consequences. On the day following 1 night without sleep, brain performances are severely decreased. The most visible behavior is an increased tendency to fall asleep, even when the person fights to remain awake. The night following the sleep deprivation is modified and a sleep rebound usually occurs. This sleep rebound triggers a lengthening of nighttime sleep, an increase in slow wave sleep, and an increase in REM sleep.

## METHODS FOR SLEEP STUDY IN INTENSIVE CARE UNIT PATIENTS

Sleep can be assessed in terms of quantity (total sleep time and time spent in each sleep stage), quality (fragmentation, sleep EEG patterns), and distribution over the 24-hour cycle.

Full polysomnography (PSG) is the only reliable tool for measuring sleep, especially in patients with marked sleep disturbances. Accurate sleep scoring requires the recording of at least 3 EEG signals (preferentially F4-A1, C4-A1, O2-A1), 2 electro-oculography signals, and a submental electromyography (EMG) signal. Additional signals are usually recorded, such as nasal and oral airflow, thoracic and abdominal belts, electrocardiogram, and pulse oximetry. Sound and light levels should be measured, although these data are not obtained routinely.

Download English Version:

<https://daneshyari.com/en/article/3108100>

Download Persian Version:

<https://daneshyari.com/article/3108100>

[Daneshyari.com](https://daneshyari.com)