

Congestive Heart Failure and Central Sleep Apnea



Scott A. Sands, PhD^{a,b}, Robert L. Owens, MD^{c,*}

KEYWORDS

- Congestive heart failure • Central sleep apnea • Cheyne stokes respiration
- Loop gain

KEY POINTS

- Congestive heart failure (CHF) is a common clinical syndrome among patients in the intensive care unit (ICU), who frequently require noninvasive or mechanical ventilation.
- CHF affects breathing control by increasing chemosensitivity and circulatory delay, predisposing to central sleep apnea, classically in a crescendo–decrescendo pattern of respiration known as Cheyne–Stokes respiration (CSR).
- Few data are available to determine prevalence of CSR in the ICU, or how CSR might affect clinical management and weaning from mechanical ventilation.

CLINICAL CONSIDERATIONS

Historical Perspective

An abnormal respiratory pattern has long been recognized as an ominous sign of congestive heart failure (CHF). The observations of 3 physicians who have lent their names to the pattern remain informative:

His breathing was very particular: he would cease breathing for twenty or thirty seconds, and then begin to breathe softly, which increased until he breathed extremely strong, or rather with violent strength, which gradually died away till we could not observe that he breathed at all. He could not lie down without running the risk of being suffocated, therefore he was obliged to sit up in his chair.

— John Hunter, 1781¹

Disclosure Statement: Dr S.A. Sands is supported by a National Health and Medical Research Council of Australia Early Career Fellowship and R.G. Menzies award (1053201). Dr R.L. Owens is supported by the National Institutes of Health K23 HL105542. He has previously consulted for Philips Respironics.

^a Division of Sleep Medicine, Brigham and Women's Hospital and Harvard Medical School, 221 Longwood Avenue, Boston, MA 02115, USA; ^b Department of Allergy, Immunology and Respiratory Medicine and Central Clinical School, Alfred Hospital and Monash University, 55 Commercial Rd, Melbourne, VIC 3004, Australia; ^c Division of Pulmonary and Critical Care Medicine, University of California San Diego, 9300 Campus Point Drive, #7381, La Jolla, CA 92037, USA

* Corresponding author.

E-mail address: rowens@ucsd.edu

Crit Care Clin 31 (2015) 473–495
<http://dx.doi.org/10.1016/j.ccc.2015.03.005>

criticalcare.theclinics.com

0749-0704/15/\$ – see front matter © 2015 Elsevier Inc. All rights reserved.

The patient suddenly developed palpitations and displayed signs of severe congestive heart failure. The only particularity in the last period of his illness, which lasted eight or nine days, was in the state of respiration. For several days his breathing was irregular; it would entirely cease for a quarter of a minute, then it would become perceptible, though very low, then by degrees it became heaving and quick, and then it would gradually cease again. This revolution in the state of his breathing occupied about a minute, during which there were about thirty acts of respiration.

—John Cheyne, 1818

This symptom [periodic breathing], as occurring in its highest degree, I have only seen during a few weeks previous to the death of the patient.

—William Stokes, 1854

The initial observations by Hunter, Cheyne, and Stokes were made in patients close to death. They were the first to note the characteristic waxing and waning respiratory pattern of “Cheyne–Stokes respiration” (CSR), a common pattern of central sleep apnea in patients with CHF. CSR is characterized by complete cessation of respiratory effort and airflow (apnea phase) alternating with profound hyperventilation (hyperventilation phase). Such patterns occur during wakefulness, but are typically more prominent during sleep (Fig. 1). The apnea phase of CSR causes arterial hypoxemia, and the hyperventilation phase produces surges in blood pressure, arousal from sleep, and dyspnea (Fig. 2).^{2–5} Typically, CSR has a periodicity of 45 to 90 seconds, and occurs during non-rapid eye movement (REM) sleep stages 1 and 2. CSR severity is typically measured by quantifying the percent of total sleep time in CSR, and by the number of apneas and central hypopneas per hour of sleep (apnea–hypopnea index).⁶ Despite advances in the treatment of CHF (eg, β -blockade, spironolactone), untreated CSR remains highly prevalent during sleep and retains its association with




Fig. 1. Cheyne–Stokes respiration in a patient with congestive heart failure. Note the crescendo–decrescendo pattern of respiratory effort and airflow. The lung-to-ear circulatory delay can be approximated by the time from resumption of airflow to the start of the increase in oxygen saturation. EEG, electroencephalogram; EKG, electrocardiogram; SpO₂, oxygen saturation.

Download English Version:

<https://daneshyari.com/en/article/3108106>

Download Persian Version:

<https://daneshyari.com/article/3108106>

[Daneshyari.com](https://daneshyari.com)