



# A computed tomographic image study on the thickness of the infrazygomatic crest of the maxilla and its clinical implications for miniscrew insertion

Eric J. W. Liou,<sup>a</sup> Po-Hsun Chen,<sup>b</sup> Yu-Chih Wang,<sup>b</sup> and James Chen-Yi Lin<sup>b</sup>

Taipei, Taiwan

**Introduction:** The purpose of this study was to measure the thickness of the infrazygomatic (IZ) crest above the maxillary first molar at different angles and positions to the maxillary occlusal plane. These measurements were then used to derive clinical implications and guidance for inserting miniscrews in the IZ crest without injuring the mesiobuccal root of the maxillary first molar. **Methods:** Computed tomographic images of 16 adults were used. For each subject, on the coronal slice of the computed tomographic image containing the IZ crest and mesiobuccal root of the maxillary first molar, the IZ crest thickness and the miniscrew insertion position were measured by postulating that the miniscrew would be inserted at each 5° increment from 40° to 75° to the maxillary occlusal plane. **Results:** The IZ crest thickness above the maxillary first molar ranged from  $5.2 \pm 1.1$  mm to  $8.8 \pm 2.3$  mm, measured at 40° to 75° to the maxillary occlusal plane and 13 to 17 mm above the maxillary occlusal plane. **Conclusions:** By adopting 6 mm as the minimal IZ crest thickness for sustaining a miniscrew well throughout treatment and avoiding injury to the mesiobuccal root of the maxillary first molar, the clinical implication for miniscrew placement in the IZ crest of an adult is to insert it 14 to 16 mm above the maxillary occlusal plane and the maxillary first molar at an angle of 55° to 70° to the maxillary occlusal plane. (Am J Orthod Dentofacial Orthop 2007;131:352-6)

The infrazygomatic (IZ) crest is a placement site in the maxilla for orthodontic miniscrews or miniplates. It has been used successfully to provide skeletal anchorage for maxillary canine retraction, anterior retraction, en-masse anterior retraction, and intrusion of the maxillary posterior teeth.<sup>1-9</sup>

Anatomically, the IZ crest is a pillar of cortical bone at the zygomatic process of the maxilla (Fig 1). Clinically, it is a palpable bony ridge running along the curvature between the alveolar and zygomatic processes of the maxilla. In younger subjects, it is between the maxillary second premolar and first molar, whereas it is above the maxillary first molar in adults. A serious complication during miniscrew insertion in the IZ crest of adults is injury to the mesiobuccal root of the maxillary first molar. The maxillary interdental sites for insertion of miniscrews were shown radiographi-

cally.<sup>10-13</sup> It is recommended to insert miniscrews in higher rather than lower positions, such as midroot, at or beyond the root apex where interseptal bone is thicker and there is less chance of root injury.

However, without injuring the mesiobuccal root of the maxillary first molar, the appropriate sites and angles for insertion of the miniscrew in the IZ crest have not yet been well documented. The purposes of this computed tomographic (CT) image study were to show the thickness of the IZ crest above the maxillary first molar and the biting depth of a miniscrew in the IZ crest at different angles and positions to the maxillary occlusal plane, and to derive clinical implications for miniscrew insertion in the IZ crest without injuring the mesiobuccal root of the maxillary first molar.

## MATERIALS AND METHODS

The CT images of 16 adults (6 women, 10 men; average age,  $27.0 \pm 5.2$  years) with no craniofacial anomalies or systemic diseases were randomly selected from the database of the craniofacial CT imaging library of Chang Gung Memorial Hospital, Taipei, Taiwan. The machine was a spiral CT scanner (SOMATOM Sensation 16, Siemens, Munich, Germany). The CT images were formatted into

From the Department of Orthodontics and Craniofacial Dentistry, Chang Gung Memorial Hospital, Taipei, Taiwan.

<sup>a</sup>Assistant professor.

<sup>b</sup>Attending staff.


Reprint requests to: Dr Eric J. W. Liou, Chang Gung Memorial Hospital, 199 Tung-Hwa North, Taipai, 105, Taiwan; e-mail, lioueric@ms19.hinet.net.

Submitted, October 2004; revised and accepted, April 2005.

0889-5406/\$32.00

Copyright © 2007 by the American Association of Orthodontists.

doi:10.1016/j.ajodo.2005.04.044



**Fig 1.** Schematic illustration of IZ crest above mesio-buccal root of maxillary first molar.

standard dicom and reconstructed into continuous slices at 1-mm thickness each.

We used Analyze (Mayo Clinic, Rochester, Minn) for the CT image analyses. The images were first adjusted to the tissue threshold to extract the soft tissue. Then the images of the hard tissue were displayed simultaneously with their coronal, axial, and sagittal slices so that the IZ crest containing the mesio-buccal root of the maxillary first molar could be accurately located (Fig 2).

On the selected coronal slice, reference lines and points were established for measurements of lateral wall thickness of the maxillary sinus and IZ crest thickness (Fig 3). The first reference line was the maxillary occlusal plane, a plane between the mesio-buccal cusps of both maxillary first molars. The second reference line was the tangent line to the buccal surface of the mesio-buccal root of the maxillary first molar. The intersection point of this tangent line and the floor of the maxillary sinus was the sinus point (S point).

Through S point, another 8 reference lines were established at increments of 5°, from 40° to 75°, to the maxillary occlusal plane (Fig 3). These 8 reference lines also were the postulated miniscrew insertion angles. The intersections between these reference lines and the lateral surface of the IZ crest were points B1 to B8. The length between S point and B1 to B8, respectively, was IZ crest thickness. The perpendicular distances from B1 to B8, respectively, to the maxillary occlusal plane were the postulated miniscrew insertion positions. The lateral wall thickness of the maxillary sinus, 5 mm above S point, was also measured.

For each measurement, both the left and right sides were measured. They were compared with the Student *t* test for differences ( $P < .05$ ). The lateral wall thickness of the maxillary sinus and the IZ crest thickness at

various miniscrew insertion angles were compared with analysis of variance (ANOVA) ( $P < .05$ ).

## RESULTS

The right and left measurements of lateral wall thickness of the maxillary sinus or the IZ crest thickness at each miniscrew insertion angle were not significantly different ( $P > .05$ ). The right and left measurements were therefore pooled.

The IZ crest thickness varied with the postulated miniscrew insertion angles. The greater the postulated miniscrew insertion angle, the thicker the IZ crest would be (Fig 4, A). The IZ crest thickness was  $5.2 \pm 1.1$  mm when the postulated miniscrew insertion angle was 40° to the maxillary occlusal plane, and it was  $8.8 \pm 2.3$  mm with a postulated insertion angle of 75° to the maxillary occlusal plane (Table I). The IZ crest thickness at each postulated miniscrew insertion angle was significantly greater than the lateral wall thickness of the maxillary sinus ( $2.9 \pm 0.9$  mm) (Table I).

The miniscrew insertion position also varied with the postulated miniscrew insertion angle. The smaller the postulated miniscrew insertion angle, the higher the miniscrew insertion position relative to the maxillary occlusal plane (Fig 4, B). The miniscrew insertion position was  $17.0 \pm 3.7$  mm above the maxillary occlusal plane when the postulated insertion angle was 40° to the maxillary occlusal plane, and it was  $12.8 \pm 4.2$  mm above the maxillary occlusal plane when the postulated insertion angle was 75° to the maxillary occlusal plane (Table II).

## DISCUSSION

The IZ crest above the maxillary first molar, as shown in this study, was significantly thicker than the lateral wall of the maxillary sinus. As an insertion site for orthodontic skeletal anchorage, the IZ crest is usually used for a single miniscrew because of its thicker bone,<sup>2,4,7-9</sup> whereas the lateral wall of the maxillary sinus is used for miniplates with several miniscrews because of its thinner wall.<sup>3,5,6</sup> A thicker bone allows greater miniscrew biting depth, more osseous contact, and better primary stability of the miniscrew.<sup>14</sup> The IZ crest has 2 cortical plates—the buccal cortical plate and the sinus floor. This anatomic advantage allows for bicortical fixation and possibly contributes to better primary stability of the miniscrew.

The IZ crest thickness is also the miniscrew biting depth when the postulated miniscrew is inserted through and in the IZ crest. To obtain a 5- to 9-mm miniscrew biting depth without injuring the mesio-buccal root of the maxillary first molar, the miniscrew insertion position and the postulated miniscrew inser-

Download English Version:

<https://daneshyari.com/en/article/3118794>

Download Persian Version:

<https://daneshyari.com/article/3118794>

[Daneshyari.com](https://daneshyari.com)