



Original Article

# Effects of transportation time after extraction on the magnetic cryopreservation of pulp cells of rat dental pulp

Mao-Suan Huang<sup>1,2,3</sup>, Wei-Jen Chang<sup>2,3†</sup>, Haw-Ming Huang<sup>4</sup>,  
Yen-Chuang Lin<sup>3</sup>, Yen-Hua Huang<sup>5,6</sup>, Jen-Chang Yang<sup>6,7</sup>,  
Sheng-Yang Lee<sup>3,5,8\*†</sup>

<sup>1</sup> Division of Oral and Maxillofacial Surgery, Department of Dentistry, Sin Kong, Wu Ho-Su Memorial Hospital, Taipei, Taiwan

<sup>2</sup> Dental Department, Taipei Medical University's Shuang Ho Hospital, Taipei, Taiwan

<sup>3</sup> School of Dentistry, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan

<sup>4</sup> Graduate Institute of Biomedical Materials and Engineering, Taipei Medical University, Taipei, Taiwan

<sup>5</sup> Department of Biochemistry, Graduate Institute of Medical Sciences, School of Medicine;

Center for Reproductive Medicine, Taipei Medical University Hospital, Taipei Medical University, Taipei, Taiwan

<sup>6</sup> Center for Teeth Bank and Dental Stem Cell Technology, Taipei Medical University, Taipei, Taiwan

<sup>7</sup> School of Dental Technology, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan

<sup>8</sup> Orthodontic Department of Wan-Fang Hospital, Taipei, Taiwan

Received 13 November 2010; accepted 17 February 2011

Available online 24 March 2011

## KEYWORDS

cell number;  
cryopreservation;  
magnetic-programmed  
freezer

**Abstract** *Background/purpose:* The aim of this study was to examine the effects of the time period between tooth extraction and the freezing procedure on pulp cells in frozen intact teeth of rats.

*Material and methods:* In total, 120 incisors from 30 rats were extracted and kept in transportation solution for 0 hour, 12 hours, and 24 hours. The tested teeth were divided into 2 experimental groups, where the extracted incisors were first frozen in a magnetic-programmed freezer (PF) or a traditional  $-20^{\circ}\text{C}$  freezer (TF). The tested teeth were then stored at  $-150^{\circ}\text{C}$  for 7 days. Incisors extracted from the opposite side of the same rat were treated as the non-frozen control. After thawing, the pulp was extracted and dissected. Hematoxylin and eosin staining was performed to observe the cell distributions. Cell densities in the odontoblast region and cell-rich zone were calculated using an optical microscope.

\* Corresponding authors. School of Dentistry, College of Oral Medicine, Taipei Medical University, 250 Wu-Hsing Street, Taipei 11042, Taiwan.

E-mail address: [seanlee@tmu.edu.tw](mailto:seanlee@tmu.edu.tw) (S.-Y. Lee).

† Wei-Jen Chang and Sheng-Yang Lee contributed equally to this work.

**Results:** When samples were immediately frozen after extraction, the cell density of odontoblastic cells was higher in the PF group compared with analogs which were frozen in the  $-20^{\circ}\text{C}$  freezer ( $P < 0.05$ ). In the cell-rich zone, cell densities of the PF group were significantly higher ( $P < 0.05$ ) than those of the TF group although the pulp cells had been stored in transportation solution for 24 hours before being frozen.

**Conclusions:** The freezing technique used in this animal study provided positive effects on pulp cell storage. In addition, the storage time before the freezing procedure is an important issue for cryopreserving pulp cells in intact teeth.

Copyright © 2011, Association for Dental Sciences of the Republic of China. Published by Elsevier Taiwan LLC. All rights reserved.

## Introduction

Regeneration is the mainstream in medical science in the 21<sup>st</sup> century, and also a goal for which scientists keep searching. The acquisition of good sources of stem cells (SCs) and accurately induced differentiation ability are 2 major factors for the success of regenerative medicine, and are also research subjects that scientists continue exploring. Meanwhile, determining ways to preserve the source tissue intact for further SC extraction is an important cornerstone of the development of SC technology.<sup>1,2</sup>

Since Gronthos et al. first discovered SCs in permanent teeth, diverse studies of pulp SCs have gradually become the focus of tissue engineering.<sup>3,4</sup> However, if mesenchymal SCs are deliberately extracted but not used in a timely manner, the entire process can be a waste of time and money. In this regard, an alternative way for dental tissue engineering is to save both the periodontal ligaments and pulp tissues at the same time, and thaw them for teeth reimplantation or mesenchymal SC extraction if needed in the future.

For cryopreservation of dental tissues, almost all studies focused on the cryopreservation of periodontal ligaments. Kaku and colleagues reported that the survival rate of reimplanted pulpless teeth after thawing was up to 97%.<sup>5</sup> This result demonstrates the possibility of the clinical application of a tooth bank.<sup>5</sup> In 2001, Laureys et al. froze pulpless teeth for 7 days and found that revascularization could be activated when the teeth were thawed and reimplanted.<sup>6</sup> However, technical developments concerning the cryopreservation of intact teeth with pulp are still limited. To cryopreserve pulp tissue, Lee and colleagues,<sup>7</sup> using the same technique as Kaku et al.<sup>5</sup> to cryopreserve extracted intact teeth, found that after thawing, dental pulp SCs could be successfully incubated from 73% of the thawed teeth. This result suggested that cryopreserved intact teeth not only can be used for autotransplantation but also can provide a source of dental pulp SCs.<sup>7</sup>

However, whether the storage time between teeth extraction and the freezing procedure influence the preserved cells is still unknown. Thus, in this animal study, we investigated the influence of storage time before cryopreservation on pulp cells thawed from cryopreserved intact teeth of the rat.

## Materials and methods

Thirty Wistar rats (with an average weight of  $91.4 \pm 19.4$  g) used in this study were purchased from the National

Laboratory Animal Center (Taipei, Taiwan). Central incisors of the maxilla and mandible of the 30 rats were extracted before the experiments. To obtain intact incisors, a rat was sacrificed, and its jawbone was excised. Then the surrounding bone of the incisor was carefully removed. Immediately after extraction, the teeth were cleaned with phosphate-buffered saline (PBS) and stored in a normal saline solution. Thirty extracted teeth were frozen in a programmed freezer (ABI, Chiba, Japan), supplied with a magnetic field (PF group), as described in a previous report.<sup>7</sup> Briefly, the teeth were transported at  $4^{\circ}\text{C}$  and then placed in a freezer at  $-5^{\circ}\text{C}$ . Teeth were maintained at that temperature for 15 min, and then cooled at a rate of  $-0.5^{\circ}\text{C}/\text{min}$  until the temperature reached  $-32^{\circ}\text{C}$ . During the freezing process, an electromagnetic field from a 75-mA current passed through a coil was applied to the samples. For the traditional freezing group (TF group), 30 extracted teeth were frozen in a traditional  $-20^{\circ}\text{C}$  freezer. During cryopreservation, the teeth were immersed in commercialized dimethyl sulfoxide (DMSO)-free cryoprotectant (modified-BAMBANKER, Lymphotec, Tokyo, Japan). After the freezing procedure, all experimental teeth were transferred to a freezer (MDF-11561, Sanyo, Osaka, Japan) and stored at  $-152^{\circ}\text{C}$  for 7 days. For each group, incisors extracted from the opposite side of the same rat were treated as the control. These control teeth were not subjected to either the freezing procedure or 7-day cryopreservation. To test the effects of transportation time between teeth extraction and the freezing procedure on pulp cells, the extracted incisors in the 2 experimental groups were kept in transportation solution for 0 hour, 12 hours, and 24 hours, with 10 samples used for each experimental condition.

After thawing at room temperature, teeth samples were fixed in 10% formalin and a decalcification solution in 10% formic acid for 48 hours according to a previous report.<sup>8</sup> Then a tooth sample was horizontally cut into cubes and dehydrated using a graded series of alcohol. After dehydration, the tooth cubes were embedded in paraffin and dissected at  $6\text{ }\mu\text{m}$  with a sliding microtome (Leica LM2500, courtesy of Meyer Instruments, Langham Creek, Houston, TX, USA). The slices were then dewaxed, stained with hematoxylin and eosin (H&E), and examined with an optical microscope.

Four microscopic images were taken of each pulp slice. For each microscopic image, two  $40 \times 15\text{-}\mu\text{m}$  squares were respectively defined at the odontoblast region and cell-rich zone (Fig. 1). Cell numbers were calculated in each square. Quantitative analysis was performed by comparing cell

Download English Version:

<https://daneshyari.com/en/article/3145842>

Download Persian Version:

<https://daneshyari.com/article/3145842>

[Daneshyari.com](https://daneshyari.com)