

Relationship between Postendodontic Pain, Tooth Diagnostic Factors, and Apical Patency

Ana Arias, DDS, Dr. Odont,* Magdalena Azabal, MD, DDS, Dr. Med,[†]
 Juan J. Hidalgo, MD, DDS, Dr. Odont,* and José C. de la Macorra, MD, DDS, Dr. Med*

Abstract

This study compares the incidence, degree, and length of postoperative pain in 300 endodontically treated teeth, with and without apical patency, in relation to some diagnostic factors (vitality, presence of preoperative pain, group, and mandible of treated tooth). Of the questionnaires received back, apical patency was maintained during shaping procedures with a #10 K-file in one group ($n = 115$) and not in the other ($n = 121$). There was significantly less postendodontic pain when apical patency was maintained in nonvital teeth. If pain appeared, its duration was longer when apical patency was maintained in teeth with previous pain or located in the mandible. Maintenance of apical patency does not increase the incidence, degree, or duration of postoperative pain when considering all variables together. (*J Endod* 2009;35:189–192)

Key Words

Apical patency, postendodontic pain, postoperative pain

From the *Department of Conservative Dentistry, Faculty (Estomatología II) of Odontology, Complutense University of Madrid, Spain; and [†]Private practice, Madrid, Spain.

Address requests for reprints to José C. de la Macorra, Department of Estomatología II, Faculty of Odontology, Complutense University, Plaza Ramon y Cajal s/n, 28040 Madrid, Spain. E-mail address: macorra@odon.ucm.es. 0099-2399/\$0 - see front matter

© 2008 Published by Elsevier Inc. on behalf of the American Association of Endodontists.
 doi:10.1016/j.joen.2008.11.014

Accumulation of soft tissue remnants or of dentinal debris in the apical region is a common event that can cause blockage of root canal, normally in its apical third. This can be avoided if patency of the apical foramen during the shaping procedure is granted (1). Currently, maintaining apical patency is recommended during shaping and cleaning endodontic procedures (2, 3).

Apical patency is a technique in which the apical portion of the canal is maintained free of debris by recapitulation with a small file through the apical foramen (4, 5). This technique allows prevention of blockage (6–9). The most predictable method is to regularly use a so-called patency file during cleaning and shaping procedures. This file can be defined as a small flexible K-file, which is passively moved through the apical constriction without widening it (10). The files used to obtain patency are often the same files initially used to negotiate canals (11).

Other advantages of this procedure are that it minimizes the risk of loss of length, reduces canal transportation and other accidents such as ledges (10), eases irrigation in the apical third of the canal (12), allows maintenance of the anatomy of the apical constriction (6), and improves the tactile sense of the clinician during apical shaping (10).

One of the alleged reasons for not using apical patency is the possible extrusion of debris through the apical foramen, a condition classically related with postoperative pain. In fact, the patency concept is controversial to some practitioners (13). Some think that the repeated pass of patency files, even of small ones, through the apex can cause by itself a periapical acute inflammatory response (9) and severe postoperative pain (13).

This procedure is taught in 50% of U. S. dental schools. In the other half, this technique is not taught, arguing that apical patency might increase the displacement of debris and subsequently irritate the periodontal ligament without producing a better healing (9). However, Tsesis et al. (14) found that maintaining apical patency did not reduce apical transportation or have an effect on loss of working length in curved root canals.

Other authors stated that maintaining apical patency would not cause more postoperative problems, providing it is satisfactorily made (9), and that its benefits exceed the possible injury it might cause (6) because it is intended exclusively to prevent dentinal chips being compacted into the apical region and forming a plug that can interfere with maintaining working length (15).

We have not found any published research assessing the incidence of postendodontic pain when apical patency was maintained in relation to when it was not. The purpose of this prospective study was to assess whether maintaining apical patency might influence the incidence, degree, or duration of postoperative pain, considering different tooth diagnostic factors such as pulpal status, preoperative pain, or the position or group of the teeth to be treated.

Materials and Methods

This research was conducted with the approval of the Ethics Committee of Clinical Research of Saint Carlos Hospital-Madrid.

Three hundred endodontic treatments were performed in uniradicular, biradicular, and multiradicular teeth by one endodontist, all of them in single visits. All patients were informed of the aims and design of the study, and written authorizations were obtained before their inclusion.

Exclusion criteria were the need for retreatment, pregnancy, failure to obtain patient's authorization, and the presence of accidents or complications during treatment (calcified canals, impossibility of achieving apical patency in any canal).

The following data were collected in clinical records. Pulpal vitality status (vital/nonvital) was assessed through thermic stimulation with ethyl chloride spray. This status was rechecked by testing the presence of bleeding during the endodontic access. If the thermic stimulation was positive and there was bleeding during endodontic access, the tooth was considered as vital and as nonvital if the stimulation was negative or there was no bleeding. The presence or absence of preoperative pain (yes/no) was noted. We asked the patients whether they had pain the days before the appointment. Group of teeth (posterior/anterior) and position (superior/inferior) were also collected.

Patients were given local anesthetics (lidocaine hydrochloride and epinephrine 1:80,000; Xilonibsa, Inibsa, Spain). The standard treatment procedure consisted of the following steps. Access was obtained by using 014 round carbide and Endo Z burs (Dentsply International, York, PA), with high-speed and water refrigeration at all moments. Full rubber dam was placed in the tooth to be treated. GLYDE (Dentsply Maillefer, Ballaigues, Switzerland) lubricant was placed at the entrance of the canals. Negotiation was done with a #10 file. Determination of working length was made with Root ZX apex locator (J Morita Europe GVBH, Frankfurt, Germany), with radiographic confirmation. Pulpal chamber was blot-dried with a cotton pellet. Lubricant was placed at the entrance of canals (ie, measurements were made along moist canals). A #10 file clamped to Root ZX apex locator was used to measure working length. Repetition of measurement was made with #12 and #15 files. If there was no agreement between measures obtained by using the 3 files, the measure that was dissimilar was reassessed. If disagreement persisted, the measure delivered with the thicker file was selected. Working length was confirmed with an intraoral periapical radiograph. In case of disagreement between radiographic and electronic measurements, the latter was selected. Shaping was done with Gates-Glidden (Dentsply Maillefer) and K-flexofile (Dentsply Maillefer). Master apical files ranged from #20–#30 in narrow and from #25–#40 in wide canals. After shaping of coronal and mid thirds, working length was confirmed by using apex locator. Cleaning with 5% NaOCl was performed during all procedures. AH-Plus sealer (Dentsply Maillefer) was deposited in canal by using an impregnated master cone twice. The #15 gutta-percha cones (Dentsply Maillefer) were laterally condensed with #20 nickel-titanium spreaders (Dentsply Maillefer) 1 mm short of working length.

Patients were randomly assigned to 1 of 2 groups: patency (P) and no patency (NP). In group P (initial $n = 150$), apical patency was maintained throughout shaping and cleaning procedures by using a #10 K-file between each instrument. In group NP (initial $n = 150$), all efforts were made to avoid surpassing the working length at all times during treatment.

TABLE 1. Chi-Square Test Results in Analysis of Incidence of Postoperative Pain (outcomes: yes/no)

Diagnostic factor	Condition	n	P value
Previous status	Vital	145	.47
	Nonvital	91	.03
Preoperative pain	Yes	76	.29
	No	160	.054
Group	Posterior	152	.07
	Anterior	84	.59
Position	Upper	121	.64
	Lower	115	.08

TABLE 2. Trend Test Results in Analysis of Degree of Postoperative Pain (outcomes: mild/moderate/severe)

Diagnostic factor	Condition	n	P value
Previous status	Vital	79	.36
	Nonvital	43	.39
Preoperative pain	Yes	44	.503
	No	78	.52
Group	Posterior	89	.37
	Anterior	33	.45
Position	Upper	63	.82
	Lower	59	.16

Patients were informed of the possible occurrence of pain for days after treatment and were given a questionnaire to be completed and returned. In it, they would record the presence or absence of postendodontic pain, its duration and level of discomfort rated as follows: mild pain: any discomfort of any duration that does not require treatment; moderate pain: pain that requires and is relieved with analgesics; and severe pain: any pain that is not relieved with treatment (analgesics).

Two hundred thirty-six of the 300 questionnaires were returned properly answered. Of these, 121 belonged to P group and 115 to NP group.

Results of groups P and NP related to incidence (yes/no), degree (mild, moderate, severe), and length (days) of postoperative pain were compared, attending to diagnostic factors: status of tooth (vital/nonvital), presence or absence of preoperative pain, group of teeth (posterior or anterior), or position (superior, inferior).

Results were analyzed with the χ^2 test for the incidence of pain, the trend test for its degree, and Mann-Whitney U test for its duration (SPSS 15 for Windows; SPSS Inc, Chicago, IL).

Results

Results are shown in Tables 1, 2, and 3.

Previous Vital Status

Differences were not statistically significant between P and NP groups regarding degree or duration of pain.

Incidence of postoperative pain differences was not statistically significant except in the group of nonvital teeth, where incidence was significantly lower ($P = .03$) when apical patency was maintained (Table 1).

Odds ratio was 2.53 (95% confidence interval [CI], 1.03–3.70). Odds of postendodontic pain in nonvital teeth in which apical patency was not maintained (NP) were between 1.03 and 3.70 times higher than in P group, in which patency was maintained.

Presence of Preoperative Pain

Differences were not statistically significant between P and NP groups regarding incidence or degree of postoperative pain.

TABLE 3. Mann-Whitney U Test in Analysis of Duration of Postoperative Pain (outcome: days)

Diagnostic factor	Condition	n	P value
Previous status	Vital	79	.48
	Nonvital	43	.89
Preoperative pain	Yes	44	.006
	No	78	.36
Group	Posterior	89	.22
	Anterior	33	.42
Position	Upper	63	.09
	Lower	59	.016

Download English Version:

<https://daneshyari.com/en/article/3149628>

Download Persian Version:

<https://daneshyari.com/article/3149628>

[Daneshyari.com](https://daneshyari.com)