



## Original Article

## Online assessment of sustained attention following sleep restriction

Avi Sadeh<sup>a,\*</sup>, Orrie Dan<sup>b</sup>, Yair Bar-Haim<sup>a</sup><sup>a</sup> The Adler Center for Research in Child Development and Psychopathology, Department of Psychology, Tel Aviv University, Tel Aviv 69978, Israel<sup>b</sup> Department of Psychology, The Max Stern Academic College of Emek Yezreel, Israel

## ARTICLE INFO

## Article history:

Received 18 November 2009

Received in revised form 24 December 2009

Accepted 4 February 2010

Available online 9 April 2010

## Keywords:

Sleep deprivation

Restriction

Actigraphy

Attention

Internet

Continuous performance test

## ABSTRACT

**Objective:** To assess the feasibility of conducting home-based sleep restriction studies with actigraphic monitoring of sleep and a new online continuous performance test (OCPT).

**Methods:** Thirty-four university undergraduate students (24 females, 19–30 years old) underwent repeated home assessments using self-administered OCPT following a regular night of sleep (8 h or more) and following sleep restriction (4 h of sleep) in a within-between subjects counter-balanced design. Actigraphy was used to monitor sleep. OCPT sessions were scheduled in the morning and the evening of days following normal and restricted sleep.

**Results:** OCPT measures demonstrated acceptable test–retest reliability. Actigraphic monitoring revealed good compliance with sleep requirements, and reported alertness reflected significant effects of sleep manipulation ( $p < .0001$ ). In comparison to performance following an 8-h sleep night, sleep restriction to 4 h was associated with a significant increase in omission errors in the high-target section of the test ( $p < .0005$ ) and with a significant increase in omission errors in the low-target section of the test ( $p < .01$ ).

**Conclusions:** These preliminary results support the feasibility of conducting home-based sleep restriction studies and the validity of the online version of the OCPT, suggesting that it may serve as a sensitive tool for assessment of sleep restriction/deprivation.

© 2010 Elsevier B.V. All rights reserved.

## 1. Introduction

The quest to unravel the function of sleep has led to numerous studies on the effect of sleep restriction/deprivation on cognitive performance [1–3]. Many studies have shown compromised neurobehavioral functioning following sleep loss, and a meta-analysis of sleep deprivation studies suggested that sleep deprived individuals function at a level that is equivalent to the ninth percentile of non-sleep-deprived individuals [1].

One domain of cognitive function that is particularly sensitive to sleep loss is sustained attention. Studies have repeatedly demonstrated that sleep loss in both children and adults leads to poor sustained attention as measured by different versions of the continuous performance test (CPT) and the psychomotor vigilance test (PVT) [3–9]. Furthermore, it has been shown that increase in sleep loss is associated with performance deterioration [4,5], and that these sustained attention tests are sensitive to circadian influences on alertness [10]. Finally, sustained attention tests are sensitive to intake of alertness promoting agents such as caffeine, amphetamines, and modafinil, reflected in improvements in cognitive function [11].

Traditionally, sleep restriction/deprivation studies have been conducted in laboratory settings to allow control over sleep schedule and cognitive testing. Recent research has shown the feasibility of conducting naturalistic sleep restriction studies using actigraphy to monitor compliance with prescribed sleep schedules [6,12,13]. Such studies are of great ecological value as they allow testing of sleep patterns in participants' natural settings. The ability to conduct studies on sleep deprivation outside the lab could be further enhanced by the development of reliable and valid Internet-based tests of cognitive function in general and of sustained attention in particular.

The goal of the present study was to assess the feasibility of conducting home-based sleep restriction study using actigraphy to monitor sleep and a new online continuous performance test (OCPT) to assess the effects of sleep restriction on cognitive function of young adults.

## 2. Methods

## 2.1. Procedure

Study design contrasted attentional capacities following two sleep-related conditions: (a) regular sleep – defined as a night of at least 8 h in bed; (b) restricted sleep – defined as a night of 4 h of sleep or less. To control for potential order effects, the

\* Corresponding author. Tel.: +972 3 6409296; fax: +972 3 6408074.

E-mail address: [sadeh@post.tau.ac.il](mailto:sadeh@post.tau.ac.il) (A. Sadeh).

participants were randomly assigned into two equal groups. One group was instructed to obtain 8 h of sleep during the first night of the protocol and to perform OCPT twice during the following day: (1) morning test, 30–60 min after their rise time and (2) evening test, between 18:00 and 21:00 h of the same day. These participants were instructed to sleep their regular sleep schedule on the following nights, and on the fourth night they were asked to sleep only 4 h and to perform the OCPT twice on the following day according to the same schedule (morning and evening tests). The other group received the same instruction with the only difference being that the first night was the restricted sleep night and the fourth night was an 8-h sleep night. All participants wore an actigraph to monitor their sleep–wake patterns. Participants were instructed to avoid any distractions including phone calls, TV, or other media or interpersonal stimulation while performing the OCPT. They were also instructed to shut down all other active windows or programs on the PC used to perform the test.

## 2.2. Participants

Forty undergraduate students enrolled and started the study. Participants received research credit hours according to their academic requirements. Four participants were excluded from the sample due to failure to complete all online tests. Two additional participants failed to comply with the sleep requirements as revealed through actigraphic monitoring. Thus, the final sample consisted of 34 participants (24 females), mean age = 23.67 years, SD = 1.99, range = 19–30. All participants reported good health and being free of prescription medications for health problems.

## 2.3. Measures

### 2.3.1. Online continuous performance test (OCPT)

The OCPT (eAgnosis Inc., Newark, DE) is a standard CPT designed and programmed for delivery over the Internet. The task is adapted for use with most Internet browsers (e.g., Internet Explorer, Fire Fox, Chrome, Apple-Safari) and can be operated from practically any computer connected to the Internet (for a demo see <http://www.checkadhd.com/onlineCPTresearch.php>). Millisecond accuracy in response time (RT) recording is achieved by using the end user's own CPU and transmitting the collected data back to a remote database at various points during task performance. Thus, the integrity of data collection does not depend on Internet speed or bandwidth. Furthermore, because the OCPT code for RT monitoring and recording is compact, the end user's computer characteristics and processor speed can rarely affect recording accuracy.

The task uses two geometric stimuli: equilateral triangles (5 cm sides) and circles (5 cm diameter), both presented in a light blue color delineated by a black one-pixel stroke contour. The geometric stimuli are presented in the middle of the screen against a gray background within a four-pixel stroke 7.5 cm × 11 cm black rectangle that is presented constantly throughout the task.

Each trial consists of a presentation of one geometric shape for 100 ms followed by a 1900 ms inter-trial interval. Participants are instructed to respond to the triangle shape as fast as possible without compromising accuracy by pressing the space bar on the computer's keyboard and to withhold responding to the circle shape. The task contains two conditions: low target frequency and high target frequency. The first half of the test (the low target frequency half) consists of 224 trials (56 targets, 168 non-targets) with a target to non-target ratio of 1:3. In this half, the task is boring and fatiguing. In the second half of the test (the high target frequency half), the target to non-target ratio is reversed and is set to 3:1 (168 targets, 56 non-targets). In this half of the test, the participant expects to respond most of the time but occasionally must inhibit the tendency to respond. These two conditions were chosen be-

cause they reflect a conceptual distinction in the attention/CPT literature between a condition that taxes sustained attention and attention focusing in a dull and boring environment (low target frequency) and a condition that taxes primarily the ability to inhibit a prepotent response (high target frequency) [12]. Throughout the task the geometric shapes are presented in a fixed, pseudo-random sequence. To minimize practice effects, each half of the task is preceded by a 2-min practice phase reflecting the target to non-target ratios of the actual test to follow. Subjects are not informed about the practice nature of these 2 min sessions and consider these parts of the test. Total net test time (including the two practice sessions) is 19 min. Three breaks are allowed (following the first and second practice sessions and following the low target frequency session). Participants are instructed not to exceed 2 min of break time. Responses with RTs that are faster than 150 ms are considered anticipatory and are removed from analyses. When a participant presses the spacebar more than once per stimulus presentation, only the data from the first response is included in data analysis.

The software automatically records omission and commission errors as well as reaction times. Three primary measures were extracted for analyses: error of omission, errors of commission, and response times. These measures are extracted per condition (low and high target frequencies) and can also be extracted by test quartiles per condition. All data is stored offline and is available as a Microsoft Excel spreadsheet.

### 2.3.2. Sleep assessment

Actigraphy and sleep diaries were used to monitor sleep. The diary data was only used to detect and remove possible artifacts from the actigraphic data. Actigraphy has been established as a reliable and valid method to assess sleep–wake patterns in infants, children, and adults [13–15]. Participants were given actigraphs (Mini Motionlogger, Ambulatory Monitoring Inc.) and were instructed to wear these on their non-dominant wrist. The actigraph was set to collect data in 1-min epochs and amplifier setting 18 which is the standard mode for sleep–wake scoring. Actigraphic raw data were translated to sleep measures using the Actigraphic Scoring Analysis (ASA) program for an IBM-compatible PC. These sleep measures have been validated against polysomnography with agreement rates for minute-by-minute sleep–wake identification higher than 90% [16].

To monitor compliance with the study's sleep manipulation we analyzed true sleep time as derived from actigraphy. True sleep time is defined as sleep time excluding all periods of wakefulness during the sleep period. Daily sleep logs, completed by the participants, were used to corroborate the actigraphic data. The sleepiness visual analogue scale (VAS) was used to assess subjective perception of sleepiness in the evening on the daily sleep logs. The VAS measures sleepiness on a continuous scale ranging from wide awake (0%) to nearly being asleep (100%). Participants are required to mark their stage of sleepiness by placing a tick mark on a long line. The specific relative location on the line represents their subjective level of sleepiness [17,18].

## 3. Results

The data analysis plan was aimed at assessing: (a) compliance of the participants with the experimentally imposed sleep schedule; (b) test–retest reliability of the OCPT and (c) sensitivity of the OCPT in detecting the effects of sleep restriction on cognitive performance.

### 3.1. Compliance with experimental sleep schedule manipulation

To assess compliance with experimental sleep requirements we examined actual sleep time as monitored by actigraphy and

Download English Version:

<https://daneshyari.com/en/article/3176376>

Download Persian Version:

<https://daneshyari.com/article/3176376>

[Daneshyari.com](https://daneshyari.com)