




ACTAS Derma-Sifiliográficas

Full English text available at
www.actasdermo.org



E-CASE REPORT

Retronychia: A Case Report Including Ultrasound Imaging and Surgical Treatment[☆]



M.L. Alonso-Pacheco,^{a,*} E. de Miguel-Mendieta,^b R. Maseda-Pedrero,^a M. Mayor-Arenal^a

^a Servicio de Dermatología, Hospital Universitario La Paz, Universidad Autónoma, Madrid, Spain

^b Servicio de Reumatología, Hospital Universitario La Paz, Universidad Autónoma, Madrid, Spain

KEYWORDS

Retronychia;
Proximal ingrowing nail;
Chronic paronychia;
Ultrasound;
Nail ablation

Abstract Retronychia is a recently described disorder caused by ingrowth of the proximal nail plate into the proximal nail fold. This situation provokes chronic paronychia refractory to antimicrobial therapy. Ultrasound has been proposed as the noninvasive method of choice to confirm the diagnosis and rule out other differential diagnoses, particularly local tumors and arthritic disease. The presence of 2 or more overlapping nail plates and a reduced distance between the root of the nail plate and the base of the distal phalanx could be the ultrasound hallmarks of this condition. Nail plate avulsion is the treatment of choice and is curative. Knowledge of retronychia is still limited among dermatologists, which can lead to diagnostic and therapeutic errors and delay. This has prompted us to present this new case.

© 2016 AEDV. Published by Elsevier España, S.L.U. All rights reserved.

PALABRAS CLAVE

Retroniquia;
Uña encarnada proximal;
Paroniquia crónica;
Ecografía;
Avulsión ungueal

Retroniquia: estudio ecográfico y tratamiento quirúrgico de un caso

Resumen La retroniquia es un proceso emergente en el que la parte proximal de la lámina ungueal se encarna en el pliegue ungueal proximal, produciendo una paroniquia crónica refractaria a terapias antimicrobianas. La ecografía se ha postulado como la prueba no invasiva de confirmación y para diagnóstico diferencial, sobre todo con afecciones tumorales y artríticas locales. La presencia de dos o más láminas ungueales superpuestas y la disminución de la distancia entre el origen de la lámina ungueal y la base de la falange distal podrían ser los criterios ecográficos distintivos de esta entidad. La avulsión quirúrgica de la lámina ungueal es la terapia de elección y curativa de la misma. Su conocimiento todavía está poco difundido entre los dermatólogos, ocasionando errores y demoras diagnósticas y terapéuticas, lo que nos ha motivado a la presentación de un nuevo caso.

© 2016 AEDV. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

[☆] Please cite this article as: Alonso-Pacheco ML, de Miguel-Mendieta E, Maseda-Pedrero R, Mayor-Arenal M. Retroniquia: estudio ecográfico y tratamiento quirúrgico de un caso. Actas Dermosifiliogr. 2016;107:e33–e37.

* Corresponding author.

E-mail address: mlalonsop@yahoo.es (M.L. Alonso-Pacheco).

Introduction

The term retronychia (proximal ingrowing nail), was coined by Berker and Rendall¹ in 1999. The condition emerged as a distinct entity in 2008, after the presentation of a cases series by members of the European Nail Society (ENS). Retronychia progresses with ingrowth of the proximal nail plate into the proximal nail fold (PNF), the presence of 2 or more overlapping generations of nail plate under the PNF, clinical signs of chronic proximal paronychia, and a yellowish nail that does not grow.^{2,3}

The pathogenesis of retronychia is not well known, although the condition is thought to be a complication of onychomadesis.³ Thus, repeated injury to the distal end of the nail triggers onycholysis that would first break the continuity and then the alignment between the nail matrix and the nail plate. By losing its proximal attachment to the nail bed and being subject to repeat injury, the nail plate gradually moves backward until it embeds itself in the cul-de-sac of the PNF. An inflammatory process with formation of granulation tissue is triggered in the cul-de-sac and nail bed, thus accounting for the clinical signs of paronychia. In addition, the presence of inflammatory phenomena would prevent the attachment of the new growing nail plate to the proximal part of the nail bed, thus leading again to onychomadesis, backward movement, and ingrowth into the PNF. This process can thus generate a vicious cycle in which up to 4 generations of nail plate overlap under the PNF.⁴

Case Description

The patient was a 22-year-old man who practiced mountaineering, for which he used tight boots. He came to the clinic with a 1-year history of absent nail growth on the great toe of the right foot and episodes of proximal exudative paronychia. Although the inflammatory signs improved to some extent with oral antibiotics, the patient continued to complain of pain in the PNF. He had taken ciprofloxacin during the previous 2 weeks because of growth of a culture of exudate of *Escherichia coli* that was sensitive to the antibiotic. He also reported what appeared to be a cut in the proximal part of the nail plate.

Examination revealed erythema and pain in the PNF on palpation, as well as proximal onychomadesis with a thick yellowish nail plate (Fig. 1A). Slight lateral deviation was observed in both great toes. No abnormal results were reported for the bacterial and fungal cultures or radiology examination of the toe.

Ultrasound study with a multifrequency linear probe (9-14 MHz) and high-resolution device (GE Medical Systems) revealed overlapping nail plates with a hypoechoic space between them, reduction of the distance between the root of the nail plate and the base of the distal phalanx at the level of the distal interphalangeal joint, as well as posterior acoustic shadowing deep under the nail plate. The PNF was thicker in the affected nail than in the healthy contralateral nail. The Doppler signal was slightly enhanced in the proximal part of the nail bed at the root of the matrix (Figs. 2 and 3).

Once the diagnosis of retronychia was confirmed, surgical avulsion of the nail was performed using an anterior approach, and the 2 nail plates were found to be overlapping at the proximal end, with granulation tissue and keratotic material between them (Fig. 4). Histopathology confirmed these findings. A new nail plate grew over the following 12 months, with slight distal thickening and yellowing (Fig. 1B).

Discussion

A total of 64 cases of retronychia have been reported, and most were included in multicenter studies carried out by experts from the ENS.^{2,4-15} Despite the wide age range reported (10-71 years), the mean age is 25 years. The condition mainly affects women (81%).

Retronychia affects the great toe unilaterally, although at least 7 cases of bilateral involvement^{1,2,4,10,13} and 9 cases of retronychia affecting the nails of the hands^{1,2,7,8} have been reported.

The most common triggers are repeated minor injuries, such as those that occur while jogging, walking, and dancing and those that result from having long nails or wearing heels or uncomfortable footwear. However, most patients do not associate their disease with these injuries. There have been at least 10 cases in which no trigger was identified^{2,4,11,12,14}; in a further 8 cases, the cause was acute injury such as



Figure 1 A, Erythema and edema in the proximal nail fold and proximal onychomadesis, with a thick, yellowish nail plate. B, Appearance of the new nail at 12 months, with distal yellowing and thickening.

Download English Version:

<https://daneshyari.com/en/article/3182334>

Download Persian Version:

<https://daneshyari.com/article/3182334>

[Daneshyari.com](https://daneshyari.com)