

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.elsevier.com/locate/apme](http://www.elsevier.com/locate/apme)

## Case Report

# A non healing wound treated with hyperbaric oxygen therapy

Tarun Sahni <sup>a,\*</sup>, Shweta Gupta <sup>b</sup>

<sup>a</sup> Senior Consultant, Department of Internal & Hyperbaric Medicine, Indraprastha Apollo Hospital, Sarita Vihar, Delhi-Mathura Road, New Delhi 110076, India

<sup>b</sup> Clinical Research Manager, Department of Hyperbaric Medicine, Hyperbaric Oxygen Unit, Indraprastha Apollo Hospital, Sarita Vihar, Delhi-Mathura Road, New Delhi 110076, India

## ARTICLE INFO

## Article history:

Received 27 January 2015

Accepted 5 February 2015

Available online 26 February 2015

## Keywords:

Wound

Healing

Hyperbaric

Ulcer

## ABSTRACT

Problem wounds represent a significant and growing challenge to our healthcare system. The incidence and prevalence of these wounds are increasing in the population resulting in growing utilization of healthcare resources. These problem wounds may arise from excessive pressure, trauma, venous insufficiency, diabetes mellitus, vascular disease, or prolonged immobilization leading to its difficult management with significant increases in cost, disability, and liability. Healing of such wounds is a dynamic pathway requiring the presence of oxygen for optimal restoration of tissue integrity and function needs good building blocks for repair and a good transport system to get the building blocks to the site of action. Hyperbaric oxygen therapy (HBOT) is used as a therapeutic modality which leads to an increase in tissue oxygen pressures at the wound site and hence allowing the reversal of a hypoxic state by increasing the oxygen diffusion within the plasma, consequently promoting angiogenesis, encouraging fibroblastic activity and supporting the tissues to resist against bacteria. It is used as an adjunctive treatment to enhance best-practice wound care and employing HBOT in a directed and appropriate way can significantly enhance wound healing efforts. We review a case of 17 year old girl with Grade I pressure ulcer over her right heel since 6 years and role of hyperbaric oxygen therapy as an adjunctive for its management.

Copyright © 2015, Indraprastha Medical Corporation Ltd. All rights reserved.

## 1. Introduction

Wound healing is a complex process involving an immediate sequence of cell migration leading to tissue repair and wound closure.<sup>1</sup> This sequence consists of removal of debris, control of infection, clearance of inflammation, angiogenesis, deposition

of granulation tissue, contraction, remodelling of the connective tissue matrix, and maturation. If wounds fail to undergo this sequence, chronic wounds may result. These are wounds that have existed for longer than 3 months and are unlikely to heal by themselves. The longer the wound exists, the less likely it is to heal with repeated uncomfortable cycle of dressings.<sup>2,3</sup>

\* Corresponding author. Tel.: +91 9810038010; fax: +91 (0) 11 26823629.

E-mail address: [aimhu@live.in](mailto:aimhu@live.in) (T. Sahni).

<http://dx.doi.org/10.1016/j.apme.2015.02.007>

0976-0016/Copyright © 2015, Indraprastha Medical Corporation Ltd. All rights reserved.

Clinically, chronic wounds are associated with pressure, trauma, venous insufficiency, diabetes mellitus, vascular disease, or prolonged immobilization.<sup>4-6</sup> Healing of such wounds needs good building blocks for repair and a good transport system to get the building blocks to the site of action. The normal healing cascade begins with an orderly process of haemostasis and fibrin deposition, which leads to an inflammatory cell cascade, followed by attraction and proliferation of fibroblasts and collagen deposition, and finally remodelling by collagen cross-linking and scar maturation.<sup>7,8</sup> Despite this orderly sequence of events responsible for normal wound healing, pathologic responses leading to fibrosis or chronic ulcers occur if any part of the healing sequence is altered.<sup>1,4,5</sup>


Currently, standard therapy for lower extremity wounds entails wound debridement, off-loading, systemic antibiotic therapy, and supportive medical therapy in an effort to heal wounds within a reasonable period of time. Hyperbaric oxygen therapy is a systemic treatment option that has emerged as a specialized and effective treatment option to manage such patients. When wound hypoxia is the systemic cause of the healing failure, providing oxygen at the wound site is, essentially, treating the cause. HBOT delivers very high concentrations of oxygen to the wound via the bloodstream, allowing it to kick start the healing process. In normal distal tissue, the partial pressure of oxygen is approximately 40 mmHg, and the partial pressure of oxygen in hypoxic wounds is about 10–20 mmHg. Following HBOT, the partial pressure of oxygen can increase to approximately 200 mm Hg.<sup>1-5</sup>

## 2. Case report

We report a case of 17 year old girl who is a known case of myelomeningocele and had undergone laminectomy when she was four month old. She has congenital vertical talus of right foot which was corrected with soft tissue release and tendon transfer (Peronei to Tibialis Posterior transfer) and Tibialis anterior lengthening. When she was 10 years old, Grade I pressure sore over right heel was detected with calcaneal deformity of the same foot. A surgery was performed for release of dorsiflexors of right foot and ankle. In 2006, she was diagnosed with infected corn in the same heel with cellulitis. It was managed with corn excision and full debridement.

The ulcer in her right heel had varied in severity over time and was deteriorating from the past few years. In 2010, there has been a large ulcer in her right foot which was treated by excision of callus. Bacterial culture revealed growth of *Escherichia coli* which was suppressed by antibiotics. Radiographical examination of right foot revealed no calcaneal spur with diminished bone density. In 2011, she was diagnosed with Osteomyelitis of calcaneum with discharging sinus.

She presented to our unit with a non healing infected wound on the heel of her right foot. Since healing was not obtained with topical antibiotic treatment and wound care, she was referred to be evaluated for hyperbaric oxygen therapy (HBOT). On examination, she had excruciating pain estimated as 5 using scale of 0–10, 0 being no pain and 10 being worse pain ever experienced.



**Fig. 1 – Wound on initial assessment.**

### 2.1. Wound profile

Past treatment of her ulcer had included pain management, debridement of wound bed with excision, but her current treatment regimen for ulcer consisted of dressings with T-Bact and betadiene. The cavity of her ulcer had soft slough with visible and deep base (Fig. 1). Wound odour was minimal and exudates presented as yellowish liquid oozing out which was being controlled by daily dressings.

### 2.2. Hyperbaric oxygen therapy

She was initially scheduled for 10 HBOT sessions. She responded slowly to the treatment but very promising progress was shown at the completion (Fig. 2). We decided to continue a further 10 treatment. In the 4th week, after 20th HBOT session, wound showed filling of cavity with minimal exudates. The surrounding skin was healthy, soft and



**Fig. 2 – After 10th HBOT session, yellow discolouration disappeared with better granulation tissue in the wound bed.**

Download English Version:

<https://daneshyari.com/en/article/3234871>

Download Persian Version:

<https://daneshyari.com/article/3234871>

[Daneshyari.com](https://daneshyari.com)