



Results of non-operative treatment following hip fracture compared to surgical intervention

M. Hossain*, V. Neelapala, J.G. Andrew

Department of Trauma and Orthopaedic Surgery, Ysbyty Gwynedd, Bangor LL57 2PW, United Kingdom

ARTICLE INFO

Article history:
Accepted 2 October 2008

Keywords:
Hip fracture
Non-operative management
Surgical fixation
Hip fracture mortality

ABSTRACT

We followed all consecutive hip fracture patients admitted between 2004 and 2006, identified cases in which the intention was to treat non-operative and compared their functional outcome and mortality with a similar cohort treated surgically over the same period. We recorded length of hospital stay, place of discharge, pre and post-fracture mobility and residence, 30 days and 1 year mortality, re-admission due to same fracture and delayed surgery. The group treated surgically was recruited and matched for age, gender, pre and post-fracture mobility, mental confusion and independence. 25 patients were treated non-operative. 22 patients treated surgically over the same time period matched the patient characteristics of the non-operative arm. The mean hospital stay was 13 days in both groups. There were 4 extra-capsular fractures (3 displaced) and 21 intra-capsular fractures (5 displaced) in the non-operative arm and 11 extra-capsular fractures and 9 intra-capsular fractures in the surgically treated arm. 4 patients from the non-operative treatment group underwent late surgery because of persisting hip pain 20 days–2 months after the index event (2 cannulated screws, 1 hemiarthroplasty, 1 total hip arthroplasty). 11 patients in the surgical treatment arm underwent dynamic screw fixation, 1 had cannulated screw, 1 had total hip replacement and 7 had hemiarthroplasty. 14 of the non-operative treated patients were mobile independently or with aid before fracture but only 9 patients retained their pre-fracture mobility following treatment, compared to 16 patients pre-fracture and 11 patients post-fracture after surgery. 16 patients treated non-operative were living independently prior to injury but only 7 went back to their own residence. Of the operatively treated patients 14 patients were living independently and 10 patients went back to their previous residence. 1 month and 1 year mortality in the non-operative treated group was 4/21 and 7/21 respectively compared to 1/20 and 5/20 in the operative fixation group. There was no statistically significant difference in mobility, residence or mortality between the two groups (Fisher exact test, $p > 0.05$). Non-operative management after hip fracture is suitable for medically unfit patients and does not result in statistically significant difference in functional outcome or mortality compared to patients treated surgically.

© 2008 Elsevier Ltd. All rights reserved.

Introduction

Most patients following hip fracture are treated surgically. However, some patients are not suitable for surgical intervention. Data from a recent Scottish hip fracture audit confirms that we are facing an increasing number of medically unfit patients presenting with hip fracture and consequently more patients are being treated non-operatively (4.3% in 2004 compared to 2.7% in 1998).¹ It may, therefore, be useful to know what happens to the group of patients treated non-surgically. There have been several historical series presenting functional outcome after non-operative management

of undisplaced hip fractures or incomplete Garden type I fractures, but we are not aware of any recent study that has presented the functional outcome of non-operative treated patients in general. We identified and followed up all consecutive patients with hip fracture treated non-operative and compared their functional outcome and mortality against an identical cohort of patients treated operatively over the same time period.

Patients and methods

Over a 3-year period we identified 25 patients with hip fracture where the initial intention was to treat the patient non-operatively. 1 patient who was deemed medically fit but refused surgery was excluded. We did not include any patients judged temporarily unfit who required medical optimisation before undergoing delayed

* Corresponding author. Tel.: +44 2920498905; fax: +44 2920498905.
E-mail address: munierh@doctors.org.uk (M. Hossain).

surgery. In our unit all patients with acute fracture deemed medically fit were treated with surgical fixation irrespective of pre-fracture mobility. Patients were not treated surgically if they were not medically fit, or minimally symptomatic following late diagnosis of hip fracture or in some cases were already bed-ridden and also had associated significant medical co-morbidity. All patients were reviewed daily by a senior elderly care physician with a view to optimising medical co-morbidities. Medical fitness and suitability for surgery was decided on the basis of American Society of Anaesthesiologists (ASA) grading. All patients were reviewed by more than one senior anaesthetist on more than one occasion who decided on medical fitness of patients and also agreed on suitability for anaesthesia and ASA grading before the patient was deemed medically unfit. Patients graded as ASA IV (severe systemic disease that is a constant threat to life) or above (ASA V – moribund patient not expected to survive for 24 h with or without surgery) were deemed medically unfit. The final decision about treatment was taken after discussion with the family and the patient (where applicable). Patients were then mobilised as soon as pain allowed. None of the patients had prolonged bed rest or traction. The plan of mobilisation was initially bed to chair followed by weight bearing on a walking frame and subsequent full weight bearing mobilisation as tolerated. Patients continued to be treated actively while an inpatient under the supervision of the elderly care team. Those previously resident in a residential or nursing home were, if possible, returned to their pre-fracture residence. Others were initially discharged to a community hospital and finally returned to the most suitable residence after social service and occupational therapy review. Fracture clinic follow-up was not arranged for any of these patients. There was, however, opportunity to re-refer any of these patients if the supervising elderly care physician felt that their medical condition had improved or minimally symptomatic patients with late diagnosed fractures were not making satisfactory progress. Patients were then readmitted, if necessary, to re-assess medical fitness and the need for surgery.

We also identified another cohort over the same time period – matched for age, gender, pre-fracture residence and mobility – who were treated with operative fixation and compared post-fracture mortality, mobility and independence between the two groups of patients. We consecutively recruited 22 patients who had operative fixation but 2 of them were visitors from an outside health region and were excluded as their final outcome could not be established. We recorded the following personal and clinical details: age, gender, mental confusion, pre and post-fracture residence, pre and post-fracture mobility, medical co-morbidities, type of fracture, fracture displacement, age of fracture, reason for non-operative management, readmission, subsequent change of decision and mortality at 1 month and 1 year. The length of hospital stay is the time spent in the acute hospital setting only and does not include time spent in a community hospital. Pre or post-fracture residence for a patient was categorised as living at own home with or without carers, or in an institutionalised setting (residential or nursing home). For patients transferred to a community hospital we considered post-fracture residence as the place where the patient was eventually transferred from the community hospital up to 3 months following discharge from the acute hospital. A patient's pre or post-fracture mobility was categorised as independently mobile indoors or outdoors, mobile with walking aids or immobile (wheel-chair or bed-bound). For patients not followed up following surgery we retrieved the clinical notes and searched entries from the elderly care clinic or physiotherapy out-patients to identify patient mobility up to 3 months following discharge from acute hospital setting. Acute confusional state was ignored, but patients were categorised as confused if chronic confusion or dementia was documented in the patient notes on admission.

Results

Over a 3-year period we identified 25 patients where the initial decision was to treat the hip fracture non-operative. There were 9 men and 16 women. The mean age was 82 (range 71–92). The mean hospital stay was 13 days (range 3–43 days). There were 4 extra-capsular fractures (3 displaced) and 21 intra-capsular fractures (5 displaced). 7 patients had valgus impaction fracture. 11 patients had delayed diagnosis of fracture and on the basis of history presumed to have sustained a fracture from 1 week to 9 weeks previously. 4 patients failed a trial of non-operative management and had to undergo late surgery because of persisting hip pain 20 days–2 months after the index event (2 cannulated screws, 1 hemiarthroplasty, and 1 total hip arthroplasty). Of the non-operative treated group, 14 patients were mobile independently or with aids before fracture, 9 of them maintained the same level of mobility afterwards. 11 out of 16 patients in the operatively treated group were mobile independently or with aids after fracture. The differences in pre and post-fracture mobility, residence and mortality between the two groups were not statistically significant (Fisher exact test, $p > 0.05$). In the non-operative treated group, 2 patients died while inpatients due to chest infection and 1 patient died due to septic shock and multiple organ failure 8–26 days following admission. No one developed pressure sores while an inpatient. None of the 4 patients who had delayed surgery died within the first year. The patient who underwent a total hip replacement in the delayed treatment group was suspected to have sustained hip injury around 6 weeks previously and presented with a healing intra-capsular fracture. Her ASA status was grade II. She was re-admitted from the clinic because of persisting hip pain and eventually underwent uneventful hip arthroplasty. 2 of the patients who had delayed surgery had delayed diagnosis of fracture. 1 of them had significant left ventricular failure (LVF) and the other chronic obstructive airway disease (COAD). These patients were categorised as ASA grade IV patients. Because of the combination of delayed diagnosis, minimal symptoms and significant medical conditions it was initially decided to treat these 2 patients non-operatively. The other patient had acute injury and was categorised as ASA grade IV with significant LVF and valvular heart disease. However, persisting uncontrolled pain on mobilisation prompted re-admission, reassessment and delayed surgery in these patients after discussion with the patient and the family regarding the high risks of surgery.

The mean time to surgery was 2 days in the operatively treated group. 11 patients underwent Dynamic Hip Screw (DHS) fixation, 1 patient had cannulated screw fixation, 1 patient with displaced intra-capsular fracture and co-existent rheumatoid arthritis had total hip arthroplasty and the rest had hemiarthroplasty. There were no cases of delayed diagnosis in the operative fixation group. Table 1 compares the pre and post-fracture characteristics of the operatively and non-operative treated groups. The patient who died within 1 month of surgery had bronchopneumonia. There were no re-admissions due to complications from surgery.

Discussion

The reported 1 year mortality in this series following non-operative management compares well with published UK national figures of mortality following surgical treatment of hip fractures (29–33%).^{3,9} The results also show that compared to a group of patients who were in better health and underwent hip fracture surgery non-operative management of hip fracture in a group of patients deemed medically unfit for surgery did not result in statistically significant difference in functional outcome or 30 day

Download English Version:

<https://daneshyari.com/en/article/3241403>

Download Persian Version:

<https://daneshyari.com/article/3241403>

[Daneshyari.com](https://daneshyari.com)