



Case Report

Reconstructive osteotomy for a malunited medial Hoffa fracture – A feasible salvage option



Binu Sasidharan*, Surendra Shetty, Satheesh Philip, Shivanand Shetty

Department of Orthopaedics, NMC Specialty Hospital, Abu Dhabi, United Arab Emirates

ARTICLE INFO

Article history:

Received 4 July 2015

Received in revised form 24 January 2016

Accepted 6 March 2016

Available online 26 March 2016

Keywords:

Intraarticular osteotomy

Malunion

Hoffa's fracture

ABSTRACT

A 35-year-old male presented with pain and restriction of movements of right knee of 9 months duration. He underwent open reduction and internal fixation for a coronal plane fracture of medial femoral condyle (Hoffa fracture) in another facility 9 months back. Radiological evaluation showed a malunited isolated medial Hoffa fracture. We did an intraarticular osteotomy as a salvage treatment and got a reasonable outcome. This case report analyses the difficulties in the management of this inherently unstable fracture pattern and the technique of intraarticular osteotomy.

© 2016 Prof. PK Surendran Memorial Education Foundation. Published by Elsevier, a division of Reed Elsevier India, Pvt. Ltd. All rights reserved.

1. Background

Hoffa fractures are intraarticular coronal plane fractures of the posterior femoral condyle. The lateral condyle is three times more commonly injured than medial one.¹ The preponderance to lateral condyle can be attributed to physiological valgus at knee.² An isolated medial Hoffa fracture is an extremely rare injury with less than 10 cases reported in the literature.³ Review of literature on Hoffa's fracture shows a case report of intraarticular corrective osteotomy done as a salvage treatment for malunited lateral Hoffa's fracture.⁴ Intraarticular corrective osteotomy for a malunited medial Hoffa fracture is technically even more challenging. Reconstructive osteotomy for a symptomatic malunited medial Hoffa fracture has not been reported in the English literature to the best of our knowledge. We report our experience with intraarticular corrective osteotomy for a malunited medial Hoffa fracture.

2. Case presentation

A 35-year-old male presented to us with pain and limitation of movements of right knee of 9 months duration. His pain aggravated with walking and was affecting his job. He suffered a road traffic accident 9 months back and had an isolated medial Hoffa fracture for which he underwent surgical fixation in another facility. He continued to have pain and limitation of movements after surgery. On examination, he had a fixed flexion deformity of

20° with further flexion up to 80° and an antalgic gait. Radiographs and computerized tomography images showed a malunited isolated medial Hoffa fracture with an intraarticular incongruity and two screws in situ (Fig. 1). He hoped to get a painless gait with improved range of motion.

3. Treatment

Reconstructive osteotomy of medial femoral condyle was planned. Knee was flexed to 40° and kept in a lazy figure of four position. Skin incision was made through the scar mark of previous surgery. The malunited fracture was exposed through a medial parapatellar arthrotomy. The Hoffa's fragment had migrated proximally by more than 10 mm (Fig. 2). A Homan's retractor was kept close to the posterior border of medial femoral condyle where it meets the femoral shaft. The two screws were removed. The osteotomy site was marked with a cautery. Intraarticular osteotomy was done keeping the medial collateral ligament and posterior oblique ligament attached to the Hoffa's fragment. The fibrosis and contracture of the posterior capsule made the reduction difficult. Two Steinmann pins were placed on the medial aspect of the medial femoral condyle and was used as a joystick to aid in anatomical reduction (Fig. 4). The reduced Hoffa's fragment was fixed with two 4 mm partially threaded cannulated cancellous screws introduced in anteroposterior direction (Figs. 3 and 4). The screw heads were countersunk. Stability of the construct was assessed and found to be satisfactory. The wound was closed in layers after achieving hemostasis. The knee was immobilized in a cylinder cast for 2 weeks.

* Corresponding author. Tel.: +971 553871350.

E-mail address: binuortho2000@gmail.com (B. Sasidharan).



Fig. 1. Computerized tomography image showing incongruity of articular surface.



Fig. 3. Post-operative radiograph – anteroposterior view.




Fig. 2. Intraoperative photograph showing depressed Hoffa fragment.

4. Outcome and follow-up

After 2 weeks, knee mobilization was started. Weight bearing walking was deferred till 2 months. Computerized tomography images at 3 months showed a congruous articular surface of distal femur (Fig. 5). At 3 months follow-up, his range of movements of knee was 5°–110° and he was walking full weight bearing. On further follow-ups at 4 months and 5 months, it is noticed that he develops pain when flexion beyond 110° is attempted. His residual pain and stiffness of knee could be due to the fibrosis and scarring from the revision surgery. However, intraarticular corrective osteotomy could moderately improve his range of motion of knee and gave a better articular congruity, which may delay the onset of secondary osteoarthritis.

5. Discussion

A written informed consent from the patient and formal permission from the institutional ethics committee is obtained for publication of this case report. A search into the currently available literature shows that several aspects of this rare injury pattern are controversial. Although it is agreed that surgical fixation is the preferred method of treatment in Hoffa's,¹ there is a lack of consensus in the surgical approach, mode of fixation, type of implant, and rehabilitation protocol in Hoffa fractures.¹ Failure of fixation, nonunion, and malunion are common complications with this intrinsically unstable fracture pattern. The vascular anatomy of the medial femoral condyle and the physiological stresses in the fixation construct are two major factors that surgeon needs to consider in the management of a medial Hoffa fracture.

The intraosseous blood supply to the medial femoral condyle has a watershed region of limited vascularity compared to lateral

Download English Version:

<https://daneshyari.com/en/article/3251742>

Download Persian Version:

<https://daneshyari.com/article/3251742>

[Daneshyari.com](https://daneshyari.com)