

Short Report

# Neuregulin 1 affects leptin levels, food intake and weight gain in normal-weight, but not obese, db/db mice

G. Ennequin<sup>a,b</sup>, N. Boisseau<sup>a,b</sup>, K. Caillaud<sup>a,b</sup>, V. Chavanelle<sup>a,b</sup>, M. Etienne<sup>a,b</sup>,  
X. Li<sup>e</sup>, C. Montaurier<sup>b,c,d</sup>, P. Sirvent<sup>a,b,\*</sup>

<sup>a</sup> EA 3533, Laboratory of Metabolic Adaptations to Exercise in Physiological and Pathological Conditions, Clermont University, Blaise Pascal University, BP 10448, 63000 Clermont-Ferrand, France

<sup>b</sup> Clermont University, Auvergne University, Human Nutrition Unit, BP 10448, 63000 Clermont-Ferrand cedex 1, France

<sup>c</sup> French National Institute for Agricultural Research (INRA), UMR 1019, UNH, CRNH Auvergne, 63000 Clermont-Ferrand, France

<sup>d</sup> UFR Médecine, University Clermont 1, 63001 Clermont-Ferrand, France

<sup>e</sup> Zensun Sci and Tech Ltd., Shanghai, China

Received 27 September 2014; received in revised form 24 November 2014; accepted 2 December 2014

Available online 5 January 2015

## Abstract

**Aim.** – Studies in vitro have highlighted the potential involvement of neuregulin 1 (NRG1) in the regulation of energy metabolism. This effect has also been suggested in vivo, as intracerebroventricular injection of NRG1 reduces food intakes and weight gain in rodents. Thus, it was hypothesised that NRG1 might affect serum leptin levels in mice.

**Methods.** – Weight, food intakes, energy expenditure, spontaneous physical activity and serum leptin levels were evaluated in normal-weight C57BL/6J mice following intraperitoneal administration of NRG1 (50 µg/kg, three times/week) or saline for 8 weeks. Based on the results of this first experiment, leptin-resistant obese db/db mice were then given NRG1 for 8 weeks.

**Results.** – Leptin serum concentrations were six times higher in C57BL/6J mice treated with NRG1 than in the animals given saline. NRG1 treatment also reduced weight gain by 10% and food intakes by 15% compared with saline treatment, while energy expenditure remained unchanged. In db/db mice, serum leptin concentrations, weight gain, food intakes, energy expenditure and spontaneous physical activity were not altered by NRG1 treatment.

**Conclusion.** – The decrease in food intakes and weight gain associated with NRG1 treatment in C57BL/6J mice may be partly explained by increased leptin levels, whereas db/db mice were not affected by the treatment, suggesting resistance to NRG1 in this pathological state.

© 2014 Elsevier Masson SAS. All rights reserved.

**Keywords:** NRG1; Obesity; Energy expenditure; Weight loss

## 1. Introduction

Neuregulin 1 (NRG1) belongs to the epidermal growth factor family. Its biological activity is mediated by erythroblastic leukaemia viral oncogene homologue (ErbB) receptors. NRG1 plays a fundamental role in both developing and mature heart tissue, and NRG1 treatment has positive effects in cardiovascular diseases [1]. Altered NRG1/ErbB signalling has been demonstrated during the development and progression of many tumours [2]. Complex crosstalk between ErbB2 receptors and leptin appears to take place in cancer [3], and ErbB2 induces leptin transcriptional upregulation in human breast epithelial cells [4].

**Abbreviations:** NRG1, Neuregulin 1; ErbB, Erythroblastic leukaemia viral oncogene homologue; RQ, Respiratory quotient; VHL, Vehicle (saline).

\* Corresponding author. Laboratoire des Adaptations Métaboliques à l'Exercice en Conditions Physiologiques et Pathologiques (AME2P), Université Blaise-Pascal, Bâtiment Biologie B, 24, avenue des Landais, BP 80026, 63171 Aubière cedex, France. Tel.: +33 4 73 40 71 33; fax: +00 33 4 73 40 50 62.

E-mail address: [pascal.sirvent@univ-bpclermont.fr](mailto:pascal.sirvent@univ-bpclermont.fr) (P. Sirvent).

<http://dx.doi.org/10.1016/j.diabet.2014.12.002>

1262-3636/© 2014 Elsevier Masson SAS. All rights reserved.

As leptin plays a major role in energy balance regulation, and given the link between leptin and NRG1/ErbB signalling in cancer, it was hypothesized that NRG1 treatment might alter leptin levels.

Therefore, the effect of chronic (8-week) intraperitoneal (i.p.) NRG1 treatment on leptin, weight status, fat mass, food intakes, physical activity and energy expenditure was assessed in normal-weight mice and, based on the results of this first experiment, in leptin-resistant db/db mice also.

## 2. Methods

### 2.1. Animals

A total of 16 C57BL/6JRJ and 16 BKS(D)-*Lepr<sup>db</sup>/JOrIRj* mice (10 weeks old), provided by CERJ Janvier (Le Genest-Saint-Isle, France), were kept in temperature-controlled (20–22 °C) cages (single-housed) with free access to water and food (A04 diet; Scientific Animal Food and Engineering, France). All animal husbandry and experimental procedures were in accordance with the current legislation on animal experimentation, and were approved by the local ethics committee (CEMEA Auvergne, CE 97-12).

### 2.2. NRG1 treatment

Recombinant NRG1 (beta-2a; Zensun Sci and Tech, Shanghai, China) [5] at 50 µg/kg body weight or an equivalent volume of 0.9% NaCl (sodium chloride) was administered by i.p. injection three times a week for 8 weeks ( $n=8$  for each group). Mice were euthanised 3 days after the end of treatment.

### 2.3. Food intakes, body mass and body composition

Body weight and food intakes were recorded every week during the treatment period. One week before sacrifice, body composition was evaluated by quantitative nuclear magnetic resonance imaging (EchoMRI; Echo Medical Systems, Houston, TX, USA).

### 2.4. Daily energy expenditure and physical activity

VO<sub>2</sub>, VCO<sub>2</sub> and physical activity were measured in mice using a PhenoMaster/LabMaster home cage system (TSE Systems, Bad Homburg, Germany). Energy expenditure was calculated using the Weir equation [6] from measurements of gas exchanges computed for each cage from data sampled every 5 min. Respiratory quotient (RQ) was determined from VCO<sub>2</sub>/VO<sub>2</sub>. Spontaneous activity was measured using three-dimensional meshes of light beams. For adaptation, mice were placed in individual calorimetric cages (22 °C) for 24 h prior to data collection. Daily energy expenditure and physical activity were computed for a 24-h period.

### 2.5. Leptin measurement

Serum leptin was evaluated with an ELISA kit (Abcam, Cambridge, MA, USA), and samples were read at 450 nm using a PowerWave spectrophotometer (US BioTek, North Shoreline, WA, USA).

### 2.6. Statistical analysis

SPSS Advanced Statistics software was used for the statistical analysis (IBM, Armonk, NY, USA). Data are presented as the mean ± SEM. One-way analysis of variance (ANOVA) with repeated measures was performed to assess weight and food intake changes following NRG1 or saline treatment. Unpaired Student's *t* test was used where appropriate. Statistical significance was set at  $P<0.05$ .

## 3. Results

### 3.1. NRG1 increases serum leptin and decreases weight gain, food intakes and spontaneous physical activity in C57BL/6JRJ mice

On evaluating NRG1 effects in C57BL/6JRJ mice (Fig. 1), serum leptin was strongly increased compared with the saline vehicle (VHL)-treated animals ( $183.4 \pm 46.6$  pg.mL<sup>-1</sup> vs.  $29.5 \pm 5.1$  pg.mL<sup>-1</sup>, respectively;  $P<0.05$ ; Fig. 1A). The VHL group also gained 9.6% in body mass (from  $25.1 \pm 0.95$  g to  $27.6 \pm 0.85$  g) whereas, in NRG1-treated mice, weight remained stable (from  $26.6 \pm 0.52$  g to  $26.1 \pm 0.43$  g) with treatment (Fig. 1B). Similarly, food intakes increased by 15.3% in the VHL group (from  $27.1 \pm 0.59$  g.week<sup>-1</sup> to  $31.5 \pm 1.5$  g.week<sup>-1</sup>;  $P<0.05$ ; Fig. 1C), but did not change in NRG1-treated mice (from  $29.3 \pm 0.57$  g.week<sup>-1</sup> to  $29.3 \pm 0.69$  g.wk<sup>-1</sup>; Fig. 1C). At the end of the treatment period, body fat percentages were lower in the NRG1 than in the VHL group ( $12.5 \pm 0.67\%$  vs.  $16.3 \pm 0.91\%$ , respectively;  $P<0.05$ ; Fig. 1D). Daily energy expenditure (Fig. 1E) and RQ (not shown) did not differ between the groups. Spontaneous activity over 24 h and at night was lower in the NRG1 than in the VHL group ( $288.2 \pm 31.5$  m vs.  $502.6 \pm 102.3$  m and  $227.7 \pm 24.9$  vs.  $373.6 \pm 56.1$  m, respectively;  $P<0.05$ ), whereas diurnal activity was similar in both groups (Fig. 1F).

### 3.2. NRG1 has no effect on serum leptin, weight gain, food intakes or spontaneous physical activity in db/db mice

As NRG1 treatment increased serum leptin levels in C57BL/6JRJ mice, it was postulated that NRG1 effects might be blunted in leptin-resistant db/db mice. Serum leptin concentrations were comparable in both NRG1- and VHL-treated db/db mice (Fig. 2A). Similarly, weight (Fig. 2B) and food intakes (Fig. 2C) remained stable in both groups of db/db mice, and the percentage of body fat was also comparable in both groups at the end of treatment (Fig. 2D). Energy expenditure (Fig. 2E),

Download English Version:

<https://daneshyari.com/en/article/3259116>

Download Persian Version:

<https://daneshyari.com/article/3259116>

[Daneshyari.com](https://daneshyari.com)