

Applied nutritional investigation

New insights about iron bioavailability inhibition by zinc

Manuel Olivares, M.D.^{a,*}, Fernando Pizarro, M.T.^a, and Manuel Ruz, Ph.D.^{a,b}

^a *Institute of Nutrition and Food Technology, University of Chile, Santiago, Chile*

^b *Department of Nutrition, Faculty of Medicine, University of Chile, Santiago, Chile*

Manuscript received November 20, 2006; January 16, 2007.

Abstract

Objective: We measured the effects of lower and higher doses of zinc (Zn) given as an aqueous solution on the bioavailability of iron (Fe).

Methods: Fourteen healthy subjects received a solution with 0.5 mg of elemental Fe as ferrous sulfate given alone or with 0.59 mg of Zn as zinc sulfate (molar ratio Zn:Fe 1:1). Fourteen days after they received a second solution with 10 mg of Fe given alone or with 11.71 mg of Zn (molar ratio Zn:Fe 1:1). Iron bioavailability was assessed by erythrocyte incorporation of iron radioisotopes ⁵⁵Fe and ⁵⁹Fe.

Results: No significant effect of Zn on Fe bioavailability was observed at lower doses; however, at higher doses Fe bioavailability was inhibited by 56% ($P < 0.001$, repeated measures analysis of variance).

Conclusion: The inhibitory effect of Zn on Fe bioavailability depends on the total amount of both minerals present in the intestinal lumen. This fact should be considered when designing a supplementation program if Fe and Zn are to be provided together. © 2007 Elsevier Inc. All rights reserved.

Keywords:

Iron absorption; Zinc; Iron; Interaction; Human

Introduction

Iron deficiency is the single most common nutritional disorder worldwide and the main cause of anemia in infancy, childhood, and pregnancy [1]. It is prevalent in most of the developing world where it coexists with other micronutrient deficiencies such as zinc, vitamin A, and folate [2,3]. The exact prevalence of zinc deficiency is not known, but it is estimated that the magnitude might not be too different from that for iron [4]. This is probably because the diet of populations in the developing world is based mainly on foodstuffs that not only have low iron and zinc concentrations, but also the bioavailability of these minerals is poor.

Combined supplementation with both micronutrients is one strategy that can be used to improve the iron and zinc status of a population. However, there is concern about the negative interactions between these two minerals. Studies

performed in humans have shown an inhibitory effect of zinc on iron absorption, but it has not been well established whether this interaction depends on the absolute amount of iron and zinc in the supplement and/or on the molar ratio between these two minerals. This information could help design rational guidelines for iron and zinc supplementation programs.

Therefore, the aim of the present study was to measure the effects of lower and higher doses of zinc on iron bioavailability when both are provided as an aqueous solution in a 1:1 molar ratio.

Materials and methods

Subjects

Fourteen healthy women 35 to 44 y of age were selected to participate in the study. None of the women were pregnant before the study, as confirmed by a negative test result for human chorionic gonadotropin in urine, and all were using an intrauterine device as a method of contraception at the time of the study. A written, informed consent was

This research was supported by grant 1040874 from FONDECYT-Chile.

* Corresponding author. Tel.: +56-2-978-1482; fax: +56-2-221-4030.

E-mail address: molivare@inta.cl (M. Olivares).

obtained from all the volunteers before the isotopic studies. The protocol was approved by the ethics committee of the Institute of Nutrition and Food Technology.

Isotopic studies

Iron isotopes (^{59}Fe and ^{55}Fe) of high specific activity were used as tracers (Du Pont de Nemours, Wilmington, DE, USA). Aqueous solutions containing iron alone, as ferrous sulfate, or iron and zinc, as zinc sulfate, were mixed with the isotopes immediately before administration to the subjects. A total amount of 50 mL of the labeled solutions was administered to each subject. The radioisotope dose and protocols had been previously approved by the Chilean Commission on Nuclear Energy.

On day 1 of the study subjects received 0.5 mg of iron labeled with 111 kBq of ^{55}Fe and on day 2 they received 0.59 mg of zinc and 0.5 mg of iron labeled with 37 kBq of ^{59}Fe (molar ratio 1:1 zinc to iron). A venous blood sample was obtained 12 d later (day 14 of the study) to measure the circulating radioactivity and to determine the iron status of the subjects. These samples also provided baseline values of ^{55}Fe and ^{59}Fe radioactivity in red blood cells for the next set of absorption studies. On day 14, subjects were given 10 mg of iron labeled with 111 kBq of ^{55}Fe and on day 15 of the study they received 11.71 mg of zinc and 10 mg of iron labeled with 37 kBq of ^{59}Fe (molar ratio 1:1 zinc to iron). A final venous sample was obtained on day 31 to determine the increase in red blood cell radioactivity.

Blood analyses

Hemoglobin (CELL-DYN 1700, Abbott Diagnostics, Abbott Park, IL, USA), transferrin saturation [5], Zn-protoporphyrin (ZP Hematofluorometer Model 206D, AVIV Biomedical Inc., Lakewood, NJ, USA), and serum ferritin [6] were assessed to evaluate the iron status of the subjects. Iron deficiency was defined if a subject had a normal hemoglobin and at least two positive indicators of deficiency (transferrin saturation <15%, Zn-protoporphyrin >1.24 $\mu\text{mol/L}$ red blood cells, and serum ferritin <12 $\mu\text{g/dL}$) and iron deficiency anemia if a subject had two positive indicators of deficiency and a hemoglobin level <120 g/L.

For the calculation of total radioactivity ingested, radioactivity was counted in sextuplicate from labeled solution aliquots. The measurement of blood radioactivity was performed from duplicate venous samples according to the technique of Eakins and Brown [7]. All samples were analyzed using a liquid scintillation counter (Tri-Carb 1500TR, Packard Instruments Co., Downers Grove, IL, USA) allowing sufficient time to obtain a counting error <3%. Radioactivity from labeled solution aliquots and venous samples were counted simultaneously at the end of the study to avoid an error in the calculation of iron absorption due to decay that had occurred between administration of the isotopes and the absorption measurement 14 d later. In addition, the absorp-

tion of iron administered on days 14 and 15 was corrected for the isotope that had been administered on days 1 and 2 by subtracting the radioactivity of the blood sample of day 14 from red blood cell radioactivity of day 28. The percentages of iron absorption were calculated on the basis of blood volumes estimated for height and weight [8] and assuming an 80% incorporation of the radioisotope into the erythrocyte [9]. This method is reproducible in our laboratory with a coefficient of variation of 5%.

Statistical methods

Because the percentages of iron bioavailability and serum ferritin have skewed distributions, these values were first converted to their logarithms before performing any statistical analyses. The results were retransformed to the antilogarithms to recover the original units and then expressed as geometric means \pm 1 SD ranges. Statistical analyses performed include repeated measures analysis of variance and Scheffé's post hoc tests to establish significant differences in iron absorption (Statistica 4.5 for Windows, StatSoft Inc., Tulsa, OK, USA). All comparisons were done at the 5% level of significance.

Results

The general characteristics and iron statuses of subjects are presented in Table 1. The means \pm SDs for age, height, and weight of the subjects were 39.4 ± 3.4 y, 156 ± 8 cm, and 63.3 ± 10.1 kg, respectively. The means \pm SDs for hemoglobin and zinc protoporphyrin concentrations and transferrin saturation were 130 ± 11 g/L, 1.27 ± 0.44 $\mu\text{mol/L}$ of red blood cells, and $22.7 \pm 13.8\%$, respectively. Furthermore, the geometric mean (1 SD range) for serum ferritin was 22.1 (8.8–55.4). Two of the 14 subjects had iron-deficiency anemia and two had iron deficiency without anemia.

When equimolar concentrations of zinc to iron (1:1) were administered to the subjects, no significant effect of zinc on iron bioavailability was observed at lower doses (0.59 mg of zinc and 0.50 mg of iron; Table 2). However, at higher doses (11.71 mg of zinc and 10 mg of iron) zinc had a 56% inhibitory effect on iron bioavailability (repeated measures analysis of variance, $F = 23.92$, $P < 0.001$; Scheffé's post hoc test, $P < 0.002$).

Discussion

Most of the information on the interaction between non-heme iron and zinc has been obtained from studies that provided these two minerals simultaneously as a supplement. We previously demonstrated that zinc impairs iron bioavailability in a dose-dependent way when both micronutrients are administered together as a water solution in

Download English Version:

<https://daneshyari.com/en/article/3277460>

Download Persian Version:

<https://daneshyari.com/article/3277460>

[Daneshyari.com](https://daneshyari.com)