

## Latest Generation, Wide-Angle, High-Definition Colonoscopes Increase Adenoma Detection Rate

ANDREAS ADLER,\* ALIREZA AMINALAI,<sup>‡</sup> JENS ASCHENBECK,<sup>‡</sup> ROLF DROSSEL,<sup>‡</sup> MICHAEL MAYR,<sup>‡</sup> MATHIAS SCHEEL,<sup>‡</sup> ANDREAS SCHRÖDER,<sup>‡</sup> TIMUR YENERIM,\* BERTRAM WIEDENMANN,\* ULRICH GAUGER,\* STEPHANIE ROLL,<sup>§</sup> and THOMAS RÖSCH<sup>||</sup>

\*Central Interdisciplinary Endoscopy Unit, Department of Gastroenterology, Charité University Medical Hospitals, Berlin; <sup>‡</sup>Private gastroenterology practice, Berlin;

<sup>§</sup>Department of Biostatistics, Charité University Medical Hospitals, Berlin; and <sup>||</sup>Central Research Institute of Ambulatory Health Care and Department of Interdisciplinary Endoscopy, University Hospital Hamburg-Eppendorf, Hamburg, Germany

**BACKGROUND & AIMS:** Improvements to endoscopy imaging technologies might improve detection rates of colorectal cancer and patient outcomes. We compared the accuracy of the latest generation of endoscopes with older generation models in detection of colorectal adenomas. **METHODS:** We compared data from 2 prospective screening colonoscopy studies (the Berlin Colonoscopy Project 6); each study lasted approximately 6 months and included the same 6 colonoscopists, who worked in private practice. Participants in group 1 ( $n = 1256$ ) were all examined by using the latest generation of wide-angle, high-definition colonoscopes that were manufactured by the same company. Individuals in group 2 ( $n = 1400$ ) were examined by endoscopists who used routine equipment (a mixture of endoscopes from different companies; none of those used to examine group 1). The adenoma detection rate was calculated on the basis of the number of all adenomas/number of all patients. **RESULTS:** There were no differences in patient parameters or withdrawal time between groups (8.0 vs 8.2 minutes). The adenoma detection rate was significantly higher in group 1 (0.33) than in group 2 (0.27;  $P = .01$ ); a greater number of patients with least 1 adenoma were identified in group 1 (22.1%) than in group 2 (18.2%;  $P = .01$ ). A higher percentage of high-grade dysplastic adenomas were detected in group 1 (1.19%) than in group 2 (0.57%), but this difference was not statistically significant ( $P = .06$ ). **CONCLUSIONS:** The latest generation of wide-angle, high-definition colonoscopes improves rates of adenoma detection by 22%, compared with mixed, older technology endoscopes used in routine private practice. These findings might affect definitions of quality control parameters for colonoscopy screening for colorectal cancer.

**Keywords:** Adenoma Detection Rate (ADR); Early Detection; Colon Cancer; CRC; Screening Trial; BECOP-6.

Screening colonoscopy has been shown to decrease colorectal cancer (CRC) incidence as well as mortality,<sup>1</sup> not only by finding cancers at an earlier stage but even more because of the detection and removal of adenomas as precursor lesions. Thus, the adenoma detection rate (ADR) has been considered to be one of the main quality outcome parameters of screening colonoscopy.<sup>2</sup> ADR has recently been shown to correlate with colonoscopy withdrawal times.<sup>3</sup> Adenoma rates reported from various countries have ranged from 8%–35%,<sup>4–10</sup> and it is not known whether these differences reflect differences in disease prevalence or perhaps also in colonoscopy quality.

Among the potential factors for improving the quality of colonoscopy and increasing the ADR, the use of new technology has often been advocated, but only a few studies have focused on this issue. Such studies, moreover, have dealt with specific refinements within one endoscope generation such as wide-angle imaging,<sup>11</sup> use of image processing such as narrow band imaging (NBI),<sup>12–15</sup> or Fuji Intelligent Color Enhancement.<sup>16,17</sup> Although these studies could not consistently show differences in outcome, instrument quality in the control groups always represented the most up-to-date standard endoscopes.

Larger randomized comparisons of different generations of instruments, which arrive with a variety of improvements in imaging, image resolution, and image processing, have not been performed yet. In daily routine and especially in private practice, colonoscopy is performed with various generations of instruments, mostly not of the latest generation. However, there are no published data on the question of whether and to what extent introduction of new technology might improve outcome as compared with daily routine. A small recent randomized study found a 3-fold increase in ADR when using the latest generation instrument as compared with the previous endoscope generation.<sup>18</sup>

We therefore compared the performance of the same 6 Berlin colonoscopists who participated in 2 prospective studies. Data from these 6 endoscopists were taken from the large prospective quality assurance study performed by a total of 21 colonoscopists that was examining the influence of case volume on adenoma detection<sup>19</sup> as well as documentation quality of main screening colonoscopy outcome parameters.<sup>20</sup> The other study was a prospective randomized comparison of high-definition television (HDTV) colonoscopies with and without NBI, which were performed by this group of 6 colonoscopists and which did not show any difference in adenoma detection.<sup>15</sup>

### Methods

Two groups of screening colonoscopy patients from 2 successive prospective study periods were analyzed with respect to ADR (all adenomas/all patients). All colonoscopies were performed during the 2 study periods in 5 private gastroenter-

**Abbreviations used in this paper:** ADR, adenoma detection rate; CRC, colorectal cancer; HDTV, high-definition television; HGIN, high-grade intraepithelial neoplasia; NBI, narrow band imaging.

© 2012 by the AGA Institute

1542-3565/\$36.00

doi:10.1016/j.cgh.2011.10.026

ology practices in Berlin by the same 6 colonoscopists, each with outpatient screening colonoscopy volumes of >500 per year; the lifetime experience of the participating examiners was >10,000 colonoscopies each. Both study periods spanned 6–7 months.

### ***Study Group I: Latest and Uniform Technology***

Data for this group was recruited from a prospective randomized study by using exclusively the latest generation Olympus (Tokyo, Japan) colonoscopes to compare conventional wide-angle HDTV colonoscopy with and without NBI, with ADR as the main outcome parameter; instrument, processor, as well as screen were of the same HDTV generation. The comparative results of this study, which did not show any differences in ADR between the 2 groups in any of the parameters analyzed, were reported elsewhere.<sup>15</sup> Because no differences were found in this comparative study between the group in which NBI was used and the group in which HDTV imaging alone was used, data of both the NBI and non-NBI groups are combined for the present comparative analysis. We consider this group to be the one in which the most modern technology, namely high-resolution HDTV imaging, was used.

### ***Study Group II: Older Generation Endoscopes***

Of 18 private practices with 21 physicians in Berlin who performed a quality assessment study of screening colonoscopy and included 12,134 cases during a total of 18 months,<sup>19</sup> colonoscopy data ( $n = 1400$ ) from 6 of 18 colleagues were extracted during the first 6 months of their study participation to arrive at an equal time period compared with the retrospective analysis. In the abovementioned study, various performance parameters, findings, and complications were assessed in this study as well as patient acceptance (evaluated by questionnaire) until study termination. ADR was one of the main quality outcome parameters of this trial. A study audit was performed for all participating centers to achieve high data completeness except for nonconsenting patients. The instruments routinely used in these 5 practices came from different companies but did not represent the latest generation instruments of the respective companies. They were from Pentax (EC-3870, EC-3940; Pentax Europe Co, Hamburg, Germany) in 3 practices and from Fujinon (EC 201 WI, EC 200 MR, EC 250 WI5; Fujinon Europe Co, Willich, Germany) and Olympus (PCF-100 and CF-145; Olympus Europe Co, Hamburg, Germany) in 1 practice each. Respective processors of the same generation were used; screens were not of HD quality.

### ***Data Acquisition and Recorded Parameters***

Ethical approval from the Charité Ethical Committee was obtained for the 2 prospective studies (EA 02/019/07 and EA 02/018/07) that involved either data acquisition/follow-up and/or use of a new scope imaging technology (HDTV/NBI).

The following parameters were recorded: (1) cecal intubation rate; (2) number of adenomas, with location, size, and histology; because of the unclear definition of flat adenomas in the retrospective analysis, this parameter was not included in our 3-group comparison; (3) number of hyperplastic polyps, with location and size; and (4) examination times (introduction and withdrawal).

### ***Outcome Parameters***

The main outcome parameter was the ADR, calculated as number of all adenomas/number of all patients. Secondary outcome parameters were the percentage of patients with at least 1 adenoma, the rate of high-grade dysplastic adenomas as special risk lesions, and the overall number of hyperplastic polyps.

### ***Data Completeness***

In the modern technology study (group I), only 2 instruments were available in the participating offices, leading to inclusion of about 75% of screening colonoscopy cases; subsequent patients were included according to availability of reprocessed scopes without further selection. Group II (extracted from the Berlin screening study) was audited for completeness by comparing included patients with the coded cases during the same period, with missing cases being supplemented as far as possible.

### ***Statistical Analysis***

To test whether percentage differences between the 2 groups were statistically significant,  $\chi^2$  tests were performed. In the case of metric variables (age, etc), analysis of variance for independent groups was carried out, followed by pair-wise comparisons in the case of statistically significant main effects. When only 2 groups were compared,  $t$  tests were performed.

## **Results**

Results concerning patient and examination data are shown in Table 1. There was only a significant difference between the 2 groups with respect to the sedation regimens, with fewer patients sedated and a lower rate of patients sedated with propofol in group I. The other parameters including endoscope withdrawal times were similar in the 2 patient groups. As shown in Table 2, ADRs (all adenomas/all patients) as well as patient rates with at least 1 adenoma were significantly higher in group I compared with group II. The percentage of small adenomas was also significantly different. There were numeric differences with respect to high-grade intraepithelial neoplasia (HGIN) lesions, but they failed to reach statistical significance, probably because of low overall numbers.

## **Discussion**

Our study focuses on a possible effect of new endoscope technology on adenoma detection during screening colonoscopy. We used data from 2 prospective studies that were performed by the same colonoscopists, but with different equipment. In one study, endoscopists used their usual instruments consisting of different generation endoscopes from different companies, exactly mimicking daily routine and reality; none of these scopes were of the latest generation. The other study, which was primarily performed to detect differences between NBI and non-NBI imaging (but adenoma rates were exactly the same in both groups<sup>15</sup>), was used as example of a homogeneous switch to the most modern scope technology, in this case from one company. Both studies were prospective and were audited, and their main outcome parameter was the ADR.

Such a study design, comparison of data of 2 trials, is naturally different from a prospective randomized study com-

Download English Version:

<https://daneshyari.com/en/article/3283640>

Download Persian Version:

<https://daneshyari.com/article/3283640>

[Daneshyari.com](https://daneshyari.com)