



Original Article

National Data of CPR Procedures Performed on Hospitalized Thai Older Population Patients[☆]

Panita Limpawattana^{1*}, Arunotai Siriussawakul², Mann Chandavimol³,
Kittisak Sawanyawisuth¹, Jarin Chindaprasirt¹, Vichai Senthong¹,
Kaewjai Thepsuthammarat⁴

¹ Department of Internal Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen, ² Department of Anesthesiology, Faculty of Medicine, Siriraj Hospital, ³ Division of Cardiology, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, ⁴ Clinical Epidemiology Unit, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand

ARTICLE INFO

Article history:

Received 29 January 2013

Received in revised form

8 November 2013

Accepted 11 April 2014

Available online 4 June 2015

Keywords:

aging,
cardiac arrest,
coronary circulation,
heart arrest,
survival rate

SUMMARY

Background: Little is known about the epidemiology of the elderly who receive cardiopulmonary resuscitation (CPR) and the subsequent survival factors in Thailand and other developing countries. The objective of this study was to determine the rate of CPR and the survival rate among hospitalized Thai elderly patients, and also factors predicting survival at discharge.

Methods: National databases from three sources were searched. These three systems accounted for 96% of the Thai population. All inpatients in the fiscal year 2010 (from October 1, 2009 to September 31, 2010) aged 60 years or over who received CPR procedure were included. Baseline characteristics were studied. The study outcomes were CPR rate, mortality rate, and survival factors.

Results: In total, 17,813 elderly patients who were hospitalized during the 2010 fiscal year received CPR (997.2/100,000 older adults). Of those, 5125 patients (28.77%) survived at discharge. Pre-existing comorbidity, asystole, time on a mechanical ventilator > 96 hours, and being admitted to a private hospital were associated with poorer outcomes ($p < 0.05$).

Conclusion: The rate of CPR in hospitalized elderly patients was 997.2 events/100,000 older adults with a survival rate of 28.77%. Factors predicting death at discharge are pre-existing comorbidities, cardiac arrhythmia type, intervention/procedure, and type of hospital.

Copyright © 2015, Taiwan Society of Geriatric Emergency & Critical Care Medicine. Published by Elsevier Taiwan LLC. All rights reserved.

1. Introduction

Cardiopulmonary resuscitation (CPR) is defined as an attempt to restore spontaneous circulation by performing chest compressions with or without ventilations¹. The survival rate of out-of-hospital CPR appeared to be less than in-hospital CPR². The overall incidence of CPR in the elderly (≥ 65 years) from the largest study to date which included 433,985 patients who underwent in-hospital CPR was 2.73 events per 1000 admissions. The rates were higher

in black and nonwhite patients³. The incidence of CPR in the elderly among developing countries is currently limited.

The outcome of CPR is diverse due to the differences in age and pre-existing conditions⁴. Nearly half of older people have a history of heart failure compared with one-fifth of younger people⁴. Prolonged hypoxemia in survivors worsens functional and cognitive status, which consequently leads to caregiver burden and poor quality of life. Therefore, it is a common dilemma of physicians to weigh up between rational and moral decisions dealing with older persons with cardiac arrest⁴.

There are several predictors of CPR outcomes such as age, sex, ethnic, pre-existing conditions, time to perform CPR, or type of arrhythmia. Aged over 70 years, male African Americans were shown to be associated with poor survival rate and functional status^{2,3,5–8}. Older adults often have multiple comorbidities, and are at risk of death during hospitalization. Advance directive plans

[☆] Conflicts of interest: All contributing authors declare that they have no conflicts of interest.

* Correspondence to: Dr Panita Limpawattana, Division of Geriatric Medicine, Internal Medicine Department, Faculty of Medicine, Khon Kaen University, Friendship Highway Road, Muang District, Khon Kaen, 40002, Thailand.

E-mail address: lpnit@kku.ac.th (P. Limpawattana).

for these patients in the event of a cardiac arrest are challenging for physicians, patients, and their families to prevent unnecessary CPR and improve satisfaction of patient care⁹. Unfortunately, the data regarding the epidemiology and prognostic factors for survival at discharge in an Asian context are limited. This study aims to determine the rate of CPR, its outcome, and survival factors in Thai elderly patients using a national database.

2. Materials and methods

2.1. Patient population

This study was a national survey over three Thailand databases including the National Health Security Office (NHSO), the Civil Servants Benefit System from the Comptroller General's Department, and the Social Security Office. These three systems accounted for 96% of the Thai population. All inpatients in the fiscal year 2010 (from October 1, 2009 to September 31, 2010) aged 60 years or over who received CPR procedure (the International Classifications of Diseases or ICD-9-99.6: cardioversion of cardiac rhythm) were included. Data were checked for accuracy by looking for: (1) overlapping information; (2) visit dates; (3) missing items; (4) incorrect coding; and (5) dating with the correct fiscal year by our analyst team. Data collection of all eligible patients was performed by using their summary discharge forms.

2.2. Patient demographics and clinical characteristics

Baseline characteristics of the patients including age, sex, primary admitting diagnosis, presence of cardiovascular bypass, duration of mechanical ventilator use, levels of hospital, regions of hospital, admission rate, prevalence rate, and mortality rate were studied. For admitting diagnosis, the principal diagnosis of participants was obtained from the ICD-10 codes; severe infection (J09-J22, A49.9, N39), stroke (I61-I67), cirrhosis with complication (K74, I85), renal failure (N17-N19), poisoning (T36-T50), drowning (W65-W74), and pulmonary embolism (I26).

2.3. Outcome measures

The study outcomes were admission rate, prevalence rate, and mortality rate per 100,000 populations in three age groups; 61–70 years, 71–80 years, and over 80 years. Survival to discharge post CPR, length of stay in days, and predictors of discharge status were analyzed.

2.4. Statistical analysis

After the primary analysis, results were sent to 10 medical specialists to check the validity of the information. Upon confirmation of validity, the data were compared to the Ministry of Public Health's Statistics Report 2010 for trend congruence as well as the hospital's mortality reporting for each age and disease group, and to the National Death Registration of the Registry Administration, Ministry of Interior Affairs¹⁰. Data were shown as descriptive statistics. CPR rate was calculated from the number of older adults divided by the total population of older adults who received CPR procedure, and survival rate at discharge was calculated from the number of population in each age group divided by the total number of population in the same age group who received CPR procedure. Predictors of CPR outcomes were analyzed using multiple logistic regressions. For univariate analysis, the Chi-square test or Fisher's exact test was used to examine all categorically associated factors. Factors with a $p < 0.20$ were then entered into a multiple logistic regression model. A p value < 0.05 was considered to indicate statistically significant differences and their 95%

confidence intervals (CI) were reported to consider the strength of association between factors and death at discharge.

Ethics approval was provided from the Ethics Committee of Medicine Faculty, Khon Kaen University, Khon Kaen, Thailand under guidelines of the Helsinki Declaration.

3. Results

In total, 17,813 elderly patients who were hospitalized during the 2010 fiscal year received CPR (997.2/100,000 older adults). Of those, 5125 patients (28.77%) survived at discharge. Age group of 70–79 years had the highest CPR rate at 41.11%, while the survival rates were not different among the three different age groups. Renal failure and severe sepsis were the two most common pre-existing conditions (Table 1). The most common established

Table 1

Percentage of patients who received CPR and survived to hospital discharge, according to patient and hospital characteristics.

	CPR (N = 17,813 patients)	Survival at discharge (N = 5125 patients)
Patient characteristics		
Prevalence rate/100,000 older adults (≥ 60 y)	997.2	286.9
Male (%)	52.8	29.4
Female (%)	47.2	28.1
Age (y)		
• 60–69	6314 (35.4)	1734 (27.5)
• 70–79	7320 (41.1)	2145 (29.3)
• 80+	4179 (23.5)	1246 (28.8)
Primary admitting diagnoses		
• MI	2922	690 (23.6)
• CHF	2811	784 (27.89)
• Severe infection	5477	1375 (25.1)
• Stroke	1271	279 (22)
• Cirrhosis with complication	382	107 (28)
• Renal failure	6482	1548 (23.9)
• Poisoning	94	30 (31.9)
• Drowning	6	0 (0)
• Pulmonary embolism	77	12 (15.6)
Established cardiac arrhythmia		
• AVB	5	4 (80)
• CHB	98	71 (72.4)
• Asystole	422	112 (26.5)
• SVT	132	107 (81.1)
• AF	139	94 (67.6)
• VT \pm VF	106	68 (64.2)
Intervention/procedure		
• CABG	150	87 (58.0)
• Bypass anastomosis	88	47 (53.4)
• Mechanical ventilator > 96 h	13,024	3350 (25.7)
LOS (d)		
• 0–6	12,740	3319 (26.1)
• 7–13	2576	934 (36.3)
• 14–20	1015	371 (36.6)
• 21–27	500	162 (32.4)
• 28+	981	338 (34.5)
Hospital characteristics		
Type of hospital		
• Primary	2160	1120 (51.9)
• Secondary	5298	1270 (24)
• Tertiary	9671	2619 (27.1)
• Private	684	116 (17)
Region		
• Central	6613	1504 (22.7)
• North	3861	1080 (28)
• Northeast	5491	1955 (35.6)
• South	1848	586 (31.7)

Data are presented as n (%) unless otherwise indicated.

AF = atrial fibrillation; AVB = atrioventricular block; CABG = coronary artery bypass graft; CHB = complete heart block; CHF = congestive heart failure; CPR = cardiopulmonary resuscitation; LOS = length of stay; MI = myocardial infarction; SVT = supraventricular tachycardia; VF = ventricular fibrillation; VT = ventricular tachycardia.

Download English Version:

<https://daneshyari.com/en/article/3325056>

Download Persian Version:

<https://daneshyari.com/article/3325056>

[Daneshyari.com](https://daneshyari.com)