



Allergologia et immunopathologia

www.elsevier.es/ai



ORIGINAL ARTICLE

Change of mean platelet volume values in asthmatic children as an inflammatory marker

T. Tuncel*, P. Uysal, A.B. Hocaoglu, D.O. Erge, O. Karaman, N. Uzuner

Dokuz Eylul University Hospital, Department of Pediatrics, Division of Allergy, Balçova, 35340, Izmir, Turkey

Received 11 January 2011; accepted 3 March 2011

Available online 28 May 2011

KEYWORDS

Asthma;
Children;
Inflammation;
Mean platelet
volume;
Platelets

Abstract

Background: Asthma is the most common chronic disease of childhood in industrialised countries. T helper-2 (Th-2) cells, mast cells and eosinophils have a role in inflammation of asthma. Recently it was shown that platelets also play a role in asthma. Mean platelet volume shows platelet size and reflects platelet activation.

Objective: The aim of this retrospective study is to evaluate levels of mean platelet volume in asthmatic patients during asymptomatic periods and exacerbations compared with healthy controls.

Methods: The study consisted of 100 asthmatic patients (male/female: 55/45, mean age: 8.2 ± 3.3) and 49 age and sex matched healthy children as a control group.

Results: Mean platelet volume values of asthmatic patients during asymptomatic period were 7.7 ± 0.8 fL while mean platelet volume values in asthmatics during exacerbation were 7.8 ± 0.9 fL. Comparison of mean platelet volume values of asthmatic patients and healthy controls both in acute asthmatic attack and asymptomatic period showed no difference ($p > 0.05$). Comparison of mean platelet volume values at asthmatic attack and asymptomatic period also had no difference ($p > 0.05$). The presence of atopy, infection, eosinophilia, elevated immunoglobulin E, and severity of acute asthmatic attack did not influence mean platelet volume values.

Conclusion: The results of our study suggest that mean platelet volume values may not be used as a marker in bronchial asthma, although prospective studies with larger number of patients are needed to evaluate the role of mean platelet volume in asthma.

© 2011 SEICAP. Published by Elsevier España, S.L. All rights reserved.

Introduction

Asthma is a worldwide problem with an estimated 300 million affected individuals.¹ The prevalence of asthma is increasing, especially in children and it is the most

* Corresponding author.

E-mail address: tuba.tuncel@deu.edu.tr (T. Tuncel).

Table 1 Clinical characteristics of asthmatic children.

Demographical data	Asthmatic patients (n = 100)
Gender (M/F)	55/45
Age (years)	8.2 ± 3.3 (2.5–16)
Disease duration (years)	2.79 ± 2 (1–15)
Total eosinophil count (/mm ³)	372 ± 395 (0–1800)
Immunoglobulin E level (IU/L)	356 ± 537 (5–2982)
Prick test positivity (+/–)	49/51
Asthma attack (/year)	2.17 ± 1.36(1–7)
Immunotherapy(+/–)	10/90

common chronic disease of childhood in industrialised countries.² Asthma is a chronic inflammatory disorder of the airways which involves several inflammatory cells and multiple mediators resulting in characteristic pathophysiological changes. Inflammatory cells such as mast cells, Th2 cells and eosinophils are defined well in the asthmatic airway. Platelet activation was demonstrated in asthma for a number of years. Clinical evidence has been supplemented with experimental data that platelets are important components of the inflammatory response. Activated platelets secrete multiple inflammatory factors such as chemokines, cytokines and coagulation factors and thus enhance the aggregation, adhesion, and thrombus formation. The induction of inflammatory mediators by activated platelets enhances also the recruitment of leukocytes including eosinophils, monocytes and neutrophils via platelet P-selectin.³ Platelets' volume increases when platelets become activated. Larger platelets contain more dense granules and have higher thrombotic potential and are able to induct the inflammation. Mean platelet volume (MPV) reflects the platelet size. Platelet size is correlated with platelet function and activation. Therefore higher MPV levels predict platelet activity and thus intensity of inflammation.⁴ There is a lot of evidence showing that platelets play a role in both allergic and non-allergic inflammatory conditions. Changes of MPV values have been studied in many chronic inflammatory diseases. Increased MPV values are related to some chronic inflammatory conditions such as familial Mediterranean fever (FMF).^{5,6} To the best of our knowledge, there is no study evaluating the relationship between asthma and MPV. The aim of this study is to investigate whether the inflammatory response affects MPV values in childhood asthma.

Methods

Patients

Case records of the Pediatric Allergy Department of Dokuz Eylul University Hospital for the period January 2007 to June 2009 were screened for patients diagnosed with asthma retrospectively. Only complete blood count (CBC) values obtained from patients both during an asthmatic attack and during the asymptomatic period were included in the study. The demographical data of patients were extracted from

the patient data system. Laboratory data were screened via hospital's computerised patient database. Sex- and age-matched healthy children who visited the "well-child outpatient clinic" for routine controls and who did not have any chronic illnesses constituted the control group. The CBC parameters of control group were also obtained from the same hospital computerised database.

The CBC analyses were performed by the Coulter analyser, which was checked every month by the central laboratory. Blood samples were collected in standard tubes, which contained EDTA. The reference values for MPV ranged between 7.0 and 11.0 fL.

Statistical Analyses

Data were evaluated using the Statistical Package for Social Sciences 13.0 (SPSS for Windows 13.0, Inc., Chicago, IL, USA) and by analysing descriptive statistics (means, standard deviation), comparing the means of quantitative data for dual groups using the Student *t*-test and paired *t*-test, chi square test. Pearson's correlation was used to evaluate the association between MPV and other laboratory values. *p* < 0.05 was considered as significant. Data were expressed as the mean ± standard deviation (SD).

Results

There were 55 boys and 45 girls in the asthmatic group, while the healthy control group was of 22 boys and 27 girls. The mean age of the asthma patients was 8.2 ± 3.3, while that of the controls was 9.1 ± 2.6 years. No significant difference was found between asthmatic patients and the control group in terms of age (*p* = 0.078) and gender (*p* = 0.991). Clinical characteristics of the asthmatic children are shown in Table 1.

Mean platelet volume values of asthmatic children during an asthmatic attack and during the asymptomatic period were compared with MPV values of healthy children. Comparison of MPV values in asthmatic patients and healthy controls both in an acute asthmatic attack (*p* = 0.434) and asymptomatic period had no statistically significant difference (*p* = 0.110). Comparison of MPV values in asthmatic patients in an asthmatic attack and in the asymptomatic period also had no statistically significant difference (*p* = 0.755) (Table 2).

The presence of atopy, infection, eosinophilia, elevated IgE, severity of asthmatic exacerbation did not influence MPV values (*p* > 0.05).

Discussion

This study demonstrated that MPV values in asthmatic children both during an asthmatic attack and during the asymptomatic period had no statistically significant difference compared to the healthy control group. Furthermore, no statistically significant difference was found between mean MPV values in asthma exacerbation and asymptomatic period.

The function of platelets is well known in haemostasis but also platelets are fully functional cells concurrently with

Download English Version:

<https://daneshyari.com/en/article/3339946>

Download Persian Version:

<https://daneshyari.com/article/3339946>

[Daneshyari.com](https://daneshyari.com)