



ORIGINAL ARTICLE

Contact allergy to propolis in beekeepers

K. Münstedt^{a,*}, M. Kalder^b

^aDepartment of Obstetrics and Gynaecology, University Hospital Giessen and Marburg, Justus Liebig University, Klinikstrasse 32, 35385 Giessen, Germany

^bDepartment of Gynaecology and Gynaecological Endocrinology and Oncology, University Hospital Giessen and Marburg, Philipps University, Baldingerstrasse, 35043 Marburg, Germany

Received 13 February 2009; accepted 13 April 2009

Available online 22 October 2009

KEYWORDS

Contact allergy;
Propolis;
Beekeeper;
Beekeeping

Abstract

Background: Allergy to propolis seems to be rare and little is known about it.

Objective: The aim of the study was to survey a subset of affected beekeepers to determine aspects such as time of onset of disease, comorbidity, and possible methods of prevention.

Methods: With the help of two German journals for beekeepers we contacted 41 beekeepers with propolis allergy. They were sent a questionnaire which assessed several aspects of the disease and was based on the current literature.

Results: 70.7% returned our questionnaire and had clear signs of propolis allergy with positive testing by their local allergologists. They reported that allergy had developed after an average of 9.5 years beekeeping. We also found a high prevalence of other allergies (72.4%). Interestingly, there were also systemic reactions to propolis in some beekeepers but not necessarily when using propolis as a medication against other diseases. Beekeepers believed that solvents used to clean the hands could play a role in the development of the disease.

Conclusion: This study provides new insights into allergy to propolis. The hypothesis that solvents used to clean the hands could play a role in the development of the disease should be addressed in future studies.

© 2009 SEICAP. Published by Elsevier España, S.L. All rights reserved.

Introduction

In 1915, the first case of propolis allergy was described in a beekeeper. Today there are many reports on the subject of

propolis allergy in the literature and its incidence may be increasing, because propolis is an ingredient of many 'over the counter' products, including cosmetics, toothpastes and ointments, and it is increasingly used as a dietary supplement. The frequency of propolis allergy in patients with contact allergy from unknown causes is reported to be 1.2 to 6.6% and there is a possible link to pollen allergy^{1,2}. A recent case series on propolis allergy in beekeepers showed that

*Corresponding author.

E-mail address: karsten.muenstedt@gyn.med.uni-giessen.de (K. Münstedt).

beekeepers often do not recognise the problem and continue their hobby without protecting themselves from contact with the substance³. It also found that allergic beekeepers even use propolis as a medication for other disorders and identified the presence of benign lung diseases, allergies other than propolis, and psychological instability as factors associated with the likelihood of suffering from propolis allergy³. Regarding lung diseases, a recent publication showed that topical spread of propolis on nasal mucosa as an adjuvant therapy for asthma for 6 months mimicked lung cancer due to the development of a pulmonary tumour in the left lower lobe with satellite nodules, mediastinal lymphadenopathy, and even an increase in the tumour marker CEA⁴.

Since our earlier study did not specifically focus on propolis allergy we initiated a new study in order to gain a better understanding of the problems associated with it⁵.

Methods

Study questionnaires

As there were no previous studies except our own we developed the "QABH-propolis allergy", which supplemented the previously used Questionnaire for the Assessment of Beekeepers' Health (QABH)^{3,5,6}. The questionnaire was tested in 10 volunteers for intelligibility. A copy of the questionnaire is available from KM.

Subjects

We asked readers of the major beekeeping journals *Die Biene*, *Der Imkerfreund*, *ADIZ*, and the *Deutsches Bienenjournal* with an allergy to propolis to contact us, if they were prepared to complete the questionnaires. A total of 41 beekeepers contacted us. They were sent a questionnaire based on the current literature and which assessed several aspects of the disease. Furthermore, they were asked if they were willing to allow us to contact their treating physician if further clarification was required.

Statistical analysis

SPSS version 14.0 (SPSS, Chicago) was used for data management and statistical analysis. A *p*-value of less than 0.05 was considered significant.

Ethical approval

The study was submitted to the ethics committee of the Justus-Liebig-University Giessen, Germany. The project was approved on September 19th, 2006 (application number 113/06).

Results

We were contacted by 41 beekeepers. Twenty-nine of these (70.7%) returned the questionnaire and had clear signs of propolis allergy. The characteristics of the responding beekeepers are given in the Table 1.

Table 1 Demographic characteristics of the beekeepers with allergy to propolis

Characteristic	Study group (n=29)
Age (years)	
Mean (SD)	58.3 (11.2)
Median	59
Range	36–81
Gender (%)	
Female	10.3
Male	89.7
Time spent as a beekeeper (years)	
Mean (SD)	20.1 (5.2)
Median	20
Range	1–75
Number of bee hives attended	
Mean (SD)	19.2 (28.2)
Median	10
Range	0–150

Beekeepers had suffered from allergy to propolis for 7.4 years on average (SD 6.2; range 1–23 years). However, beekeepers were not confronted with the disease when they first began beekeeping. On average, allergy to propolis started after 9.5 years of beekeeping (SD 9.0; range 0.1–35.0 years). In most cases, the diagnosis was verified by skin tests. Interestingly, one person reported that he had had repeated skin reactions on his hands although the skin test failed to verify the allergy to propolis. In two cases (6.9%) family members were also allergic to propolis. Two beekeepers (6.9%) suffered from atopic dermatitis and 21 (72.4%) were allergic to other substances as well, most frequently to grass pollen (*n* = 6; 20.7%), bee venom (*n* = 4; 13.8%), and perubalsam (*n* = 3; 10.3%). Allergy to bee dust was not found in this group.

Reactions to propolis occurred after various time intervals ranging between 5 min and 48 h (mean 10.9 h, SD 13.9 h, median 5 h). The various types of reactions reported are shown in Figure 1. This shows that allergy to propolis in beekeepers is not only a local problem but may cause systemic reactions as well. These may be less severe (urticaria and difficulty swallowing) or more severe (dyspnoea). The symptoms lasted for different time intervals ranging from 5 h to 20 days (mean 4.6 days, SD 4.7 days, median 3 days). Symptoms in beekeepers worsened over time in seven cases (24.1%), improved in six cases (20.7%) and remained the same in the remaining cases. In four cases (13.8%) symptoms were more severe or less severe in spring after beekeepers had less contact with propolis.

We asked the beekeepers what measures they took in order to prevent the problems of propolis allergy. All of them proved to be effective. The majority always wore rubber gloves (*n* = 12; 41.4%) or leather gloves (*n* = 5; 17.2%), some wore gloves only occasionally (*n* = 10; 34.5%), and two never wore gloves (6.9%). In the latter two cases, sensitivity to propolis was present but not when in contact

Download English Version:

<https://daneshyari.com/en/article/3340125>

Download Persian Version:

<https://daneshyari.com/article/3340125>

[Daneshyari.com](https://daneshyari.com)