



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Brief communication

Quality of life and pain multidimensional aspects in individuals with HTLV-1



**Maíra Carvalho Macêdo^{a,*}, Renata de Sousa Mota^a, Naiane Araújo Patrício^b,
Ana Paula Campos dos Santos^c, Selena Márcia Dubois Mendes^d,
Cristiane Maria Carvalho Costa Dias^b, Abrahão Fontes Baptista^e, Katia Nunes Sá^{c,f}**

^a Escola Bahiana de Medicina e Saúde Pública, Programa de Pós-Graduação em Medicina e Saúde Humana, Salvador, BA, Brazil

^b Escola Bahiana de Medicina e Saúde Pública, Programa de Pós-Graduação em Tecnologias da Saúde, Salvador, BA, Brazil

^c Escola Bahiana de Medicina e Saúde Pública, Salvador, BA, Brazil

^d Universidade Federal da Bahia, Faculdade de Medicina da Bahia, Programa de Pós-Graduação em Medicina e Saúde, Salvador, BA, Brazil

^e Universidade Federal da Bahia, Salvador, BA, Brazil

^f Universidade Católica de Salvador, Salvador, BA, Brazil

ARTICLE INFO

Article history:

Received 9 January 2016

Accepted 27 May 2016

Available online 26 July 2016

Keywords:

Pain

Quality of life

HTLV-1

ABSTRACT

HTLV-1 creates a chronic health condition that involves moderate to severe pain with a negative impact on quality of life (QoL). There is no consensus on which attitudes to pain are more related to the worsening of QoL in HTLV-1 infected patients. The aim of this study was to investigate the correlation between QoL and multidimensional aspects of pain in patients with HTLV-1. A cross-sectional study was conducted in Salvador, Bahia, Brazil. The study included individuals diagnosed with HTLV-1. The Short Form 36 Questionnaire was used to analyze QoL, and the Brief Pain Inventory was used to assess multidimensional aspects of pain. The mean pain intensity was 4.88 ± 3.06 on the visual pain scale, and the average impact on QoL corresponded to a loss of approximately 40%. Moderate to high correlations between pain intensity and all domains of QoL were observed and compared reaction attitudes for general activity, mood, ability to walk, ability to work, relationships, sleep, and ability to enjoy life ($r > 0.40$; $p < 0.05$). Moderate correlations were found between all domains of QoL, pain intensity, and reactive attitudes to pain. The greatest pain intensity impacts involved difficulty to walk and to work, and interpersonal relationships in the emotional aspect of QoL.

© 2016 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The Human T-cell Lymphotropic Virus Type 1 (HTLV-1) is a retrovirus endemic in several regions of the world.¹⁻³ In Brazil, its prevalence is higher in Salvador city, corresponding

to 1.76% of the population.⁴ Only 5% of patients with HTLV-1 develop symptoms associated with such retroviruses. It frequently manifests with spasticity, gait disturbances,

* Corresponding author.

E-mail address: mcarvalhomacedo@gmail.com (M.C. Macêdo).

<http://dx.doi.org/10.1016/j.bjid.2016.05.010>

1413-8670/© 2016 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

weakness and stiffness of the lower limbs, impaired dynamic balance, and pain.^{5,6}

In Salvador, 84.3% of infected individuals experience pain, which is more than twice the prevalence in the general population (41.4%).^{7,8} The pain in these individuals mainly affects the lower back and legs, worsens with long periods in one position and physical effort, and presents daily. The pain, lack of bladder control, and changes in gait patterns lead to limitations that have a high impact on activities of daily living (ADLs), contributing to a reduced quality of life (QoL).^{9,10}

Because HTLV-1 is a neglected health condition, aspects such as the impact on the QoL of infected individuals have been poorly explored, especially using specific tools for delineation of multidimensional aspects of pain and assessment of different domains of QoL. Studies on QoL can guide education and health promotion programs focusing on self-care and the functional independence of individuals affected by HTLV-1.¹¹

As the impact of different sensory and reactive characteristics of pain on QoL of infected individuals remain unclear, this study aimed to correlate the multidimensional aspects of pain and levels of QoL for individuals with HTLV-1.

A cross-sectional observational study was conducted in the Advanced Physical Therapy Clinic (CAFis) of the Bahia School of Medicine and Public Health (EBMSP) in Salvador, Bahia, Brazil. The study included men and women aged 18 and older who were diagnosed with HTLV-1 according to criteria defined by the World Health Organization, classified as defined and probable for HAM/TSP, and able to remain upright without support. Patients additional disorders such as arthritis, orthopedic impairments, and other neurological disorders, or who had difficulty understanding the questionnaires were excluded. The clinical team responsible for the Integrative Care and Research Center selected the participants.

Given the scarcity of studies in the literature that served as the basis for this study, the sample size was calculated based on a correlation coefficient of 0.5. A sample of 43 subjects was estimated with a power of 80% and an alpha of 0.05. This sample was increased by 20% due to the large biological variability in this condition.

Data were collected in a standardized manner after the participants signed an Informed Consent form. The project was approved by the Ethics Committee in Research of the Bahia School of Medicine and Public Health (registration number CAAE 13568213.8.0000.5544).

Social class was defined according to criteria of the Brazilian Association of Survey Companies (ABEP), which divides the

population into six socioeconomic strata namely A, B1 (high), B2, C1 (middle), C2, D, and E (low).

The Brief Pain Inventory (BPI) was used to assess pain intensity and interference.¹² To assess pain intensity we used two of the four sensory questions of this instrument (maximum and average pain intensity in the last 24 h). To assess pain interference we used the seven items of the pain interference section of the BPI (general activity, mood, walking ability, normal work, relations, sleep, and enjoyment of life).

The Short Form 36 (SF-36) Questionnaire was used to assess quality of life according to the following domains: functional capacity, physical aspect, pain, emotional aspect, general health, social aspect, mental health, and vitality.¹³

After assuring data had normal distribution, the Spearman test was used to assess the correlation of each of the two sensory items of the BPI with the seven interference aspects of the BPI and to each domain of the SF-36. Numerical variables were expressed as mean and standard deviation. All analyses were performed using the Statistical Package for Social Sciences (SPSS) version 14.0 for Windows (Chicago, USA).

Of the 56 participants, 66.1% were female. The mean age was 52 ± 11.1 years old, and the body mass index (BMI) was 24.5 ± 4.5 kg/m². There was a higher proportion of non-white individuals (92.9%) and individuals belonging to class C (51.8%). Of these, 42.9% used a device such as a cane or crutch for walking, and 26.8% made systematic use of muscle relaxants for pain relief. The mean disease duration was 10.7 ± 7.9 years. The mean most pronounced pain in the last 24 h was reported to be 4.88 ± 3.06 in intensity, while the mean pain score was reported to be 4.66 ± 3.75 . The sites most affected by pain were the lumbar region, reported by 37 participants (66.1%); the knee, reported by 33 (58.9%); the leg, reported by 18 (32.1%); and the feet, reported by 26 (48.4%).

Table 1 shows how sensory aspects of pain impacted interference aspects. Moderate to high correlations were observed between the level of pain intensity in the last 24 h and the average pain in the last 24 h with all areas of pain interference. A correlation analysis between the most pronounced level of pain and average pain in the last 24 h with the domains of the SF-36 showed that a higher intensity of pain corresponded to worse scores in different domains of the QoL (Table 2). Table 2 shows correlations between the SF-36 domains and the BPI interference of pain items. All but the social aspects of the SF-36 domains were negatively correlated with pain interference.

This study confirmed the hypothesis that pain appears to be moderate in this population; with increased pain intensity,

Table 1 – Correlation between reaction and sensory aspects of pain, according to the Brief Pain Inventory.

Interference pain	Worst level of pain (last 24 h)	p	Average pain (last 24 h)	p
General activity	0.625	<0.001	0.584	<0.001
Humor	0.590	<0.001	0.527	<0.001
Ability to walk	0.422	0.001	0.548	<0.001
Capacity to work	0.561	<0.001	0.620	<0.001
Interpersonal relationships	0.589	<0.001	0.507	<0.001
Sleep	0.413	0.002	0.569	<0.001
Ability to appreciate life	0.478	<0.001	0.479	<0.001

Spearman correlation test, alpha 5%.

Download English Version:

<https://daneshyari.com/en/article/3343672>

Download Persian Version:

<https://daneshyari.com/article/3343672>

[Daneshyari.com](https://daneshyari.com)