



# Active screening and surveillance in the United Kingdom for Middle East respiratory syndrome coronavirus in returning travellers and pilgrims from the Middle East: a prospective descriptive study for the period 2013–2015



Sowsan F. Atabani<sup>a,\*</sup>, Steven Wilson<sup>a</sup>, Clare Overton-Lewis<sup>a</sup>, Judith Workman<sup>a</sup>, I. Michael Kidd<sup>b</sup>, Eskild Petersen<sup>c,d</sup>, Alimuddin Zumla<sup>e</sup>, Erasmus Smit<sup>a</sup>, Husam Osman<sup>a</sup>

<sup>a</sup> Public Health England Birmingham Laboratory, National Infection Service, Heart of England NHS Foundation Trust, Birmingham B9 5SS, UK

<sup>b</sup> Department of Virology, University College London Hospitals NHS Foundation Trust, London, UK

<sup>c</sup> Aarhus University, Aarhus, Denmark

<sup>d</sup> The Royal Hospital, Muscat, Oman

<sup>e</sup> Division of Infection and Immunity, University College London, London, and UK National Institute for Health Research Biomedical Research Centre, UCL Hospitals National Health Service Foundation Trust, London, UK

## ARTICLE INFO

### Article history:

Received 18 April 2016

Accepted 18 April 2016

**Corresponding Editor:** Eskild Petersen, Aarhus, Denmark.

### Keywords:

Surveillance

Middle East respiratory syndrome coronavirus (MERS-CoV)

Respiratory viruses

Mass gatherings

Pilgrimage

## SUMMARY

**Background:** Over 25 000 pilgrims from the UK visit Saudi Arabia every year for the Umrah and Hajj pilgrimages. The recent outbreak of Middle East respiratory syndrome coronavirus (MERS-CoV) in South Korea and the continuing reports of MERS-CoV cases from Saudi Arabia highlight the need for active surveillance for MERS-CoV in returning pilgrims or travellers from the Middle East. Public Health England Birmingham Laboratory (PHEBL) is one of a few selected UK public health laboratories responsible for MERS-CoV screening in travellers returning to the UK from the Middle East who present to hospital with severe respiratory symptoms. The results of the PHEBL MERS-CoV screening and surveillance over the past 3 years is presented.

**Methods:** UK travellers/pilgrims who returned from the Middle East and presented to a hospital with respiratory symptoms were studied over the period February 1, 2013 to December 31, 2015. Patients with respiratory symptoms, who satisfied the Public Health England MERS-CoV case algorithm, were tested for MERS-CoV and other respiratory tract viruses on admission to hospital.

**Results:** Two hundred and two patients suspected of having MERS-CoV were tested. None of them had a laboratory-confirmed MERS-CoV infection. A viral aetiology was detected in half (50.3%) of the cases, with rhinoviruses, influenza A (H1N1 and H3N2), and influenza B being most frequent. Peak testing occurred following the annual Hajj season and in other periods of raised national awareness.

**Conclusions:** Respiratory tract infections in travellers/pilgrims returning to the UK from the Middle East are mainly due to rhinoviruses, influenza A, and influenza B. Whilst MERS-CoV was not detected in the 202 patients studied, heightened awareness of the possibility of MERS-CoV and continuous proactive surveillance are essential to rapidly identify cases of MERS-CoV and other seasonal respiratory tract viruses such as avian influenza, in patients presenting to hospital. Early identification and isolation may prevent outbreaks in nosocomial settings.

© 2016 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The Middle East respiratory syndrome coronavirus (MERS-CoV) was first isolated in a patient with fatal pneumonia and renal

failure in the Kingdom of Saudi Arabia (KSA) in June 2012.<sup>1</sup> A further nine patients in Jordan were detected retrospectively.<sup>2</sup> The disease was named Middle East respiratory syndrome (MERS) and it presents as a clinical spectrum ranging from asymptomatic to a severe fulminant multisystem disease affecting all organs.<sup>3</sup> MERS-CoV has remained on the radar of global public health authorities since its first discovery in 2012 because of recurrent nosocomial and community outbreaks and its association with severe disease

\* Corresponding author.

E-mail address: [Sowsan.Atabani@phe.gov.uk](mailto:Sowsan.Atabani@phe.gov.uk) (S.F. Atabani).

and high mortality rates, especially in patients with co-morbidities. MERS-CoV is endemic throughout the Arabian Peninsula,<sup>4</sup> although Saudi Arabia appears to bear the majority of reported cases. MERS-CoV cases have been reported in travellers to the Middle East returning to Europe, North Africa, North America, and Asia.<sup>3,5–7</sup> Whilst primary transmission of MERS-CoV appears to be from camels, evidence of person-to-person and nosocomial transmission of MERS-CoV is well documented.<sup>8,9</sup>

The largest outbreak of MERS-CoV outside the Middle East occurred in South Korea in 2015 and was attributed to poor hospital infection control measures.<sup>9</sup> This outbreak only serves to highlight the continued threat of this novel virus to global health security and calls for all health systems to have proactive MERS-CoV surveillance and screening systems in place for ill returning travellers from the Middle East.

Three of the four laboratory-confirmed cases of MERS-CoV reported in the UK in 2013 were initially identified by the Public Health England Birmingham Laboratory (PHEBL).<sup>8</sup> The first of these cases was in a traveller returning from KSA following a religious pilgrimage. The UK is home to over two million Muslims, who make up 4.8% of the overall population of England and Wales,<sup>10</sup> and these Muslims will visit Mecca and Medina in the KSA for religious purposes at least once in a lifetime. There are three main religious ceremonies or events that occur in KSA,<sup>11</sup> for which over 25 000 pilgrims from the UK travel to KSA each year: (1) the Hajj, which is the main obligatory pilgrimage and attracts up to three million pilgrims per year, (2) Umrah, which is a mini pilgrimage and is undertaken at any time during the year, and (3) Ramadan, the month of fasting. During these events, mass gatherings of pilgrims group together and simultaneously carry out the various religious rituals over a period of days or weeks. These mass gatherings are known to be associated with the transmission of a range of infectious diseases, with respiratory tract infections being exceedingly common.<sup>11</sup>

Birmingham is the second largest city in the UK, and 21% of its population are Muslims.<sup>10</sup> The city also acts as a hub for travel to and from the Middle East from the surrounding areas. The regional public health laboratory situated in Birmingham provides a 7-day service. It is one of the few laboratories in the UK that has been responsible for enhanced surveillance covering the South West, the Midlands and the North of England, for returning travellers with severe respiratory symptoms that fit the clinical and epidemiological criteria of the Public Health England (PHE) MERS-CoV algorithm.<sup>12</sup> A detailed analysis of the results from this continued surveillance for MERS-CoV and other viral respiratory tract infections was conducted, which included the period over the three Hajj seasons of 2013, 2014, and 2015.

## 2. Materials and methods

### 2.1. Patients

From February 1, 2013 to December 31, 2015, returning travellers presenting to hospitals in the Midlands, South West, and North England with respiratory symptoms were actively investigated. Risk assessments were performed routinely by liaison between the PHE Birmingham consultant virologist and the referring hospital microbiologist, including advice on infection control precautions for healthcare workers, sample preparation, and transport arrangements. Patients were admitted or moved to hospital side rooms and staff instructed to use FFP3 masks and personal protective equipment (PPE) until MERS-CoV had been excluded by laboratory testing. Local health protection teams were routinely involved, both at the time of presentation and when the results of testing became available.

### 2.2. Samples

Both upper respiratory tract (URT) samples, such as nose and throat swabs and nasopharyngeal aspirates, and lower respiratory tract (LRT) samples, such as sputum and bronchoalveolar lavage samples, were tested. The sample type provided was dependent on the severity of infection and urgency of testing. PHE ensured that all samples were transported using category B shipping regulations, in accordance with World Health Organization guidance on the regulation for the transport of infectious agents 2013–14,<sup>13</sup> using a specialized courier with a ‘same day’ delivery service. Microbiology investigations were performed routinely in the local laboratories using PHE guidance on the handling and processing of samples suspected of MERS-CoV.<sup>14</sup> Clotted blood was also requested and serum stored for possible future serological testing.

### 2.3. Respiratory virus screening

Investigation for MERS-CoV RNA was included in the routine panel of respiratory viruses tested for by qualitative real-time reverse transcription PCR (rRT-PCR), as described elsewhere.<sup>15</sup> The routine panel included testing for influenza A (both H1 and H3), influenza B, respiratory syncytial virus (RSV), rhinoviruses, parainfluenza virus types 1–4, human metapneumovirus, and adenoviruses.

### 2.4. Statistical methods

The comparison between the likelihood of virus detection and the destination of travel (KSA vs. Dubai) was performed using Z-ratio proportion analysis. The comparison of the duration of symptoms with the detection of a viral agent was performed using the Mann–Whitney *U*-test.

## 3. Results

### 3.1. Patients

Two hundred and fourteen patients fulfilled the criteria of the PHE case definition algorithm and were tested for MERS-CoV infection. Twelve patients were tested elsewhere due to a 5-month cessation of testing at Birmingham as a result of a PHE policy change; these patients were excluded from the analysis. The remaining 202 patient samples were tested at Birmingham. Patient demographic characteristics are shown in Table 1. The median age of the patients was 54 years (range 4 months to 85 years).

### 3.2. Sample types

All (100%) of the 202 samples were tested for MERS-CoV and other respiratory viruses and the results were reported within 24 h of sample receipt, which corresponded to the defined PHE turnaround time. Amongst the positive samples, 57% were from the LRT. Since one of the criteria for MERS-CoV testing is evidence of LRT involvement, 39 patients had a sample taken from the LRT in addition to the URT. In this subset, 21 (53.8%) had a virus detected

**Table 1**  
Demographic characteristics of 202 patients tested, February 2013 to December 2015

	0–16 years	17–65 years	>65 years	Total (%)
Male	5	84	28	117 (57.9)
Female	5	55	25	85 (42.1)
Total (%)	10 (5.0)	139 (68.8)	53 (26.2)	202 (100)

Download English Version:

<https://daneshyari.com/en/article/3361776>

Download Persian Version:

<https://daneshyari.com/article/3361776>

[Daneshyari.com](https://daneshyari.com)