



Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: [www.elsevier.com/locate/ijid](http://www.elsevier.com/locate/ijid)

# The annual Hajj pilgrimage—minimizing the risk of ill health in pilgrims from Europe and opportunity for driving the best prevention and health promotion guidelines



Shuja Shafi<sup>a,\*</sup>, Osman Dar<sup>b,c</sup>, Mishal Khan<sup>d,e</sup>, Minal Khan<sup>f</sup>, Esam I. Azhar<sup>g</sup>, Brian McCloskey<sup>h</sup>, Alimuddin Zumla<sup>i</sup>, Eskild Petersen<sup>j,k</sup>

<sup>a</sup> Muslim Council of Great Britain, London, UK

<sup>b</sup> Public Health England, London, UK

<sup>c</sup> Chatham House Centre on Global Health Security, Royal Institute of International Affairs, London, UK

<sup>d</sup> London School of Hygiene and Tropical Medicine, London, UK

<sup>e</sup> Saw Swee Hock School of Public Health, National University of Singapore, Singapore

<sup>f</sup> Imperial College London, London, UK

<sup>g</sup> Special Infectious Agents Unit, King Fahd Medical Research Centre, and Medical Laboratory Technology Department, Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah, Saudi Arabia

<sup>h</sup> Global Health Department, Public Health England, London, UK

<sup>i</sup> Division of Infection and Immunity, University College London, and NIHR Biomedical Research Centre, UCL Hospitals NHS Foundation Trust, London, UK

<sup>j</sup> Institute of Clinical Medicine, University of Aarhus, Denmark

<sup>k</sup> The Royal Hospital, Muscat, Oman

## ARTICLE INFO

### Article history:

Received 8 June 2016

Accepted 8 June 2016

### Keywords:

Mass gathering

Hajj

public health

prevention

## SUMMARY

Mass gatherings at religious events can pose major public health challenges, particularly the transmission of infectious diseases. Every year the Kingdom of Saudi Arabia (KSA) hosts the Hajj pilgrimage, the largest gathering held on an annual basis where over 2 million people come to KSA from over 180 countries. Living together in crowded conditions exposes the pilgrims and the local population to a range of infectious diseases. Respiratory and gastrointestinal tract bacterial and viral infections can spread rapidly and affect attendees of mass gatherings. Lethal infectious disease outbreaks were common during Hajj in the 19th and 20th centuries although they have now been controlled to a great extent by the huge investments made by the KSA into public health prevention and surveillance programs. The KSA provides regular updated Hajj travel advice and health regulations through international public health agencies such as the WHO, Public Health England, the Centers for Disease Control and Prevention, and Hajj travel agencies. During the Hajj, an additional 25 000 health workers are deployed; there are eight hospitals in Makkah and Mina complete with state-of-the-art surgical wards and intensive care units made specifically available for pilgrims. All medical facilities offer high quality of care, and services are offered free to Hajj pilgrims to ensure the risks of ill health to all pilgrims and KSA residents are minimal. A summary of the key health issues that arise in pilgrims from Europe during Hajj and of the KSA Hajj guidelines, together with other factors that may play a role in reducing the risks to pilgrims and to wider global health security, is provided herein.

© 2016 Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Mass gathering sporting and religious events pose important public health challenges, including the transmission of infectious diseases, exacerbation of non-communicable diseases, and

disorders related to climate change.<sup>1–5</sup> Every year, the Kingdom of Saudi Arabia (KSA) hosts the Hajj pilgrimage, which is the largest mass gathering in the world held on a recurrent annual basis.<sup>3</sup> The Hajj occurs annually from the eighth to the 12<sup>th</sup> of Dhul al-Hijah, the last (12<sup>th</sup>) month of the Islamic calendar, and two to three million people perform the pilgrimage during this period.<sup>3</sup> A further seven million complete a ‘mini’ pilgrimage, known as Umrah, outside the Hajj period throughout the year.

\* Corresponding author.

E-mail address: [msshafi12@gmail.com](mailto:msshafi12@gmail.com) (S. Shafi).

In light of the huge number of pilgrims from all around the world – of which thousands come from low-income countries with minimal access to healthcare – mixing closely for several days in a difficult terrain, it is remarkable that the majority of pilgrims complete the Hajj without experiencing any major health issues. Infectious disease outbreaks were common during Hajj in the 19<sup>th</sup> and 20<sup>th</sup> centuries and have been controlled to a great extent, although proactive surveillance of the transmission of potential epidemic threats at Hajj is critical to preserving global health security.<sup>3,5</sup> Crush injuries and stampedes, which can pose major risks at mass gatherings, are infrequent during the Hajj relative to its size and logistical complexity. A summary of the key health issues that arise in pilgrims from Europe during Hajj and of the KSA Hajj guidelines, together with other factors that may play a role in reducing the risks to pilgrims and to wider global health security, is provided herein.

## 2. Health risks during the Hajj

### 2.1. Communicable diseases

Respiratory and gastrointestinal tract bacterial and viral infections spread rapidly and affect almost all pilgrims during Hajj.<sup>6–13</sup> Respiratory tract infections – whose spread through coughing and sneezing is exacerbated by the crowded Hajj conditions – include community-acquired pneumonia, influenza, and tuberculosis (TB). While bacterial and viral pneumonia are well-documented causes of hospital admission in pilgrims,<sup>13</sup> quantifying the increase in risk of TB transmission is more challenging owing to the longer time period between infection and the development of symptoms. The elderly and those with comorbid diseases such as diabetes are particularly vulnerable to morbidity from respiratory illnesses.

### 2.2. Gastrointestinal disorders

Acute food poisoning is common during the Hajj and is caused by toxins produced by *Staphylococcus aureus* and *Bacillus cereus*. Gastroenteritis due to *Salmonella* spp and viruses such as rotavirus and norovirus are common during Hajj.<sup>3,12</sup> Factors responsible for increasing the spread of gastrointestinal diseases during Hajj include contamination of food through unhygienic preparation, prolonged storage of food, drinking from contaminated water sources, and a shortage of water for hand washing. The risks of dehydration are heightened when Hajj occurs during summer months, owing to the extremely hot climate in Saudi Arabia.<sup>14</sup>

Other infectious disease risks include meningococcal disease, which caused outbreaks during Hajj in the early 2000s owing to overcrowding and high carrier rates of *Neisseria meningitidis* among pilgrims.<sup>3,15–17</sup> Mosquito species responsible for the transmission of malaria and the arbovirus that causes dengue are present in the KSA, although the country has been classified by the World Health Organization (WHO) as a low, geographically restricted malaria transmission area since 2008.<sup>3</sup>

### 2.3. Non-communicable diseases

Historically, infectious diseases were the largest cause of morbidity and mortality during Hajj, but non-communicable diseases are now a major burden.<sup>3</sup> Many pilgrims both elderly and young have existing non-communicable diseases such as diabetes, hypertension, arthritis, epilepsy, liver and kidney disease, which can be worsened by strenuous Hajj conditions or if regular medications are neglected during the spiritual activities. In addition to cardiovascular disease, heat exhaustion and heatstroke are important causes of death; again health-related morbidity is exacerbated when Hajj occurs during the summer months.

### 2.4. Fire and trauma

The risk of injury from fires has been reduced since tents were replaced with fibreglass and cooking in tents was prohibited following a fire in 1997. However, risks from stampedes and crush injuries remain due to the overcrowding. As the events of the falling cranes and the stampede in the 2015 Hajj illustrate, trauma can be a major cause of injury and death during Hajj. Furthermore, many pilgrims who walk long distances as part of the rituals invariably are injured by motor vehicles.

## 3. Factors reducing health risks (Table 1)

Table 1

### 3.1. Pre-Hajj preparations

The KSA Government employs a well-coordinated, inter-sectoral approach to the planning, communication, public health, and safety issues of the Hajj.<sup>3</sup> Planning for Hajj starts soon after the end of the current Hajj and the KSA Ministry of Hajj and Ministry of Health liaise with the governments of all countries from which pilgrims come to the KSA. Recommendations discussed are pre-travel health regulations and advice about vaccinations, health checks, and specific immunizations at the port of entry. Well before each Hajj, the Saudi Ministry of Health generates the health requirements for the issue of travel visas. It also provides health education campaigns through the provision of reading materials and through travel agents, pilgrim group leaders, websites, and the media, both before and during the Hajj.

The KSA Government also provides regular updated Hajj travel advice and health regulations through international public health agencies such as the WHO, Public Health England, the Centers for Disease Control and Prevention (CDC), and Hajj travel agencies. For the 2015 Hajj, for example, alerts were raised for the prevailing global threats from Middle East respiratory syndrome coronavirus (MERS-CoV), influenza and other respiratory tract viruses, multidrug-resistant TB, multi-antibiotic-resistant bacteria, and Ebola.<sup>18–21,6,22,23</sup>

### 3.2. Measures taken during Hajj

During the Hajj, an additional 25 000 health workers are deployed; there are eight hospitals in Makkah and Mina complete with state-of-the-art surgical wards and intensive care units made specifically available for pilgrims.<sup>3</sup> All medical facilities offer high quality of care, and services are offered free to Hajj pilgrims to ensure the risks of ill health to all pilgrims and KSA residents are minimal.

In terms of preventative measures, in addition to the vaccination requirements described below, measures are put in place to ensure food safety and the Ministry of Health ensures strict enforcement of the regulation that pilgrims are not allowed to bring fresh food or agricultural products into the country. Other coordinated activities include targeted insecticide spraying to control mosquito populations, the distribution of health promotion materials to pilgrims, and electronic surveillance of infectious diseases.<sup>3</sup>

### 3.3. Hajj regulations for incoming pilgrims

The health requirements for pilgrimage to Mecca (Hajj and Umrah) are published annually by the KSA Government. Information for pilgrims is made available on the Saudi Arabia Ministry of Health website.<sup>24–26</sup>

Download English Version:

<https://daneshyari.com/en/article/3361787>

Download Persian Version:

<https://daneshyari.com/article/3361787>

[Daneshyari.com](https://daneshyari.com)