



## Development of real-time PCR assay for specific detection of cowpox virus

Elena V. Gavrilova\*, Dmitrii N. Shcherbakov, Rinat A. Maksyutov, Sergei N. Shchelkunov

SRC VB "Vector", Koltsovo, Novosibirsk Region, 630559, Russia

### ARTICLE INFO

#### Article history:

Received 14 November 2009  
Received in revised form 31 May 2010  
Accepted 2 June 2010

#### Keywords:

Orthopoxvirus  
Cowpox virus  
Real-time PCR  
Detection

### ABSTRACT

**Background:** The number of recorded human cowpox cases are recently increasing. The symptoms caused by cowpox virus (CPXV) in a number of human cases are close to the symptoms characteristic of the orthopoxviral human infections caused by monkeypox or smallpox (variola) viruses. Any rapid and reliable real-time PCR method for distinguishing cowpox from smallpox and monkeypox is yet absent.

**Objectives:** The aim of this study was to develop a quick and reliable real-time TaqMan PCR assay for specific detection of cowpox virus and to determine the sensitivity and specificity of this method.

**Study design:** Based on aligned nucleotide sequences of orthopoxviruses, we found a virus-specific region in the CPXV genome and selected the oligonucleotide primers and hybridization probe within this region. The specificity of the developed method was tested using a panel of various orthopoxvirus (OPV) DNAs. The sensitivity was determined using the recombinant plasmid carrying a fragment of CPXV DNA and genomic DNA of the CPXV strain GRI-90.

**Results:** The analytical specificity of this method was determined using DNAs of 17 strains of four OPV species pathogenic for humans and amounted to 100%. The method allows 6 copies of plasmid DNA and 20 copies of CPXV DNA in the reaction mixture to be detected.

**Conclusion:** A quick and reliable TaqMan PCR assay providing for a highly sensitive and specific detection of CPXV DNA was developed.

© 2010 Elsevier B.V. All rights reserved.

### 1. Background

Human cowpox is a zoonotic disease; the human cowpox cases are mainly recorded in European countries.<sup>1</sup> The natural reservoir of cowpox virus (CPXV) is rodents.<sup>2</sup> The virus is transmitted via a direct contact with infected animal, be it a rodent<sup>3,4</sup> or the animals contacting rodents, such as cows,<sup>5</sup> cats,<sup>6,7</sup> dogs,<sup>8</sup> and others.<sup>9,10</sup> In the majority of cases, the human disease has a benign course with development of local lesions, mainly on hands, forearms, face, and less frequently, other body parts.<sup>11–14</sup> The infection course is accompanied by indisposition and fever; in some cases, generalization of the process is observed.<sup>15</sup> In rare cases, when the disease affects eczema patients, its course can be accompanied by severe complications.<sup>1,4</sup> A similar pattern is observed in immunodeficiencies of various etiologies.<sup>5,16,17</sup> In rare cases, cowpox can have a lethal outcome.<sup>18</sup>

Along with monkeypox (MPXV), variola (VARV), and vaccinia (VACV) viruses pathogenic for humans, CPXV belongs to the mem-

bers of the genus *Orthopoxvirus*. All the representatives of this group are genetically close; their antigenic structures are also very similar.<sup>19–22</sup>

In 1980, the WHO officially declared smallpox eradicated on the globe and recommended to stop the vaccination against this disease; in particular, this was determined by the complications accompanying the vaccination. However, the cessation of vaccination had its negative consequences. The human cohort completely lacking the immunity to VARV and other orthopoxviruses (CPXV and MPXV) increases annually; consequently, the reports on human cases of these orthopoxvirus infections have become numerous.<sup>23–27</sup>

Recently, the number of recorded human cowpox cases in European countries has increased.<sup>18</sup> An unprecedented outbreak of human cowpox occurred in December 2008 to January 2009 in France and Germany<sup>28,29</sup> with 18 cases in Germany and 12 cases in France.<sup>30</sup> This demonstrates the need in special efforts aimed at training of medical staff and search for the antivirals specific for orthopoxviruses.<sup>18</sup> However, the most important problem is to develop a rapid and reliable method for CPXV diagnosis. This is determined by several reasons. A correct diagnosing is necessary for choosing the treatment strategy. Moreover, the quickness of diagnosis is no less important than the accuracy. All the orthopoxviruses pathogenic for humans cause a similar clinical pattern at the early stages; however, they display different contagiousness and pathogenicity. It is most important to obtain the information about

**Abbreviations:** CPXV, cowpox virus; OPV, orthopoxvirus; MPXV, monkeypox virus; VARV, variola virus; VACV, vaccinia virus; ECTV, ectromelia virus; CMLV, camelpox virus; ORF, open reading frame; RFLP, restriction fragment length polymorphism.

\* Corresponding author. Tel.: +7 9231901843; fax: +7 3833367409.

E-mail address: [lgavrilova@yandex.ru](mailto:lgavrilova@yandex.ru) (E.V. Gavrilova).

the infectious agent to undertake the corresponding anti-epidemic measures. Finally, note that numerous rapid and accurate specific diagnostic tools have been developed for VARV and MPXV,<sup>31–34</sup> whereas such method for detection and simultaneous differentiation of CPXV is absent.

Currently, CPXV is diagnosed by serological<sup>35</sup> and biological<sup>6</sup> methods. PCR in combination with RFLP assay<sup>36–38</sup> and sequencing<sup>39</sup> were proposed for CPXV identification. However, both these methods are not free from shortcomings. As has been demonstrated, RFLP assay of DNA of various CPXV strains gives a polymorphic pattern, which hinders interpretation of results.<sup>36</sup> Sequencing gives a precise information; however, this method requires expensive equipment and is inapplicable under field conditions. Of the molecular diagnostic methods, the real-time PCR is becoming ever more popular. This method combines quickness and reliability, which allowed this approach to be successfully applied to VARV and MPXV diagnostics.<sup>34,31</sup> Earlier attempts to use real-time PCR for diagnosing human cowpox gave insufficient specificity only to a level of genus identification.<sup>6,35</sup> In this work, we for the first time propose a species-specific CPXV identification based on real-time PCR assay.

## 2. Objectives

The aim of this study was to develop a quick and reliable real-time TaqMan PCR assay for specific detection of cowpox virus and to determine the sensitivity and specificity of this method.

## 3. Study design

To calculate the oligonucleotide primers and probe, all the currently known 85 complete genome sequences of the orthopoxviruses pathogenic for humans as well as sequences of other members of the genus Orthopoxvirus were aligned. The nucleotide sequences of orthopoxvirus strains were extracted from the GenBank database (<http://www.ncbi.nih.gov>). The nucleotide sequence of ectromelia virus strain Naval was obtained at [www.sanger.ac.uk](http://www.sanger.ac.uk). The nucleotide sequences were aligned using the BioEdit v.7.0 and MUSCLE v.3.6 programs. The region of CPXV GRI-90 ORF D11L, unique for cowpox virus,<sup>40</sup> was chosen as the target for the oligonucleotide fluorescent probe. The criteria for primer selection were a specific amplification of orthopoxvirus DNA independently of the presence of related species and the length of the produced amplicon not exceeding 250 bp. The proposed primer pair was analyzed using the program Oligo v.6.31 and tested for the presence of homology to the nucleotide sequences of other viruses and human genome using the BLAST program. The oligonucleotide primers CPXV\_D11L\_upper (5'-AAAACCTCTCCACTTCCATCTTCT-3'), CPXV\_D11L\_lower (5'-GCATTCAGATACGGATACTGATTC-3') and the probe CPXV\_D11L\_probe (HEX-5'-CCACAATCAGGATCTGTAAAGCGAGC-3'-BHQ1) were synthesized in an ABI 3400 DNA/RNA Synthesizer (DNA sintez, Moscow, Russia).

A recombinant plasmid pCD11 carrying CPXV GRI-90 ORF D11L<sup>40</sup> was constructed. The fragment of CPXV DNA was obtained by PCR using the primer pair 5'-CCCAAGCTTTTATTTTCTAACGAATGTAACGA-3' and 5'-CGCGGATCCGAACGTATCAACATGGTCAAG-3'. For further cloning, the sites for restriction endonucleases HindIII and BamHI were introduced into the primers. The amplification product with a length of 1944 bp was hydrolyzed with the restriction endonucleases HindIII and BamHI and purified with a Gel Extraction Kit (QIAGEN). The resulting DNA fragment was ligated with the plasmid vector pBluescript II SK(+), also hydrolyzed with

**Table 1**

The list of virus strains whose DNA was used in real-time PCR.

Species	Strain	DNA source
Cowpox virus	GRI-90	1
	OPV-Claus	2
	88-Lunge	2
	OPV-90/2	2
	OPV-90/5	2
	OPV-89/3	2
	OPV-89/4	2
Vaccinia virus	LIVP	1
	Elestree 3399	1
	Western Reserve	1
Monkeypox virus	CDC#v79-1-005	3
	CDC#v97-1-004	3
Variola virus	Ngami	1
	65/58	1
	Butler	1
	Congo 9	1
Chickenpox virus	VZV No. 4	4

(1) Virus DNAs were isolated from the strains deposited with the collection of the SRC VB Vector.

(2) Virus DNAs were kindly provided by H. Meyer (Munich, Germany).

(3) Virus DNAs were kindly provided by J. Esposito (Atlanta, United States).

(4) Virus DNA was kindly provided by M.A. Susloparov, SRC VB Vector.

HindIII and BamHI. Competent *Escherichia coli* strain Dh5 $\alpha$ F' cells were transformed with the ligation product and plated onto the solid selective medium ST1 containing 200 mg/ml ampicillin and 40 mg/ml X-gal to select the transformants. The recombinant plasmid pCD11 was isolated using a QIAprep Spin Miniprep Kit (QIAGEN). The presence of the target insert in the selected hybrid plasmids was verified by restriction analysis, PCR assay, and sequencing. The concentration of purified plasmid pCD11 was determined spectrophotometrically in an Ultrospec 3000 pro.

For iQ 5 (BioRad, USA) and Rotor-Gene 6000 (Corbett Research, Australia), PCR analysis with 5'-terminal label cleavage (TaqMan system) was performed in 25  $\mu$ l of the reaction mixture containing 200  $\mu$ M dNTP, 10 mM Tris-HCl, 50 mM KCl, 0.08% Nonidet P40, 5 mM MgCl<sub>2</sub>, 300 nM primers, and 250 nM hybridization probe, 1 U of Taq DNA polymerase (Fermentas, USA), and 1  $\mu$ l of the analyzed DNA solution. For a Real-Time PCR System 7500 (Applied Biosystems, USA), the reaction mixture contained 1  $\times$  TaqMan<sup>®</sup> Buffer A, 200  $\mu$ M dNTP, 5 mM MgCl<sub>2</sub>, 300 nM primers, 250 nM hybridization probe, 0.5 U of AmpliTaq Gold<sup>®</sup> DNA polymerase, and 1  $\mu$ l of the analyzed DNA solution.

PCR with recording of fluorescence intensity in iQ 5 and Rotor-Gene 6000 was conducted according to the following mode: 10 min at 95 °C and 40 cycles of 15 s at 95 °C, 30 s at 58 °C, and 30 s at 60 °C; with Real-Time PCR System 7500: 10 min at 95 °C and 40 cycles of 15 s at 95 °C and 1 min at 58 °C.

The specificity of the assay was assessed using a panel of orthopoxvirus DNAs (Table 1). Chickenpox virus (Table 1) and human (Medigen, Novosibirsk, Russia) DNAs were used as negative controls. Blood and tissue samples of experimental animals infected with CPXV strain GRI-90 were used as clinical specimens.

## 4. Results

The TaqMan assay occupies a top position among the real-time PCR formats. This method utilizes a 5'-exonuclease activity of the Taq polymerase. The fluorescence signal is accumulated due to liberation of the fluorescent label conjugated with oligonucleotide hybridization probe.<sup>41</sup> For the first time, this approach to OPV diagnostics was applied to VARV.<sup>42</sup> Recently, TaqMan assay was used to develop several methods for detecting VARV<sup>32,43</sup> and MPXV<sup>31,33</sup>

Download English Version:

<https://daneshyari.com/en/article/3369490>

Download Persian Version:

<https://daneshyari.com/article/3369490>

[Daneshyari.com](https://daneshyari.com)