

# Osteoarthritis and Cartilage



## Prevalence and distribution of intervertebral disc degeneration over the entire spine in a population-based cohort: the Wakayama Spine Study



M. Teraguchi †, N. Yoshimura ‡, H. Hashizume †\*, S. Muraki §, H. Yamada †, A. Minamide †, H. Oka ‡, Y. Ishimoto †, K. Nagata †, R. Kagotani †, N. Takiguchi †, T. Akune §, H. Kawaguchi ||, K. Nakamura ¶, M. Yoshida †

† Department of Orthopaedic Surgery, Wakayama Medical University, Wakayama, Japan

‡ Department of Joint Disease Research, 22nd Century Medical & Research Center, Faculty of Medicine, The University of Tokyo, Tokyo, Japan

§ Department of Clinical Motor System Medicine, 22nd Century Medical & Research Center, Faculty of Medicine, The University of Tokyo, Tokyo, Japan

|| Department of Sensory & Motor System Medicine, Faculty of Medicine, The University of Tokyo, Tokyo, Japan

¶ Rehabilitation Services Bureau, National Rehabilitation Center for Persons with Disabilities, Saitama, Japan

### ARTICLE INFO

#### Article history:

Received 23 May 2013

Accepted 29 October 2013

#### Keywords:

Magnetic resonance imaging  
Population-based cohort  
Intervertebral disc degeneration  
Entire spine  
Prevalence  
Distribution

### SUMMARY

**Objectives:** The purposes of this study were to investigate the prevalence and distribution of intervertebral disc degeneration (DD) over the entire spine using magnetic resonance imaging (MRI), and to examine the factors and symptoms potentially associated with DD.

**Design:** This study included 975 participants (324 men, mean age of 67.2 years; 651 women, mean age of 66.0 years) with an age range of 21–97 years in the Wakayama Spine Study. DD on MRI was classified into Pfirrmann's system (grades 4 and 5 indicating DD). We assessed the prevalence of DD at each level in the cervical, thoracic, and lumbar regions and the entire spine, and examined DD-associated factors and symptoms.

**Results:** The prevalence of DD over the entire spine was 71% in men and 77% in women aged <50 years, and >90% in both men and women aged >50 years. The prevalence of an intervertebral space with DD was highest at C5/6 (men: 51.5%, women: 46%), T6/7 (men: 32.4%, women: 37.7%), and L4/5 (men: 69.1%, women: 75.8%). Age and obesity were associated with the presence of DD in all regions. Low back pain was associated with the presence of DD in the lumbar region.

**Conclusion:** The current study established the baseline data of DD over the entire spine in a large population of elderly individuals. These data provide the foundation for elucidating the causes and mechanisms of DD.

© 2013 Osteoarthritis Research Society International. Published by Elsevier Ltd. All rights reserved.

### Introduction

Intervertebral disc degeneration (DD) is thought to be the first step in degenerative spinal changes<sup>1</sup>, and is typically followed by the gradual formation of osteophytes, disc narrowing, and spinal stenosis<sup>2,3</sup>. Furthermore, DD is considered to be one of the causes of several symptoms (neck pain or low back pain)<sup>4–7</sup>. Therefore, in terms of developing preventive strategies for spinal disorders, it will be important to obtain fundamental data on DD (prevalence, distribution, associated factors, etc.) in a population-based cohort.

We believe that the analysis of DD over the entire spine would provide more useful data than that of DD in the cervical, thoracic, or lumbar regions, separately. In particular, investigations on the extent of DD in these three regions using whole spine magnetic resonance imaging (MRI) could provide useful data concerning intra-individual factors in the development of DD. Several studies have examined degenerative changes in only cervical and lumbar discs because of the high susceptibility to DD in these regions<sup>8–12</sup>. As well, several previous studies have investigated the aging process of the intervertebral discs in the cervical and lumbar regions using MRI in population-based cohorts<sup>13,14</sup>. However, degenerative changes in the thoracic region and correspondingly over the entire spine are poorly understood, because DD in the thoracic region is considered to be an uncommon problem<sup>15,16</sup>. In particular, the stabilization of the thoracic region by the thoracic cage, which

\* Address correspondence and reprint requests to: H. Hashizume, Wakayama Medical University, 811-1 Kimiidera, Wakayama City, Wakayama 641-8510, Japan. Tel: 81-73-447-2300; Fax: 81-73-448-3008.

E-mail address: hashizum@wakayama-med.ac.jp (H. Hashizume).

reduces the mechanical stress imposed on the intervertebral discs, is believed to reduce the incidence of degenerative diseases in this region<sup>17</sup>.

Consistent with the above-mentioned previous studies, a population-based cohort analysis of DD in the different spinal regions using MRI could be used to examine the distribution of DD over the entire spine. However, to our knowledge, no previous studies have performed this type of investigation with a population-based cohort.

From the perspective of discogenic pain, the association between DD and symptoms remains controversial, although several reports have found that DD was a source of low back pain<sup>4,5</sup>. Moreover, reports on the association between the presence of DD in the cervical and thoracic regions and neck pain are rare<sup>6,7</sup>. Further, these studies were not performed with population-based cohorts and did not use whole spine MRI. Thus, no study has assessed neck pain and low back pain within individuals using whole spine MRI. To clarify the points described above, we established a population-based cohort study in which participants underwent whole spine MRI and were examined for symptoms associated with spinal disorders. This is our first report of DD over the entire spine based on a cross-sectional examination of a baseline population.

The aims of this study were to examine (1) the prevalence and distribution of DD over the entire spine using MRI in a population-based cohort, (2) the factors associated with DD (age, gender, and body mass index [BMI]) in the cervical, thoracic, and lumbar regions, and (3) the association between DD and symptoms (neck pain and low back pain).

## Methods

### Participants

The present study, entitled the Wakayama Spine Study, was performed with a sub-cohort of the second visit of the ROAD (Research on Osteoarthritis/osteoporosis Against Disability) study, which was initiated as a nationwide, prospective study of bone and joint diseases in population-based cohorts; the cohorts were established in three communities with different characteristics (i.e., urban, mountainous, and coastal regions) in Japan. A detailed profile of the ROAD study has already been described elsewhere<sup>18,19</sup>. Here, we briefly summarize the profile of the present study. The second visit of the ROAD study began in 2008 and was completed in 2010. All the participants in the baseline study were invited to participate in the second visit. In addition to the former participants, inhabitants aged 60 years and older in the urban area and those aged 40 years and younger in the mountainous and coastal areas who were willing to participate in the ROAD survey were also included in the second visit (both the mountainous and coastal areas were in Wakayama prefecture). Finally, 2674 individuals (900 men, 1774 women) participated in the second visit of the ROAD study, and comprised 1067 individuals (353 men, 714 women) in the urban area, 742 individuals (265 men, 477 women) in the mountainous area, and 865 individuals (282 men, 583 women) in the coastal area. Among these three communities in the ROAD study, the mountainous and coastal areas from which we invited all 1607 participants (547 men, 1060 women) to the Wakayama Spine Study are located in Wakayama prefecture. Of the 1607 participants, a total of 1011 individuals provided written informed consent and attended the Wakayama Spine Study with MRI examinations<sup>20,21</sup>. Among the 1011 participants, those who had MRI-sensitive implanted devices (e.g., pacemakers) and other disqualifiers were excluded. Consequently, 980 individuals underwent MRI of the whole spine. Furthermore, one participant who had undergone a previous cervical operation and four participants

who had undergone a previous posterior lumbar fusion were excluded from the analysis. Finally, whole spine MRI results were available for 975 participants (324 men, 651 women) with an age range of 21–97 years (mean, 67.2 years for men and 66.0 years for women). Table 1 shows the demographic and baseline characteristics of the 975 participants in the present study.

For the purpose of analysis, the participants were divided into five age groups: (1) under 50 years, (2) 50–59 years, (3) 60–69 years, (4) 70–79 years, and (5) 80 years and over. The anthropometric measurements included height, weight, and BMI (weight [kg]/height<sup>2</sup> [m<sup>2</sup>]). BMI was categorized according to the guidelines for Asians proposed by the World Health Organization and was thus defined as follows: underweight, less than 18.5; normal, 18.5–23; overweight, 23–27.5; and obesity, greater than 27.5<sup>22</sup>. Experienced orthopedists also asked all participants the following question regarding neck pain and low back pain: “Have you experienced neck pain on most days during the past month, in addition to now?” and “Have you experienced low back pain on most days during the past month, in addition to now?” Those who answered “yes” were defined as having neck pain or low back pain based on previous studies<sup>23–26</sup>.

### MRI

A mobile MRI unit (Excelart 1.5 T, Toshiba, Tokyo, Japan) was used in the present study, and whole spine MRI was performed for all participants on the same day as the examination. The participants were supine during the MRI, and those with rounded backs used triangular pillows under their head and knees. The imaging protocol included sagittal T2-weighted fast spin echo (FSE) (repetition time [TR]: 4000 ms/echo, echo time [TE]: 120 ms, field of view [FOV]: 300 × 320 mm), and axial T2-weighted FSE (TR: 4000 ms/echo, TE: 120 ms, FOV: 180 × 180 mm).

Sagittal T2-weighted images were used to assess the intervertebral space from C2/3 to L5/S1. C2/3 to C7/T1, T1/2 to T12/L1, and L1/2 to L5/S1 were defined as the cervical region, thoracic region, and lumbar region, respectively. DD grading was performed by an

**Table 1**  
Characteristics of participants

	Overall	Men	Women
<b>No. of participants</b>	<b>975</b>	<b>324</b>	<b>651</b>
<b>Age strata (years)</b>			
<50	125	38	87
50–59	175	59	116
60–69	223	65	158
70–79	261	89	172
≥80	191	73	118
<b>Demographic characteristics</b>			
Age, years	66.4 ± 13.5	67.2 ± 13.9	66.0 ± 13.4
Height, cm	156.4 ± 9.4	164.6 ± 7.2	151.5 ± 7.2
Weight, kg	56.8 ± 11.5	64.5 ± 11.6	53.0 ± 9.4
BMI (kg/m <sup>2</sup> )	23.3 ± 3.6	23.6 ± 3.4	23.1 ± 3.7
<b>BMI (WHO-Asian category) (N)</b>			
Underweight	61	16	45
Normal	425	124	300
Overweight	361	139	221
Obesity	128	44	84
<b>Baseline characteristics</b>			
<b>Symptoms (%)</b>			
Neck pain	24.9	19.4	27.7
Low back pain	43	36.7	42.1
<b>Life style (%)</b>			
Smoking	10.7	25.2	4.1
Alcohol consumption	31.4	56.8	18.8

BMI category for Asian was based on World Health Organization (WHO) guidelines defining underweight (<18.5), normal (18.5–23), overweight (23–27.5), and obese (>27.5). Values are the means ± standard deviation.

Download English Version:

<https://daneshyari.com/en/article/3379401>

Download Persian Version:

<https://daneshyari.com/article/3379401>

[Daneshyari.com](https://daneshyari.com)