

A family cluster of giardiasis with variable treatment responses: refractory giardiasis in a family after a trip to India

A. Requena-Méndez¹, P. Goñi², S. Lóbez², I. Oliveira¹, E. Aldasoro¹, M.-E. Valls³, A. Clavel², J. Gascón¹ and J. Muñoz¹

1) Barcelona Centre for International Health Research (CRESIB Hospital Clínic-Universitat de Barcelona), Barcelona, 2) Area of Parasitology, Department of Microbiology, Preventive Medicine and Public Health, Faculty of Medicine, University of Zaragoza, Zaragoza and 3) Department of Microbiology, Hospital Clínic, Barcelona, Spain

Abstract

Persistence of giardiasis after some of the recommended drugs is occurring with increasing frequency. We describe the follow-up of four members of a family with giardiasis through microscopic observation, immunochromatography and PCRs of *tpi* and *β-giardin* genes. Three patients did not respond to tinidazole but they were cured after quinacrine. However, PCR became negative at 2 months after negativization of stools in two patients and after 1 year in one patient. In all cases *Giardia* assemblage B was characterized with high homology between all isolates. Further studies are needed to determine the value of PCR in the diagnosis of *Giardia* infections.

Keywords: *Giardia*, nitroimidazole, polymerase chain reaction, quinacrine, refractory giardiasis

Original Submission: 26 April 2013; **Revised Submission:** 27 June 2013; **Accepted:** 3 July 2013

Editor: E. Bottieau

Article published online: 9 July 2013

Clin Microbiol Infect 2014; **20**: O135–O138
10.1111/1469-0691.12327

Corresponding author: A. Requena-Méndez, Barcelona Centre for International Health Research (CRESIB, Hospital Clínic-Universitat de Barcelona), Carrer Roselló 132, 4^a, 08036 Barcelona, Spain
E-mail: ana.requena@cresib.cat

Giardia duodenalis is a common parasitic infection in returning travellers [1]. Nitroimidazoles appear to be the drug of choice [2] but, persistence of the infection in humans after treatment is not uncommon [3]. Resistance of *G. duodenalis* to nitroimidazoles has been demonstrated or induced *in vitro*, although its contribution to the persistence of the parasite after treat-

ment remains unclear [2,4,5]. In addition, it is known that *G. duodenalis* infection can be caused by mixed genotypes [6] and as a result antiparasitic treatment could select resistant strains. Finally, in patients who fail to respond to treatment, a re-infection should also be considered, particularly in areas of high endemicity.

In India, *Giardia* infection with diarrhoea is highly prevalent, ranging from 0.4% to 70%, and asymptomatic cyst passage has been found to be as high as 50% in rural southern areas [7].

We report the case follow-up of a family of four members including the father (59 years), the mother (57 years), son 1 (16 years) and son 2 (14 years) that presented with giardiasis after a trip to India in July 2011. All presented watery diarrhoea, abdominal pain and aerophagia. Stool samples from each one were collected before and after each treatment.

Samples were analysed by microscopy following standard formalin–ether concentration, and by rapid immunochromatography (Stick *Cryptosporidium/Giardia/Entamoeba histolytica*; Operon, Zaragoza, SA, Spain). DNA from all the samples was extracted using a DNA stool kit (IBIAN[®] DNA Stool Kit; IBIAN Technologies, Zaragoza, Spain) and stored at –20°C until processing. *Giardia duodenalis* assemblage was determined by a PCR of the triosephosphate isomerase (*tpi*) gene [8]. A fragment-nested PCR was also used to amplify a fragment of the *β-giardin* (*bg*) gene according to Lalle *et al.* [9]. Additionally, PCR products were purified with a GFX[™] PCR DNA Gel Band Purification Kit (GE Healthcare, Chalfont St Giles, UK) and then directly sequenced in both directions. The nucleotide sequences obtained were analysed using the BioEDIT alignment program (<http://www.mbio.ncsu.edu/bioedit/bioedit>) and compared with randomly obtained assemblage B sequences deposited in GenBank.

The DNA sequences obtained have been deposited in the Genetic sequence database (GenBank) at the National Center for Biotechnical Information under accession numbers JQ782391–JQ782407.

All four patients presented *G. duodenalis* cysts in the microscopic examination of stool samples and received a single dose of 2 g of tinidazole. One of the sons improved significantly—the stool examination and the immunochromatography fecal test were both negative after 1 month. The others remained symptomatic after treatment and a new microscopic examination of stools revealed persistence of *G. duodenalis* cysts. In all three cases, HIV test was negative and the presence of IgA deficiency was ruled out. The three relatives with persistence of symptoms received treatment with quinacrine (100 mg three times a day for 5 days) and showed progressive improvement of symptoms. The control stool examinations of the three patients did not show any

Giardia cyst in any repeatedly control samples but the mother's and father's samples showed *Entamoeba histolytica/dispar* cysts and treatment with paromomycin (500 mg three times daily for 7 days) was prescribed. (Fig. 1 shows the test results and the treatment given to each member of the family).

A total of 18 samples (four from both children and five from the parents) were analysed (see Table 1). Of these, 12 yielded positive amplification for a 140-bp fragment of the *tpi* gene corresponding to Assemblage B whereas none were amplified for Assemblage A. Samples from three family members gave positive amplification for *tpi* gene up to 2 months after the negativization of the stool examination by microscopy but they were negative 1 year after the therapy in the two samples that were analysed. All *tpi* sequences were almost identical, with

similarities ranging between 99.2% and 100%. Only five samples gave positive amplification for *bg* gene. With this PCR, the persistence of *Giardia* DNA in the last sample from the father was confirmed and positive amplification was also obtained for the 3-month sample of the asymptomatic son and for the other son after quinacrine treatment. Analysis of the sequences showed a range of similarities between 97.8% and 99.1%. Similarity between β -giardin proteins ranged from 96.4% to 100%. Comparing them with the sequences registered in GenBank, all the β -giardin sequences of this work showed high identity with that of the isolate BAH8 (AY072727), with only one to three single nucleotide polymorphisms, indicating that the present isolates are classified into sub-assemblage BIII.




FIG. 1. Clinical management of cases with imported giardiasis. Samples numbered as 0, 1, 2, 3, 4 correspond to 0: before treatment, 1, 2, 3 and 4: month 1, 2, 3 and 12 after treatment); +: Stool examination positive for cyst of *Giardia duodenalis*; -: Stool examination negative for cysts of *G. duodenalis*.

Download English Version:

<https://daneshyari.com/en/article/3396587>

Download Persian Version:

<https://daneshyari.com/article/3396587>

[Daneshyari.com](https://daneshyari.com)