



The Egyptian Society of Chest Diseases and Tuberculosis
Egyptian Journal of Chest Diseases and Tuberculosis

www.elsevier.com/locate/ejcdt
www.sciencedirect.com



ORIGINAL ARTICLE

The effect of adding long acting beta 2 agonists to inhaled corticosteroids versus increasing dose of inhaled corticosteroids in improving asthma control



Nisreen M. Nabil ^{a,*}, Assem F. Elessawy ^b, Khaled M. Hosny ^c,
Shaaban M. Ramadan ^d

^a Clinical Pharmacy Department, Faculty of Pharmacy, Beni-Suef University, Beni-Suef, Egypt

^b Chest Department, Faculty of Medicine, Fayoum University, Fayoum, Egypt

^c Pharmaceutics and Industrial Pharmacy Department, Faculty of Pharmacy, Beni-Suef University, Beni-Suef, Egypt

^d Chest Department, Faculty of Medicine, Beni-Suef University, Beni-Suef, Egypt

Received 3 July 2014; accepted 8 July 2014

Available online 30 July 2014

KEYWORDS

Bronchial asthma;
Inhaled-corticosteroids;
Long-acting- β_2 -agonist;
Formetrol/budesonide combination therapy;
Asthma control

Abstract To asthmatic patients with moderate to severe persistent asthma, two main treatment options are recommended: The combination of a long-acting inhaled β_2 -agonist with inhaled corticosteroids or the use of a higher dose of inhaled corticosteroids. The aim of this study was to evaluate which drug option is more favorable.

Patients and methods: This study included 60 asthmatic patients uncontrolled on low doses of ICSs. They were randomized into two groups. Group (1): 30 patients received twice daily inhaled formetrol and budesonide in a dose of 12 mcg and 400 mcg, respectively. Group (2): 30 patients received two fold the previous dose of budesonide 800 mcg/BID alone. A comparative study was carried out at Outpatient Chest Clinic of Fayoum Hospital University for a period of 24 weeks using the spirometric data of patients of the two groups before and after treatment.

Results: Results showed that the combination therapy of inhaled formetrol and budesonide is modestly more effective in the reduction of symptoms and in improving the lung functions than with a higher dose of budesonide alone.

Conclusion: Adding formetrol in a dose of 12 μ g plus budesonide in a dose 400 μ g b.i.d. is more favorable in treatment of asthma than a higher dose of budesonide (800 μ g b.i.d).

© 2014 The Egyptian Society of Chest Diseases and Tuberculosis. Production and hosting by Elsevier B.V. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

* Corresponding author.

E-mail address: nesrin_nabil@hotmail.com (N.M. Nabil).

Peer review under responsibility of The Egyptian Society of Chest Diseases and Tuberculosis.

<http://dx.doi.org/10.1016/j.ejcdt.2014.07.007>

0422-7638 © 2014 The Egyptian Society of Chest Diseases and Tuberculosis. Production and hosting by Elsevier B.V.

Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Introduction

Asthma is a chronic inflammatory disorder of the airways in which many cells and cellular elements play a role. The chronic inflammation is associated with airway hyperresponsiveness that leads to recurrent episodes of wheezing, breathlessness, chest tightness and coughing, particularly at night or in early morning. These episodes are usually associated with wide spread, but variable, airflow obstruction within the lung that is often reversible either spontaneously or with treatment [5].

A number of factors that influence a person's risk of developing asthma have been identified. These can be divided into host factors (primarily genetic) and environmental factors [3]. Measurements of lung function (spirometry) provide an assessment of the severity of airflow limitation, its reversibility, and its variability provide confirmation of the diagnosis of asthma [1].

The aim of asthma treatment is to achieve and maintain clinical control for prolonged periods [2].

Medications to treat asthma can be classified as controllers or relievers. Controllers are medications taken daily on a long-term basis to keep asthma under clinical control chiefly through their anti-inflammatory effects.

Relievers are medications used on an as-needed basis that act quickly to reverse bronchoconstriction and relieve its symptoms [5].

To reach clinical control, add on therapy of inhaled LABA to ICS is preferred over increasing the dose of inhaled glucocorticosteroids. Long-acting inhaled β_2 -agonist, including formoterol should not be used as monotherapy in asthma as these medications do not appear to influence airway inflammation in asthma. The addition of the long-acting inhaled β_2 -agonist to the daily regimen of inhaled glucocorticosteroids improves symptom scores, decreases nocturnal asthma symptoms, improves lung function, decreases the use of rapid-acting inhaled β_2 -agonist [10] and achieves clinical control of asthma in more patients, more rapidly, and at a lower dose of ICS than ICS alone [2].

The study was conducted on asthmatic patients among those attending the outpatient chest clinic in Fayoum Hospital University during the period from January 2010 till December 2012.

The study included 60 patients with moderate to severe persistent asthma, uncontrolled on low doses of ICSs (budesonide or beclomethasone dry powder inhaler, 400 mcg/day).

All of these patients were > 18 years old.

Exclusion criteria

Patients using systemic corticosteroids, those having respiratory infections affecting asthma control within the previous 4 weeks, patients with severe cardio-pulmonary disease or other concomitant disease and *smoking patients* were excluded. The patients included were randomized into two groups:

- *Group 1*: 30 patients received twice daily inhaled formoterol and budesonide in the form of dry powder inhaler (aerolizer) in a dose of 12 mcg and 400 mcg,
- *Group 2*: 30 patients received two fold the previous dose of budesonide which is the maximum in the form of dry powder inhaler (aerolizer) in a dose of 800 mcg/BID.

Treatment with systemic anti-histaminic or other anti asthma products will not be permitted.

These individuals were subjected to the following:

1. Full history taking with proper analysis of the complaint and full history of atopic state.
2. Full clinical examination: general and local.
3. Chest X-ray.
4. Routine laboratory investigation.
5. Pulmonary function tests: The patients were instructed to stop medications 24 h before performing the test to avoid faulty results. The spirometric parameters estimated by using the Compact Vitalograph were:
 - a. Forced vital capacity (FVC).
 - b. Forced expiratory volume in the first second (FEV₁).
 - c. FEV₁/FVC ratio.
 - d. Peak expiratory flow rate (PEFR). The test was repeated 3 times and the highest result was recorded.
6. Reversibility testing using repeated nebulization by 5 mg salbutamol and 500 mcg of ipratropium.

Any patient < 12% improvement in FEV₁ was considered COPD patient and excluded.

The patients were started on predetermined therapy for a period of 6 months during which they were weekly supervised in the out-patient Chest Clinic at Fayoum Hospital University. At these visits, the patient's questions were discussed, and any problems with asthma and its treatment are reviewed. Checking of the inhaler device technique with correction and re-checking if it is inadequate were done until the period of treatment ended.

At the end of the test pulmonary function tests (spirometry) were repeated by comparing their results with those of the tests previously done before the therapy.

Adherence to therapy was assessed by reviewing patient's questionnaire which was done at the clinic before initiating the treatment and by the end of the predetermined period of treatment.

At the end of the period of the study all the data were collected, tabulated and statistically analyzed.

Statistical analysis

Statistical analysis for the obtained data was carried out to determine the effect of adding inhaled formoterol to inhaled budesonide compared to a higher dose of inhaled budesonide in improving asthma control using descriptive analysis of the results, Cross tabulation test, Student's *t*-test, Fisher's exact test, and Paired samples *t*-test. Statistical package for social science (SPSS) software version 17 was used.

Results

This study was conducted on asthmatic patients among those attending the outpatient chest clinic in Fayoum Hospital University. Their ages ranged from 20 to 60 years (with moderate to severe asthma).

Table 1 shows comparison between mean (SD) of FVC, FEV₁, FEV₁ ratio, and PEFR before and after therapy of the two groups shows that; mean (SD) of FVC of group 1

Download English Version:

<https://daneshyari.com/en/article/3400209>

Download Persian Version:

<https://daneshyari.com/article/3400209>

[Daneshyari.com](https://daneshyari.com)