



The Egyptian Society of Chest Diseases and Tuberculosis  
Egyptian Journal of Chest Diseases and Tuberculosis

[www.elsevier.com/locate/ejcdt](http://www.elsevier.com/locate/ejcdt)  
[www.sciencedirect.com](http://www.sciencedirect.com)



## ORIGINAL ARTICLE

# On improving assessment of in-hospital mortality and ICU admission in community-acquired pneumonia patients using the eCURB

Lamia Shaaban \*, Mohamed Metwally

Chest Diseases Department, Assiut University Hospital, Assiut, Egypt

Received 6 May 2012; accepted 16 May 2012

Available online 30 January 2013

### KEYWORD

Community-acquired pneumonia eCURB

**Abstract** *Background:* Assessment of severity of the disease in community-acquired pneumonia (CAP) is very important to decide the site of care. The conventional CURB-65 score is composed of five separate elements namely, Confusion, Uremia, Respiratory rate, BP, and age  $\geq 65$  years. These elements could be calculated electronically. The electronic CURB (eCURB) utilizes the 5 CURB-65 data elements as continuous, weighted variables. The aim of this study was to evaluate the performance of eCURB elements in predicting in-hospital mortality and ICU admission in comparison to the conventional CURB-65.

*Material and methods:* This study was conducted upon 134 adult patients diagnosed as CAP and confirmed by radiographic findings, admitted to chest department, Assiut University Hospital, Egypt. The CURB-65 elements were retrospectively extracted from the medical records. The eCURB variables were introduced to electronically calculate the risk using the Excel appendix model (provided by Prof. *Nathan Dean*, University of Utah, Salt Lake city, USA) and its predictive values and area under the receiver-operating characteristic (ROC) curve were compared with the conventional CURB-65 in predicting in-hospital mortality and the need for ICU admission.

*Results:* The study revealed that the conventional CURB-65 score could predict in-hospital mortality with an area under the curve (AUC) of 0.81 and the need for ICU admission with an AUC of 0.87. Using the eCURB-65 elements proved to be superior to the conventional CURB-65 in predicting in-hospital mortality with cut off point  $> 7.5$  and an AUC of 0.83 ( $P < 0.0001$ ). Also, eCURB was better than conventional CURB-65 in predicting ICU admission with cut off point  $> 3.8$  and an AUC of 0.89 ( $P < 0.0001$ ).

\* Corresponding author. Mobile: +20 01002681478.

E-mail address: [lamiashaban@yahoo.com](mailto:lamiashaban@yahoo.com) (L. Shaaban).

Peer review under responsibility of The Egyptian Society of Chest Diseases and Tuberculosis.



Production and hosting by Elsevier

*Conclusions:* Using the eCURB proved to be a valuable tool in predicting in-hospital mortality and ICU admission in patients with CAP with a significant superiority over conventional CURB-65 in both variables. Further prospective studies on a larger cohort are recommended.

© 2012 The Egyptian Society of Chest Diseases and Tuberculosis. Production and hosting by Elsevier B.V. Open access under CC BY-NC-ND license.

**Introduction**

The outcome of CAP is extremely variable and depends upon the affected host’s response, the underlying pathogen, and the treatment delivered. Assessment of severity of the patient’s disease is very important to decide the site of care in CAP patients [4,5]. However; this decision could be variable according to the need for hospital admission [1]. Therefore, accurate severity assessment during initial management is critical.

Two severity assessment tools have become widely used by clinicians and approved for use by the *ATS guidelines* [1] to help distinguish high-risk patients who require inpatient management from those able to thrive with outpatient management. The first one is the pneumonia severity index (PSI) developed by *Fine and colleagues* [2] is a prognostic model that calculates a severity-of-illness score based on 20 separate patient characteristics, including underlying co-morbidities. The second one is the CURB-65 score that is composed of five separate elements: Confusion, Uremia, Respiratory rate, BP, and age  $\geq 65$  years [3]. Although the PSI has been shown to be slightly more accurate at predicting outcome [4], CURB-65 is simpler to use. Additionally, all elements of the CURB-65 are routinely entered into the medical record, making it possible to generate an electronic mortality prediction for each patient at the point of care.

CURB-65 attributes a point to each criterion in an equally weighted fashion. However, excluding confusion, CURB-65 elements are actually continuous variables that may not be of equal predictive value. In other words, a systolic blood pressure of 85 mmHg is not like 70 mmHg although both are less than 90 mmHg and would be evaluated the same by the conventional CURB-65. So, the CURB-65 may be more accurate if calculated with continuous and weighted variables in the e-CURB model [5]. Instead of a severity score, a computer could generate an individualized mortality risk estimate using data elements from the electronic medical record. Generating

an automated, accurate mortality estimate immediately available to providers could improve severity assessment and thus improve care [5]. The aim of this study was to validate the accuracy of the new, electronic version of CURB-65 (eCURB) to predict ICU admission and in-hospital mortality compared to conventional CURB-65.

**Material and methods**

The study was conducted upon 134 patients diagnosed with CAP attending chest department of Assiut University Hospital; a tertiary care teaching university hospital. Retrospective analysis of data from the electronic medical record was done to identify all adult patients with CAP from August 2010 to December 2011. All patients must have radiographic evidence of CAP otherwise excluded. Patients diagnosed with aspiration pneumonia, having immuno-compromised conditions, hematologic malignancies, and those meeting criteria for health-care-associated pneumonia were all excluded from the study. Vital signs, orientation status at presentation and the routine laboratory results that were done within the first 12 h were extracted from the electronic medical record.

The conventional CURB-65 score was calculated and its performance in predicting the need for ICU admission and in-hospital deaths was evaluated. The methodology of the eCURB required a specific value for blood urea nitrogen (BUN). Also, the Systolic BP was used, as it was found to be non-significantly better than using diastolic BP, and the latter did not add any additional predictive value [5]. The performance of eCURB risk score in predicting ICU admission and in-hospital mortality was calculated using an Excel appendix model (lasso penalized logistic regression model) that was provided to us from the original developer of the scoring system (Prof. *Nanthan Dean* from University of Utah, Salt Lake city, USA, Personal communication with Dr. *Mohamed Metwally*, ATS meeting, Denver 2011) Fig. 1.

Age	Confused	First BUN	First RR	First SBP	Int	Age1	Age2	Age3	Confused	BUN1	BUN2	BUN3	RR1	RR2	RR3	SBP	Estimate	Final	
35	0	12.88	21	100	1	-0.115	0.440	-0.282	0	-0.309	0.677	-0.344	-0.049	0.647	-0.340	100	-4.394	1.2%	
70	0	13.44	25	140	1	0.539	0.361	-0.093	0	-0.314	0.703	-0.357	0.209	0.542	-0.255	140	-3.589	2.7%	
40	0	9.24	20	110	1	-0.093	0.519	-0.332	0	-0.257	0.539	-0.274	-0.121	0.667	-0.352	110	-4.917	0.7%	
65	0	17.92	26	110	1	0.478	0.407	-0.190	0	-0.285	0.730	-0.370	0.257	0.520	-0.230	110	-3.243	3.8%	
53	0	6.16	19	100	1	0.167	0.537	-0.339	0	-0.162	0.332	-0.169	-0.181	0.671	-0.395	100	-5.211	0.5%	
38	0	20.72	25	110	1	-0.106	0.492	-0.315	0	-0.242	0.716	-0.362	0.209	0.542	-0.255	110	-3.436	3.1%	
70	0	25.2	25	140	1	0.539	0.361	-0.093	0	-0.174	0.691	-0.348	0.209	0.542	-0.255	140	-2.919	5.1%	
66	0	11.2	40	130	1	0.495	0.397	-0.172	0	-0.298	0.641	-0.325	0.359	0.355	0.229	130	-3.337	3.4%	
19	50	0	11.2	20	120	1	0.085	0.556	-0.354	0	-0.298	0.641	-0.325	-0.121	0.667	-0.352	120	-4.892	0.7%
28	0	11.2	26	130	1	-0.093	0.281	-0.180	0	-0.298	0.641	-0.325	0.257	0.520	-0.230	130	-3.707	2.4%	
35	0	13.16	20	120	1	-0.115	0.440	-0.282	0	-0.314	0.703	-0.357	-0.121	0.667	-0.352	120	-4.623	1.0%	

**Figure 1** The excel appendix model (lasso penalized logistic regression model) for calculation of eCURB risk.

Download English Version:

<https://daneshyari.com/en/article/3400373>

Download Persian Version:

<https://daneshyari.com/article/3400373>

[Daneshyari.com](https://daneshyari.com)