




Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Brief report

Cost/efficacy analysis of preferred Spanish AIDS study group regimens and the dual therapy with lopinavir/ritonavir plus lamivudine for initial ART in HIV infected adults



Josep María Gatell Artigas^a, José Ramón Arribas López^b, Pablo Lázaro y de Mercado^{c,*}, Antonio Javier Blasco Bravo^c

^a Head Infectious Diseases and AIDS Units, Hospital Clinic-IDIBAPS, Barcelona, Spain

^b HIV Unit, La Paz Hospital, IdiPAZ, Madrid, Spain

^c Independent Researcher, Madrid, Spain

ARTICLE INFO

Article history:

Received 5 January 2015

Accepted 27 January 2015

Available online 3 March 2015

Keywords:

Cost
Efficacy
Efficiency
HIV
AIDS
Antiretroviral therapy

ABSTRACT

Introduction: The National AIDS Plan and the Spanish AIDS study group (GESIDA) proposes “preferred regimens” (PR) of antiretroviral treatment (ART) as initial therapy in HIV-infected patients. In 2013, the recommended regimens were all triple therapy regimens. The Gardel Study assessed the efficacy of a dual therapy (DT) combination of lopinavir/ritonavir (LPV/r) plus lamivudine (3TC). Our objective is to evaluate the GESIDA PR and the DT regimen LPV/r + 3TC cost/efficacy ratios.

Methods: Decision tree models were built. Efficacy: probability of having viral load <50 copies/mL at week 48. ART regime cost: costs of ART, adverse effects, and drug resistance tests during the first 48 weeks.

Results: Cost/efficacy ratios varied between 5,817 and 13,930 euros per responder at 48 weeks, for the DT of LPV/r + 3TC and tenofovir DF/emtricitabine + raltegravir, respectively.

Conclusions: Taking into account the official Spanish prices of ART, the most efficient regimen was DT of LPV/r + 3TC, followed by the triple therapy with non-nucleoside containing regimens.

© 2015 Elsevier España, S.L.U. and Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. All rights reserved.

Análisis de coste/eficacia de las pautas preferentes de GESIDA/Plan Nacional sobre el Sida y de biterapia con lopinavir/ritonavir más lamivudina para el tratamiento antirretroviral inicial en adultos infectados por el virus de la inmunodeficiencia humana

RESUMEN

Introducción: El Plan Nacional sobre el Sida y el grupo de estudio del SIDA-SEIMC (GESIDA) propone «regímenes preferentes» (RP) de tratamiento antirretroviral (TAR) como terapia inicial en pacientes infectados por VIH. Todos los regímenes recomendados en el año 2013 eran de terapia triple. El Estudio Gardel evaluó la eficacia de una doble terapia (DT) que combina lopinavir/ritonavir (LPV/r) más lamivudina (3TC). Nuestro objetivo es evaluar los ratios de coste/eficacia de los RP de GESIDA y el régimen de DT LPV/r + 3TC.

Métodos: Construcción de árboles de decisión como modelos de evaluación económica. Eficacia: probabilidad de tener una carga viral <50 copias/ml en la semana 48. Coste del régimen de TAR: coste del TAR, efectos adversos y tests de resistencia durante las primeras 48 semanas.

Palabras clave:

Coste
Eficacia
Eficiencia
VIH
SIDA
Terapia antirretroviral

* Corresponding author.

E-mail address: plazaro@gmx.es (P. Lázaro y de Mercado).

Resultados: Los ratios de coste/eficacia variaron entre 5.817 € y 13.930 € por respondedor a las 48 semanas, para la DT LPV/r + 3TC y tenofovir DF/emtricitabina + raltegravir, respectivamente.

Conclusiones: Con los precios oficiales españoles de TAR, el régimen más eficiente fue la DT con LPV/r + 3TC, seguida de la terapia triple con regímenes que no contienen nucleósidos.

© 2015 Elsevier España, S.L.U. y Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Todos los derechos reservados.

Introduction

The National AIDS Plan and the Spanish AIDS study group (GESIDA) panel of experts propose, every year, “preferred regimens” (PR) of antiretroviral treatment (ART) as initial therapy for HIV infected patients. The PR for the year 2013 were all triple therapy regimens.¹ After the publication of the 2013 GESIDA preferred regimens, the GARDEL (Global AntiRetroviral Design Encompassing Lopinavir (LPV)/ritonavir (r) and Lamivudine (3TC) vs LPV/r based standard therapy) Study was published.² The GARDEL trial assessed the efficacy and safety of a dual therapy (DT) combination of LPV/r 400/100 mg + 3TC 150 mg, both twice daily compared to triple therapy regimens of LPV/r 400/100 mg twice daily plus 3TC or emtricitabine (FTC) plus one other, investigator selected nucleos(t)ide reverse transcriptase inhibitor. The objective of this study is to evaluate the costs and efficiency of initiating treatment with the GESIDA PR and the DT of LPV/r + 3TC.

Methods

Design: Economic assessment of the costs and efficiency (cost/efficacy) by building decision trees with deterministic sensitivity analysis. The decision trees for calculating the costs, efficacy, and efficiency of the 2013 PR have been previously published.³ A new decision tree was built for the DT following the same methodology. The design, methods for building the trees, sources of data, definition of efficacy, costs and efficiency, use of resources, and sensitivity analysis have been the same as those used previously for the 2013 PR.³ In summary, such methods have been the following.

Perspective: The Payer (Spanish National Health System) perspective was applied. The following differential direct costs were considered: (1) ART (Laboratory sale price + 4% VAT – 7.5% obligatory legal reduction). In the case of DT, for 3TC, the generic price (65.27 euros: Laboratory sale price + 4% VAT) was considered, resulting in a cost of 5,041 euros for 48 weeks; (2) Adverse events (AE) management (drug treatment, emergency room visits, additional visits to the HIV specialist, visits to other specialists, diagnostic tests, and hospital admissions). Each unitary cost was calculated as the mean of the official prices of the Autonomous Communities (regions) Health Services; and (3) Genotypic study of drug resistance and HLA B*5701 testing.

Time horizon: 48 weeks.

Cost of initiating a regimen: Cost of ART and all the consequences (adverse effects, changes of ART regimen and drug resistance tests) incurred in 48 weeks due to the decision of initiating ART with that regimen. The substitution regimens for DT depending of the cause were decided by two of the authors (JMG and JRA): Tenofovir DF (TDF)/FTC + LPV/r for viral failure and pregnancy, TDF/FTC/efavirenz (EFV) for adverse effect, TDF/FTC + darunavir (DRV)/r for lack of adherence, and DT for lost to follow-up and others causes.

Efficacy: Quotient of the number of patients with undetectable viral load (<50 copies/mL) at week 48 post-ART (i.e., responders) (numerator) and the number of patients initiated on ART (denominator). It was estimated based on an intention-to-treat analysis of

the exposed (“Intent-to-treat exposed” [ITT-E]), “missing or non-completer = failure”).

Efficiency: Defined in terms of cost/efficacy and calculated for each regimen as the quotient of the cost of initiating treatment with that regimen (numerator) and efficacy (denominator). It represents the cost of achieving one responder by week 48.

Sources of information: Clinical trials (CTs): Data on efficacy, AE and withdrawals. Data of CTs included in a previous study³ and the GARDEL Study.² The expert opinion was used when scientific evidence was not available (substitution regimens and resources used in AE management).

Uncertainty management: Deterministic sensitivity analysis, building scenarios with 95% confidence intervals for efficacy and AE probability, and $\pm 15\%$ for costs.

Results

In the base case scenario, the cost of initiating treatment ranges from 5,138 euros for DT of LPV/r + 3TC to 12,059 euros for TDF/FTC + raltegravir (RAL). The efficacy ranged between 0.66 for abacavir (ABC)/3TC + LPV/r and ABC/3TC + atazanavir (ATV)/r, and 0.88 for DT of LPV/r + 3TC. The efficiency, in terms of cost/efficacy, varied between 5,817 and 13,930 euros per responder at 48 weeks, for DT of LPV/r + 3TC and TDF/FTC + RAL respectively. Moreover the DT regimen was the most efficient regimen in the most favorable (5,503 euros per responder) and most unfavorable (6,169 euros per responder) scenarios (Table 1 and Fig. 1).

Since local cost of a specific hospital may be different to the costs used in the model, a software application was designed for allowing the calculation of ART costs, regimen initiation costs, efficiency (cost/efficacy), and relative efficiency of initiating treatment with the different regimens at each individual hospital setting. The application is available free of charge at: <https://www.dropbox.com/s/875w8ye95j640ci/Aplicacion-TARV-VIH-2013-TerapiaDoble.exe?dl=0>.

Discussion

According to the findings of this analysis, considering the ART official Spanish prices, the most efficient regimen was LPV/r + 3TC, followed by triple therapy with non-nucleoside containing regimens recommended in 2013 by the Spanish GESIDA expert panel. This finding should be interpreted carefully taking into account some limitations. The specific limitations affecting the GARDEL trial and the cost/efficacy analysis performed for the GESIDA recommended regimens are described in their respective articles.^{2,3} But some limitations should be mentioned. For example, the analyses presented here are based on clinical trials performed in different countries, in different years (published between 2006 and 2014), with different inclusion and exclusion criteria, and even with different presentations for the same drug in some regimens. Thus, the results may have differed if all regimens had been administered in similar populations and years. Actually, recent studies include lower percentages of patients with poor prognosis, i.e., those with low CD4 counts (<100/200 cells/ μ L) and elevated plasma viral load (>100,000 copies/mL). This leads to results with

Download English Version:

<https://daneshyari.com/en/article/3400509>

Download Persian Version:

<https://daneshyari.com/article/3400509>

[Daneshyari.com](https://daneshyari.com)