

HOSTED BY



ELSEVIER

Available at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/IJMYCO

Hookworm infection among patients with pulmonary tuberculosis: Impact of co-infection on the therapeutic failure of pulmonary tuberculosis

Ahmad Farooq Alsayed Hasanain^{a,*}, Ali Abdel-Azeem Hasan Zayed^b, Reem Ezzat Mahdy^c, Amany Mohamed Adawi Nafee^d, Rasha Abdel-Monem Hassan Attia^e, Asmaa Omar Mohamed^f

^aDepartment of Tropical Medicine and Gastroenterology, Faculty of Medicine, Assiut University, Assiut, Egypt

^bDepartment of Chest Diseases, Faculty of Medicine, Assiut University, Assiut, Egypt

^cDepartment of Internal Medicine, Faculty of Medicine, Assiut University, Assiut, Egypt

^dDepartment of Microbiology and Immunology, Faculty of Medicine, Assiut University, Assiut, Egypt

^eDepartment of Parasitology, Faculty of Medicine, Assiut University, Assiut, Egypt

^fDepartment of Clinical Pathology, Faculty of Medicine, Assiut University, Assiut, Egypt

ARTICLE INFO

Article history:

Received 28 August 2015

Accepted 2 September 2015

Available online 1 October 2015

Keywords:

Coinfection

Hookworm

Therapeutic failure

Tuberculosis

ABSTRACT

Objective/background: The aim of this study is to determine the rate of hookworm infection among patients with pulmonary tuberculosis (TB) and to find out if there is a relation between hookworm infection and the therapeutic failure of pulmonary TB.

Methods: We carried out a prospective, hospital-based study. The study included 231 naïve patients with pulmonary TB, consecutively. Patients were evaluated at the 4th month of therapy for persistence of *Mycobacterium tuberculosis* infection. All patients had clinical evaluation, laboratory investigations (including sputum culture and stool microscopic examination), and imaging studies (abdominal ultrasonography and chest radiography).

Results: The study population mean age was 42.7 ± 13.9 years old with 26.8% of them 40 years old or more. Out of 231 patients, 133 (57.6%) were men. Therapeutic failure rate of pulmonary TB was 29.4%. Hookworm infection was diagnosed among 16.5% of patients and 27.7% had diabetes mellitus (DM). Using multivariate analysis, it was found that age of 40 years or more (odds ratio [OR] 8.4; 95% confidence interval [CI] 1.7–41.3; $p = .009$), hookworm infection (OR 7.6; 95% CI 1.2–49.9; $p = .034$), and DM (OR 5.9; 1.2–28; $p = .027$) were independently associated with therapeutic failure of pulmonary TB among the study population with pulmonary TB.

Conclusion: In conclusion, the rate of therapeutic failure of pulmonary TB is high. Besides older age and DM, hookworm infection can reduce the therapeutic response of pulmonary TB. Screening for and control of DM and hookworm infection among patients with pulmonary TB may improve their therapeutic response.

© 2015 Asian African Society for Mycobacteriology. Production and hosting by Elsevier Ltd.

All rights reserved.

* Corresponding author at: Department of Tropical Medicine and Gastroenterology, Assiut University Hospital, 6th Floor, Assiut, 71515, Egypt.

E-mail address: a.fh@outlook.com (A.O. Mohamed).

Peer review under responsibility of Asian African Society for Mycobacteriology.

<http://dx.doi.org/10.1016/j.ijmyco.2015.09.002>

2212-5531/© 2015 Asian African Society for Mycobacteriology. Production and hosting by Elsevier Ltd. All rights reserved.

Introduction

Tuberculosis (TB) is a major global health problem, with an estimated 9 million new cases and 1.5 million deaths during 2013 according to the World Health Organization (WHO) [1]. The standard four-drug regimen (isoniazid, rifampin, ethambutol, and pyrazinamide for 2 months followed by isoniazid and rifampin for 4 months) has high efficacy in achieving cure rates (90–95%) [2]. To monitor pulmonary TB patients, culture examinations are recommended at monthly intervals until two consecutive culture specimens are negative at the end of treatment [3]. According to WHO, the therapeutic failure of pulmonary TB is defined as positive sputum culture at the 4th month or later during treatment [4]. The gold standard for the diagnosis of TB is culture [5].

More than 500 million people are infected with hookworm globally. The diagnosis of hookworm infection is made with microscopic identification of the characteristic ova in stool [6]. The overlapping distribution of hookworm infection with other pathogens results in a high rate of coinfection [7]. It was speculated that helminth infections, including hookworm, may adversely influence the host-immune response to other infections, especially to intracellular pathogens such as *Mycobacterium tuberculosis* (Mtb) [8,9]. This has led to the suggestion that helminth infection may increase both the susceptibility to, and the disease progression of, bacterial diseases. This hypothesis is based on the observation that helminth infections results in eosinophilia, elevated immunoglobulin E, and mast cell activation, suggestive of the enhanced cytokine response by type 2-T-helper cells (Th2) and a concomitant suppression of a type 1-T-helper cells (Th1) response, which is often involved in the protection against intracellular pathogens [10].

Both pulmonary TB and hookworm infections are common diseases in Egypt. The therapeutic failure of pulmonary TB is not uncommonly encountered in upper Egypt. We lack studies exploring the impact of hookworm infection on the rate of therapeutic failure among patients with pulmonary TB. The aim of this study is to determine the rate of hookworm infection among patients with pulmonary TB, and to find out if there is a relation between hookworm infection and the therapeutic failure of pulmonary TB.

Materials and methods

We carried out a prospective, hospital-based study at Assiut University Hospital during the period of June 2011 to November 2014. The study included 231 naïve patients with pulmonary TB, consecutively. Pulmonary TB was diagnosed based on positive sputum culture on Löwenstein–Jensen medium performed by an experienced microbiologist, with or without evidence on chest radiography [5]. All patients received the WHO-recommended therapy: isoniazid (5 mg/kg/day), rifampin (10 mg/kg/day; maximum dose was 600 mg/day), ethambutol (20 mg/kg/day; maximum dose was 1600 mg/day), and pyrazinamide (30 mg/kg/day; maximum dose was 2000 mg/day) for 2 months, followed by isoniazid

and rifampin for 4 more months [11]. Therapeutic failure of pulmonary TB was defined as positive sputum culture at the 4th month or later during treatment [4]. The diagnosis of hookworm infection was made with microscopic identification of the characteristic ova in stool [6] performed by an experienced parasitologist using three consecutive stool samples. Patients with extrapulmonary TB or intestinal parasitic infection other than hookworm infection were excluded from the study.

Patients were evaluated at the 4th month of therapy for persistence of Mtb infection. All patients had clinical evaluation, laboratory investigations, and imaging studies. Laboratory investigations included sputum culture on Löwenstein–Jensen medium, stool microscopic examination, complete blood count, liver chemistry, kidney chemistry, fasting level of serum glucose, and testing for antibody to human immunodeficiency virus. Imaging studies included abdominal ultrasonography and chest radiography. Long-term use of either proton pump inhibitors or corticosteroids was defined as use for 1 month or more during treatment for pulmonary TB.

Ethical considerations

The study was approved by the Faculty of Medicine Clinical Research Ethical Committee (Assiut University), and was carried out in accordance with the code of ethics of the World Medical Association (Declaration of Helsinki). Before enrollment in the study, all the participants signed a consent certificate. Before signing, they were able to discuss in detail with the investigator the certificate subjects and the study aim. The participants were clearly informed that refusing to participate in the study will not affect having full benefit of the available medical service and treatment. Data were collected by personal interview with the participants taking in consideration data confidentiality.

Statistical analysis

The data were entered and analyzed using the Statistical Package for Social Sciences (SPSS, Chicago IL, USA) for windows, version 22). Results were expressed as mean \pm standard deviation or frequency (percentage) as appropriate. To compare the distribution of demographic criteria between patients with and without therapeutic failure of pulmonary TB, we used Chi-square test. Predictors of the therapeutic failure of pulmonary TB were identified using univariate analysis (Yates' corrected Chi-square test) then multivariate analysis (stepwise binary logistic regression) to assess the specific effect of each predictor. Multivariate analysis included factors with significant level ($p < .05$) in univariate analysis.

Results

The study included 231 patients diagnosed as having pulmonary TB. The demographic criteria of the study population with pulmonary TB are shown in Table 1. Their mean age was 42.7 ± 13.9 years old, with 26.8% of them being 40 years of age

Download English Version:

<https://daneshyari.com/en/article/3404957>

Download Persian Version:

<https://daneshyari.com/article/3404957>

[Daneshyari.com](https://daneshyari.com)