



## Plasmodial ortholog of *Toxoplasma gondii* rhoptry neck protein 3 is localized to the rhoptry body

Daisuke Ito<sup>a</sup>, Eun-Taek Han<sup>b</sup>, Satoru Takeo<sup>a</sup>, Amporn Thongkukiatkul<sup>c</sup>, Hitoshi Otsuki<sup>d</sup>, Motomi Torii<sup>e</sup>, Takafumi Tsuboi<sup>a,f,\*</sup>

<sup>a</sup> Cell-Free Science and Technology Research Center, Ehime University, Matsuyama, Ehime 790-8577, Japan

<sup>b</sup> Department of Parasitology, Kangwon National University College of Medicine, Chuncheon, Republic of Korea

<sup>c</sup> Department of Biology, Faculty of Science, Burapha University, Chonburi 20131, Thailand

<sup>d</sup> Division of Medical Zoology, Faculty of Medicine, Tottori University, Yonago, Tottori 683-8503, Japan

<sup>e</sup> Department of Molecular Parasitology, Ehime University Graduate School of Medicine, Shitsukawa, Toon, Ehime 791-0295, Japan

<sup>f</sup> Venture Business Laboratory, Ehime University, Matsuyama, Ehime 790-8577, Japan

### ARTICLE INFO

#### Article history:

Received 23 November 2010

Received in revised form 24 December 2010

Accepted 7 January 2011

Available online 13 January 2011

#### Keywords:

Cell-free expression

Erythrocyte invasion

Merozoite

*Plasmodium falciparum*

Rhoptry

### ABSTRACT

The proteins in apical organelles of *Plasmodium falciparum* merozoite play an important role in invasion into erythrocytes. Several rhoptry neck (RON) proteins have been identified in rhoptry proteome of the closely-related apicomplexan parasite, *Toxoplasma gondii*. Recently, three of *P. falciparum* proteins orthologous to *TgRON* proteins, *PfRON2*, 4 and 5, were found to be located in the rhoptry neck and interact with the micronemal protein apical membrane antigen 1 (*PfAMA1*) to form a moving junction complex that helps the invasion of merozoite into erythrocyte. However, the other *P. falciparum* RON proteins have yet to be characterized. Here, we determined that “PFL2505c” (hereafter referred to as *pfron3*) is the ortholog of the *tgron3* in *P. falciparum* and characterized its protein expression profile, subcellular localization, and complex formation. Protein expression analysis revealed that *PfRON3* was expressed primarily in late schizont stage parasites. Immunofluorescence microscopy (IFA) showed that *PfRON3* localizes in the apical region of *P. falciparum* merozoites. Results from immunoelectron microscopy, along with IFA, clarified that *PfRON3* localizes in the rhoptry body and not in the rhoptry neck. Even after erythrocyte invasion, *PfRON3* was still detectable at the parasite ring stage in the parasitophorous vacuole. Moreover, co-immunoprecipitation studies indicated that *PfRON3* interacts with *PfRON2* and *PfRON4*, but not with *PfAMA1*. These results suggest that *PfRON3* partakes in the novel *PfRON* complex formation (*PfRON2*, 3, and 4), but not in the moving junction complex (*PfRON2*, 4, 5, and *PfAMA1*). The novel *PfRON* complex, as well as the moving junction complex, might play a fundamental role in erythrocyte invasion by merozoite stage parasites.

© 2011 Elsevier Ireland Ltd. All rights reserved.

### 1. Introduction

Malaria is caused by the replication of protozoan parasites of the genus *Plasmodium* in circulating host erythrocytes [1]. The invasion process of merozoite stage parasite into erythrocyte requires the discharge of contents of apical secretory organelles (micronemes and rhoptries) to form an irreversible contact, called a tight junction, between the merozoite and the erythrocyte membrane. This tight junction migrates from the anterior to posterior poles of the merozoite. According to this moving junction, the host membrane invaginates the merozoites to eventually form a parasitophorous vacuole [2,3]. The moving junction is one of the most distinctive features of apicomplexan invasion into host cells, and was first

observed in *Plasmodium* species [4]. Studies in apicomplexan parasite *Toxoplasma gondii* identified a total of four proteins from distinct apical secretory organelles to form a moving junction complex; micronemal protein apical membrane antigen 1 (AMA1) and three rhoptry neck (RON) proteins, RON2, RON4, and RON5 [5,6]. Recently, this RON–AMA1 complex (*PfRON2*, *PfRON4*, *PfRON5*, and *PfAMA1*) was also identified in *Plasmodium falciparum* [7–9]. Attempts to knockout the AMA1 gene locus were unsuccessful in both *Plasmodium* [10] and *T. gondii* [11]. AMA1-binding peptide R1 not only prevents RON–AMA1 complex interaction, but also blocks *P. falciparum* merozoite invasion into erythrocytes [9].

Although the cumulative evidence above strongly suggests that the conserved RON–AMA1 complex plays an essential role in merozoite invasion, it is yet to be clarified whether molecules other than RON2, RON 4, and RON 5, play roles in the formation of the moving junction complex and in the invasion process. The report on *T. gondii* rhoptry proteome identified the presence of other RON proteins, RON1 and RON3 [12]. Therefore, we were interested in

\* Corresponding author. Cell-Free Science and Technology Research Center, Ehime University, Matsuyama, Ehime 790-8577, Japan. Tel.: +81 89 927 8277; fax: +81 89 927 9941.

E-mail address: [tsuboi@ccr.ehime-u.ac.jp](mailto:tsuboi@ccr.ehime-u.ac.jp) (T. Tsuboi).

identifying the ortholog of *tgron3* in *P. falciparum* and in characterizing its protein expression profile, subcellular localization, and role in the formation of the RON–AMA1 complex.

## 2. Materials and methods

### 2.1. Parasites

*P. falciparum* asexual stages (3D7 strain) were maintained in human erythrocytes (blood group O<sup>+</sup>) *in vitro*, as previously described [13].

### 2.2. RNA isolation and cDNA synthesis

Total RNA was extracted from *P. falciparum* schizont-infected erythrocytes (purified by differential centrifugation on a 70%/40% Percoll/sorbitol gradient) using the TRIzol Reagent (Invitrogen, Carlsbad, CA). Following DNase treatment, cDNA was generated with a random hexamer using the SuperScriptIII® First Strand Synthesis System (Invitrogen).

### 2.3. Antibodies

Recombinant *Pf*RON3 proteins were produced using the wheat germ cell-free translation system (CellFree Sciences, Matsuyama, Japan) as described previously [14–16]. Briefly, regions of the *Pf*RON3 gene encoding the deduced amino acid sequence, 927–1056 (*Pf*RON3\_1) and 1686–1884 (*Pf*RON3\_2), were PCR-amplified from *P. falciparum* 3D7 blood-stage cDNA and ligated into pEU-E01-GST-(TEV)-N2 (CellFree Sciences), an expression plasmid with an N-terminal glutathione S transferase (GST)-tag followed by a tobacco etch virus (TEV) protease cleavage site, designed specifically for the wheat germ cell-free protein expression. Oligonucleotide primers used in the PCR amplification were *Pf*RON3\_XhoIF1 (5′-ctcggagGATATTCATTAAAAGAAACCTATAAATT-3′) and *Pf*RON3\_BamHIR1 (5′-ggatccCTAATGTGGGAACATTTTCATGATTGGTA-3′) for *Pf*RON3\_1, and *Pf*RON3\_XhoIF2 (5′-ctcggagGATTTAAAGATAAATCAGATGATGATC-3′) and *Pf*RON3\_BamHIR2 (5′-ggatccCTATTTTTAGGTACATATATATTATATGGTC-3′) for *Pf*RON3\_2 (XhoI and BamHI restriction sites are underlined). Both GST-*Pf*RON3\_1 and GST-*Pf*RON3\_2 were captured using a glutathione-Sepharose 4B column (GE Healthcare, Camarillo, CA), and eluted by on-column cleavage with 60 U of AcTEV protease (Invitrogen) after extensive washing with PBS. To generate anti-*Pf*RON3\_1 and anti-*Pf*RON3\_2 sera, female BALB/c mice were immunized subcutaneously with 20 µg of purified *Pf*RON3\_1 or *Pf*RON3\_2 emulsified with Freund's adjuvant. A Japanese white rabbit was also immunized subcutaneously with 250 µg of purified *Pf*RON3\_1 or *Pf*RON3\_2 emulsified with Freund's adjuvant. All immunizations were performed 3 times at 3-week intervals, and then antisera were collected 2 weeks after the third immunization. In a similar manner, mouse anti-*Pf*RAP1 (aa 1–782) antibody, mouse anti-*Pf*EXP2 (aa 25–287) antibody, mouse and rabbit anti-*Pf*AMA1 (aa 25–546) antibodies, and rabbit anti-GST antibody, were generated as control. Rabbit antisera against the *Pf*RON3\_1 and *Pf*RON3\_2 proteins were affinity purified using a column conjugated with recombinant *Pf*RON3\_1 or *Pf*RON3\_2 as ligands, respectively. Briefly, recombinant *Pf*RON3\_1 or *Pf*RON3\_2 was covalently linked to a HiTrap™ NHS-activated HP column (GE Healthcare) as manufacturer's recommendation. Rabbit antiserum was then applied to either the *Pf*RON3\_1- or the *Pf*RON3\_2-conjugated column. After an extensive washing step with 20 mM phosphate buffer (pH 7.0), antigen-specific IgGs were eluted with 0.1 M glycine–HCl (pH 2.5), and then immediately neutralized with 1 M Tris (pH 9.0). Mouse monoclonal anti-*Pf*RON4 antibody (26C64F12) [7] and anti-*Pf*RESA antibody (23/9) [17] were kind gifts from Jean F. Dubremetz (Université de Montpellier 2, France) and Robin F. Anders (La Trobe University, Australia), respectively.

### 2.4. SDS-PAGE and western blot analysis

*P. falciparum* cultured parasites were harvested after tetanolysin (List Biological Laboratories, Inc., Campbell, CA) treatment that can remove the hemoglobin without loss of parasite proteins present in the parasitophorous vacuolar space [18]. The parasite proteins were then extracted in SDS-PAGE loading buffer, incubated at 4 °C for 6 h, and subjected to electrophoresis under reducing condition on a 12.5% polyacrylamide gel (ATTO, Tokyo, Japan). Proteins were then transferred to a 0.2-µm PVDF membrane (GE Healthcare). The proteins were immunostained with antisera followed by horseradish peroxidase-conjugated secondary antibody (GE Healthcare) and visualized with Immobilon Western Chemiluminescent HRP Substrate (Millipore, Billerica, MA) on a LAS 4000 mini luminescent image analyzer (GE Healthcare). The relative molecular masses of the proteins were estimated with reference to Precision Plus Protein Standards (BioRad, Hercules, CA).

### 2.5. Immunoprecipitation

Immunoprecipitation was carried out as previously described [19]. Briefly, proteins were extracted from late schizont parasite pellets in PBS with 1% Triton X-100 containing Complete Proteinase Inhibitor Cocktail (Roche, Indianapolis, IN). Supernatants (50 µl) were pre-incubated at 4 °C for 1 h with 40 µl of 50% protein G-conjugated beads (GammaBind Plus Sepharose; GE Healthcare) in NETT buffer (50 mM Tris–HCl, 0.15 M NaCl, 1 mM EDTA, and 0.5% Triton X-100) supplemented with 0.5% BSA (fraction V; Sigma-Aldrich Corporation, St. Louis, MO). Aliquots of recovered supernatants were incubated either with rabbit anti-*Pf*RON3\_1, anti-*Pf*AMA1, anti-*Pf*RON2, or anti-GST antibody, and then 40 µl of 50% protein G-conjugated bead suspension was added. After one-hour incubation at 4 °C, the beads were washed once with NETT–0.5% BSA, once with NETT, once with high-salt NETT (0.5 M NaCl), once with NETT, and once with low-salt NETT (0.05 M NaCl and 0.17% Triton X-100). Finally, proteins were eluted from the protein G-conjugated beads with 0.1 M glycine–HCl (pH 2.5), and then immediately neutralized with 1 M Tris pH 9.0. The supernatants were used for western blot analysis.

### 2.6. Indirect immunofluorescence assay

Thin smears of ring or schizont-enriched *P. falciparum*-infected erythrocytes were prepared on glass slides and stored at –80 °C. The smears were thawed, fixed with 4% formaldehyde at room temperature for 10 min, permeabilized with PBS containing 0.1% Triton X-100 at room temperature for 15 min, and blocked with PBS containing 5% non-fat milk at 37 °C for 30 min. The smears were then incubated with rabbit anti-*Pf*RON3 antibodies (1:500 dilution) and control mouse antibodies at 37 °C for 1 h, followed by incubation with both Alexa Fluor 488-conjugated goat anti-rabbit IgG and Alexa Fluor 546-conjugated goat anti-mouse IgG (Invitrogen) as secondary antibodies (1:500) at 37 °C for 30 min. Nuclei were stained with 4',6-diamidino-2-phenylindole (2 µg/ml, DAPI) mixed with a secondary antibody solution. Slides were mounted in ProLong Gold Antifade reagent (Invitrogen) and viewed under ×63 oil-immersion lens. High-resolution image-capture and processing were performed using a confocal scanning laser microscope (LSM5 PASCAL or LSM710; Carl Zeiss MicroImaging, Thornwood, NY). Images were processed in Adobe Photoshop (Adobe Systems Inc., San José, CA).

### 2.7. Immunoelectron microscopy

Parasites were fixed for 15 min on ice in a mixture of 1% paraformaldehyde and 0.1% glutaraldehyde in 0.1 M phosphate buffer (pH 7.4). Fixed specimens were washed, dehydrated, and embedded in LR White resin (Polysciences, Inc., Warrington, PA) as previously

Download English Version:

<https://daneshyari.com/en/article/3418183>

Download Persian Version:

<https://daneshyari.com/article/3418183>

[Daneshyari.com](https://daneshyari.com)