




available at www.sciencedirect.com



journal homepage: www.elsevierhealth.com/journals/trst



Cockroaches as carriers of human intestinal parasites in two localities in Ethiopia

Addisu Kinfu^a, Berhanu Erko^{b,*}

^a St Lideta Health Science College, P.O. Box 120004, Addis Ababa, Ethiopia

^b Akilu Lemma Institute of Pathobiology, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia

Received 21 January 2008; received in revised form 13 May 2008; accepted 15 May 2008

Available online 25 June 2008

KEYWORDS

Cockroaches;
Blattellidae;
Carriers;
Intestinal parasites;
Ethiopia

Summary A study was undertaken to assess the role of cockroaches as potential carriers of human intestinal parasites in Addis Ababa and Ziway, Ethiopia. A total of 6480 cockroaches were trapped from the two localities from October 2006 to March 2007. All the cockroaches trapped in Addis Ababa ($n=2240$) and almost 50% (2100/4240) of those trapped in Ziway were identified as *Blattella germanica*. The rest of the cockroaches trapped in Ziway were identified as *Periplaneta brunnea* (24.52%), *Pycnoscelus surinamensis* (16.03%) and *Supella longipalpa* (9.90%). Microscopic examination of the external body washes of pooled cockroaches and individual gut contents revealed that cockroaches are carriers of *Entamoeba coli* and *Entamoeba histolytica/dispar* cysts as well as *Enterobius vermicularis*, *Trichuris trichiura*, *Taenia* spp. and *Ascaris lumbricoides* ova. Besides their role as a nuisance, the present study further confirms that cockroaches serve as carriers of human intestinal parasites. The possible association of cockroaches with allergic conditions such as asthma is also discussed. Hence, appropriate control measures should be taken particularly to make hotels and residential areas free of cockroaches as they represent a health risk.

© 2008 Royal Society of Tropical Medicine and Hygiene. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Cockroaches are insects with a worldwide distribution, thriving best in warm, humid, lowland areas throughout the tropics. Of 4000 species of cockroaches (Dictyoptera: Blattodea) known to exist, approximately 30 species are

cosmopolitan inhabitants of human dwellings and approximately 16 species are associated with human health problems (Naumann, 1991). The cockroach species that are closely associated with human dwellings, food processing industries, service rendering facilities and/or occupational environments include *Blattella germanica*, *Blatta orientalis*, *Periplaneta australasiae* (Fabricius), *Periplaneta americana* (Linnaeus) and *Supella supellectilium* (Serville) (Kang, 1990; Ross, 1965).

Domestic pest cockroaches have been documented to affect human health in several ways. They serve as

* Corresponding author. Tel.: +251 11 276 30 91;

fax: +251 11 275 52 96.

E-mail address: berhanue@yahoo.com (B. Erko).

intermediate hosts of some helminths (Beaver et al., 1984) and as mechanical vectors of numerous human intestinal protozoa and helminth parasites (Ash and Greenberg, 1980; Chan et al., 2004; Tatfeng et al., 2005) and are also known to harbour a wide variety of microbial pathogens for which they serve as potential transit hosts (Baumholtz et al., 1997; Roth and Willis, 1957; Salehzadeh et al., 2007). Besides serving as mechanical transmitters of parasites or pathogens, large indoor cockroach populations are also one of the leading causes of allergies, asthma and other bronchial disorders in humans (Kang, 1990; Miller and Koehler, 2003).

The filthy breeding habits, feeding mechanisms and indiscriminate travel between filth and food make cockroaches and other synanthropic insects efficient vectors of human enteric protozoan parasites (Graczyk et al., 2005). Cockroaches frequently feed on human faeces (Fotedar et al., 1991; Graczyk et al., 2005; Pai et al., 2003) and hence can disseminate cysts, ova and larvae of gastrointestinal parasites mechanically in the environment. In Ethiopia, a number of human intestinal parasites were isolated from cycloraphan flies (Getachew et al., 2007). However, no work has been done to elucidate the role of cockroaches as carriers of human intestinal parasites. The aims of this study were therefore to assess the role of cockroaches as potential carriers of human intestinal parasites and to identify species that serve as mechanical vectors of the parasites.

2. Materials and methods

This study was carried out in Addis Ababa and Ziway, Ethiopia, from October 2006 to March 2007. The city of Addis Ababa lies in the central highlands at an altitude of approximately 2500 m above sea level. The average annual temperature and rainfall are 21 °C and 1800 mm, respectively. Addis Ababa has a relative humidity that varies from 70–80% during the rainy season and from 40–50% during the dry season. The human population of Addis Ababa is estimated to be approximately 5 million.

Ziway is a town in central Ethiopia, located approximately 160 km to the south of Addis Ababa at an elevation of 1643 m above sea level. The average annual temperature of the area is approximately 20.5 °C and the mean annual rainfall is approximately 702 mm but can be as high as 1400 mm. The annual relative humidity of Ziway area ranges from 65% to 75%. Based on figures from the Central Statistical Agency of 2005, Ziway has an estimated total population of 35 931, of whom 19 034 were males and 16 897 were females.

2.1. Trapping and identification of cockroaches

After a pilot inspection to estimate the population and concentrated locations, indoor cockroaches were trapped from houses and their surroundings both in Addis Ababa and Ziway. Cockroaches were trapped during the night in empty jars coated with a thin film of Vaseline baited with a slice of bread soaked in beer or water. The outsides of the jars were wrapped with paper to allow the grasp of cockroaches as they climb up the sides of the jars. The sticky traps were placed in and around houses at different sites (kitchens, toilets and bedrooms) overnight along lines where the vertical and horizontal planes meet. Trapped

cockroaches from both localities were placed in labelled jars and transported to the laboratory for identification and processing for parasite examination. Cockroaches processed after a few days were placed in a refrigerator at –4 °C. Identification of cockroaches was then done using the procedures and keys described by Ross (1965). Furthermore, the identified species were confirmed by an experienced entomologist from Addis Ababa University.

2.2. Processing of cockroaches for parasite examination

Cockroaches trapped from the study localities were killed in a jar using diethyl ether-soaked cotton. Cockroaches were grouped into species and pooled into batches. Based on size difference among the species, *B. germanica* and *Supella longipalpa* were pooled in a batch of ten and *Periplaneta brunnea* and *Pycnoscelus surinamensis* were pooled in a batch of four. The cockroaches in a pool were immersed in sterile physiological saline, stirred for 5 min and vortexed for 2–3 min to wash off the parasite cysts, eggs or larvae that might be attached to the external body of the cockroaches. The washed cockroaches were then removed with forceps and the whole gut of each washed cockroach was dissected out on an autoclaved microscope slide under a stereoscopic microscope using entomological needles and macerated to liberate the lumen contents. To avoid cross-contamination, the needles and forceps were dipped in ethanol and flamed between dissections. Both the pooled external washes and the gut contents of cockroaches were microscopically examined for parasite material after being processed using the formol-ether concentration technique (WHO, 1991). The modified Ziehl–Neelsen staining method was used for examination of *Cryptosporidium* spp. (Adegbola et al., 1994). The specimens were then examined under the microscope at 10× or 40× magnification for helminth ova/larvae and for protozoan cysts/oocysts. The parasites were then identified based on morphology of their eggs and cysts (WHO, 1991) and their number was recorded.

2.3. Statistical analysis

All data obtained from this study were entered into Microsoft Excel (Microsoft Corp., Redmond, WA, USA) and analysed using SPSS version 13.0 statistical software (SPSS Inc., Chicago, IL, USA). Both descriptive and analytical statistics were employed. Comparison based on localities, cockroach species and body parts with respect to parasite burden was performed using χ^2 test.

3. Results

A total of 6480 cockroaches were trapped from Addis Ababa and Ziway. Cockroach species and associated protozoan and helminthic parasites from the body wash and gut contents of the cockroaches as well as parasite load and number of pools examined/positive are presented in Table 1. All cockroaches trapped from Addis Ababa and approximately 50% of cockroaches trapped from Ziway were identified as *B. germanica*. The rest of the cockroaches trapped in Ziway were

Download English Version:

<https://daneshyari.com/en/article/3420642>

Download Persian Version:

<https://daneshyari.com/article/3420642>

[Daneshyari.com](https://daneshyari.com)