

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Asian Pacific Journal of Reproduction

journal homepage: www.apjr.netOriginal research <http://dx.doi.org/10.1016/j.apjr.2016.06.004>

Natural honey as a cryoprotectant to improve Arab stallion post-thawing sperm parameters

Reda I. El-Sheshtawy¹, Diya A. El-Badry², Gamal A. El-Sisy¹, Walid S. El-Nattat^{1*}, Amal M. Abo Almaaty¹¹Animal Reproduction and AI Department, Veterinary Division, National Research Center, Dokki, Giza, Egypt²Department Artificial Insemination and Embryo Transfer, Animal Reproduction Research Institute, Agriculture Research Center, Giza, Egypt

ARTICLE INFO

Article history:

Received 17 Mar 2016

Received in revised form 6 Jun 2016

Accepted 7 Jun 2016

Available online 20 Jun 2016

Keywords:

Arabian stallion

Semen

Preservation

Honey

ABSTRACT

Objective: To investigate the effect of extender supplementation with different concentrations of a honey bee on post-thawed sperm motility, viability index, membrane and acrosome integrities in Arab stallion.

Methods: Five ejaculates from each of four Arabian stallions were subjected to cryopreservation with a modified INRA-82, without any supplementation (control) or supplemented with 1%, 2%, 3%, 4% and 5% honey bee. After thawing, all samples were maintained at 37 °C, while analyses were performed at 0, 1, and 2 and 3 h. Sperm motility percentage, viability index percentage, membrane integrity percentage and acrosome integrity percentage of each sample were determined by conventional laboratory methods.

Results: Relative to the control group, supplementation with honey (2%, 3% and 4% significantly improved ($P < 0.01$ at least) post-thaw sperm motility, viability index ($P < 0.001$ at least) and had a positive effect on membrane integrity and intact acrosome percentage ($P < 0.001$ at least) at 0, 1, 2 and 3 h post-thawing. For all semen parameters, the lower concentration of honey (1%) and higher concentration (5%) did not show significant differences ($P > 0.05$) compared with the control.

Conclusion: Honey bee supplementation in Arab stallion semen extenders provided a better protection of sperm parameters against cryopreservation injury, in comparison to the control groups.

1. Introduction

There are worldwide renewed research interest, in improving reproductive performance in livestock artificial insemination (AI) programs and maximizing the use of AI sires with valuable genomes. Such progress can be made through improvement of semen collection, analysis, processing and cryopreservation [1]. Semen cryopreservation induces certain detrimental structural effects on spermatozoa as a result of its exposure to thermal, mechanical, chemical osmotic and oxidative stress [2] during freeze-thawing. These changes resulted in reduced sperm motility, plasma membrane functionality and acrosome integrity [3], leading to diminished spermatozoa fertilizing ability [4]. There is an international interest concerning the application of

natural medical sources in animal production fields. The beneficial effects of honey bee (HB) in reproductive health protection have been strongly evidenced by many authors [5–7]. This effects are mainly attributed to its nutrient-rich content, e.g., sugars such as fructose and glucose; minerals such as magnesium, potassium, calcium, sodium chloride, sulphur, iron and phosphates, caffeic acid, caffeic acid phenethyl ester (CAPE) and flavonoid glycones; as well as vitamins B1, B2, C, B6, B5 and B3 [5,6]. HB has a potent antioxidant and antibacterial properties [7,8]. HB flavonoids possess free radicals (FR) scavenging activity, thereby inhibiting FR-induced DNA damage [9]. Olayemi *et al.* concluded that the addition of honey to egg yolk extender improve the motility and live dead ratio and viability of liquid storage goat semen [10]. Fakhridin and Alsaadi concluded that supplementation of honey bee (10%) to cryoprotectant solution show enhancement sperm parameter post-thawing in man [11]. Ogretmen and Inananobserved that using the honey in cryomedia is effective for cryopreservation especially about hatching success of egg fertilized by the frozen-thawed sperm of common carp [12]. El-Sheshtawy *et al.* mentioned that addition of 10% HB solution

*Corresponding author: Walid S. El-Nattat, Animal Reproduction and AI Dept., Veterinary Research Division, 13 National Research Centre, Dokki, Giza, Egypt.

Tel: +202 33371635

Fax: +202 37601877.

E-mail: elnattat@gmail.com

Peer review under responsibility of Hainan Medical College.

to cattle bull semen extender improved sperm motility in chilled and frozen semen and also improved the conception rate [13]. Jerez-Ebensperger *et al.* reported a significant high ram sperm quality values when extender contained pasteurized egg yolk and honey at 0 h and 2 h after thawing [14]. Jerez-Ebensperger *et al.* concluded that the addition of honey to ram extender reduces sperm deterioration when stored at 4 °C [15]. Up to date, there is a lack of data concerning the effect of the addition of honey to semen extender on post-thawing semen quality of Arab stallion. The present study aimed to determine the effects of addition of honey to semen extender on post-thawing sperm quality in Arab stallion.

2. Materials and methods

2.1. Preparation of extender

Modified INRA-82 [mINRA-82] described by El-Badry *et al.* [16] was used as base and control extender. This extender consists of 25 g/L glucose monohydrate, 1.5 g/L lactose monohydrate, 1.5 g/L raffinose pentahydrate, 0.4 g/L potassium citrate monohydrate, 0.3 g/L sodium citrate dihydrate, 4.76 g HEPES, pH 7.0, 500 mg/L gentamycin, 0.035% SDS and 0.15% skim milk. Aliquots of mINRA-82 extenders were supplemented with different concentrations of bee honey.

2.2. Honey bee mINRA-82 (*Apis mellifera lamarckii*) extender (HEMI)

Honey solution was prepared by adding 1 mL honey to 9 mL bi-distilled water to obtain a honey solution of 10% concentration. This solution is added to mINRA-82 aliquots in concentrations of 0.5/4.5 [1% honey enriched mINRA-82 (HEMI 1%)], 1/4 (HEMI 2%), 1.5/3.5 (HEMI 3%), 2/3 (HEMI 4%), 2.5/2.5 (HEMI 5%) mL (v/v) of honey/mINRA-82 to obtain a final volume of 5 mL in each tube [13].

2.3. Animals and semen collection

On a once weekly collection schedule, five ejaculates per stallion were obtained from four Arabian stallions, aged 8–14 years, and individually housed at Police Academy stud, Cairo, Egypt. At the time of collection, early in the morning, a mare in estrus was used as a mounted animal. Semen was collected using a lubricated and pre-warmed (45–50 °C) Colorado model artificial vagina with an inline filter to separate the gel fraction.

2.4. Processing of semen

Immediately following collection, the gel-free portion of the ejaculate was evaluated for volume and progressive motility, and concentration was determined with a hemocytometer. Only ejaculates with at least 60% progressively motile sperm and 2.5×10^8 sperm cell/mL were used for freezing. The semen was extended 1:1 (semen: extender) in INRA-82 extender that had been warmed to 38 °C. The diluted samples were placed into 15 mL tubes and centrifuged for 10 min at $400 \times g$ [17]. At least 95% of the supernatant was

removed [18] and each pellet was re-suspended with modified INRA-82 (containing 5% glycerol and 15% egg yolk) with honey (1%, 2%, 3%, 4%, 5%) and without honey (0%, control) to a final sperm concentration of 1.0×10^8 motile sperm/mL. Each aliquot was cooled slowly to 5 °C over 1 h under aerobic conditions and then incubated at 5 °C for 30 min [19]. The extended semen was drawn into 0.5 mL straws (Minitube, Germany) and sealed thermally and placed 4 cm above liquid nitrogen in the vapor phase in foam box for 10 min before being plunged into the liquid phase [20]. The straws were then stored in goblets on canes and kept immersed in liquid nitrogen. For thawing, two straws per treatment were warmed in a water bath at 38 °C for 30 s.

2.5. Evaluation of frozen-thawed semen

Spermatozoa motility was examined and recorded using a pre-warmed stage of phase contrast microscope ($\times 200$) just after 0, 1, 2 and 3 h post-thawing for frozen-thawed semen. The post-thawing viability indices were estimated according to Milovanov [21]. The percentage of HOS-positive cells and acrosomal integrity in each sample was determined according to Nie and Wenzel [22] and Wells and Awa [23], respectively.

2.6. Statistical analysis

One way analysis of variance and Duncan's multiple range tests (using SPSS program version 16.0) were done for the obtained data of frozen-thawed semen qualities after transformation of percentages to their corresponding arcsin values [24]. $P < 0.05$ was considered as statistically significant.

3. Results

The effects of different concentrations of honey in extenders on post-thawing Arab stallion sperm motility at different time are reported in Table 1. There were significant differences among different concentrations in total motility at 0, 1, 2 and 3 h post-thawing. A significant increase in the percentage of total motile sperm was observed in the presence of 2%, 3% and 4% bee honey as compared with control (0%). Enrichment of the basic extender (mINRA-82) for horse with the bee honey (HEMI 3%) had significantly ($P < 0.01$ at least) enhanced post-thawing (0, 1, 2 and 3 h) sperm motility percentage in comparison with the control and HEMI (1% and 5%) as shown in Table 1. While, the enrichment with HEMI 2% (at 0, 1 and 2 h post-thawing) and HEMI 4% (at 0, 1, 2 and 3 h after thawing) have non-significantly differed from the HEMI 3% results at 0, 1, 2 and 3 h post-thawing as shown in Table 1.

In the same consent, there were also significant differences among extenders in viability index, membrane integrity (HOS-positive cells) and acrosome integrity at different times post-thawing as shown in Table 2. The enrichment of the mINRA-82 with the bee honey (HEMI 3%) showed the highest significant ($P < 0.001$ at least) mean records concerning the viability index, the HOS-positive cells and the acrosome integrity in comparison with the means recorded for the control and HEMI (1%, 2% and 5%) as shown in Table 2. A non-significant ($P < 0.001$ at least) difference was found between the mean records of the HEMI 3% and 4% as revealed in Table 2.

Download English Version:

<https://daneshyari.com/en/article/3453440>

Download Persian Version:

<https://daneshyari.com/article/3453440>

[Daneshyari.com](https://daneshyari.com)