



Document heading doi: 10.1016/S2305-0500(14)60009-9

## Correlation between embryo morphology and development and chromosomal complement

Vy Phan<sup>\*</sup>, Eva Littman, Dee Harris, Antoine La

Red Rock Fertility Center, 6420 Medical Center ST, STE 100, Las Vegas, Nevada, USA

### ARTICLE INFO

#### Article history:

Received 21 February 2014  
 Received in revised form 25 February 2014  
 Accepted 25 February 2014  
 Available online 20 June 2014

#### Keywords:

Aneuploidy  
 Blastomere biopsy  
 Cleavage rate  
 Chromosomal abnormalities  
 Embryo fragmentation  
 Embryo morphology  
 CGH

### ABSTRACT

**Objective:** To analyze the correlation between embryo morphology and the chromosomal status using the array comparative genomic hybridization [array comparative genomic hybridization (a-CGH)] technique for screening 23 chromosome pairs in a single blastomere biopsy from Day 3 embryos. **Methods:** One thousand five hundred and fifty seven embryos were included from 203 cycle ICSI patients undergoing preimplantation genetic screening. The 23 chromosome pairs were analyzed by blastomere biopsy from day 3 embryos using a-CGH array method. Embryo development rate, fragmentation rate and chromosome status of the analyzed blastomeres were recorded and correlated with the aCGH results. **Results:** The incidence of chromosomal abnormalities was significantly higher in slow- and fast cleaving embryos at day 3 after insemination. The incidence of fragmentation and the type of fragmentation was associated with an increased incidence of chromosomal abnormalities. The symmetry of the blastomeres also correlated with the aneuploidy rates. **Conclusions:** Embryo development rate and morphological parameter such as degree, type of fragmentation and the symmetry of the blastomeres to a large extent reflect the cytogenetic status of the embryo and thus are important in the selection of embryos with the highest implantation potential.

## 1. Introduction

One of the key factors for a successful outcome in an assisted reproductive program is the selection of embryos with the highest developmental potential. For many years embryos have been selected based on parameters considered important quality indicators, such as fragmentation, cell number and cell size<sup>[1-7]</sup>.

The development and implementation of techniques for preimplantation genetic diagnostic programs have made it possible to assess the chromosomal constitution without destroying the embryo. It has been suggested that pronuclear morphology could be indicative of embryo quality<sup>[1,8]</sup>. Using FISH technique for pre-implantation genetic screen for

5–7 chromosomes, previous studies showed that embryos containing unevenly sized blastomeres have an increased aneuploidy rate<sup>[9,10]</sup>. Further, there is evidence that growth retardation in addition to accelerated cleavage could be an indication of chromosomal abnormalities<sup>[11-13]</sup>. Other studies have demonstrated increased chromosomal abnormality rates with increased degree of fragmentation or poor embryo morphology<sup>[11-13]</sup>.

Most studies published have been based on the technique of Fluorescence In Situ Hybridization (FISH) results. This form of evaluation consists of testing 5 to 12 chromosomes from either single or dual cell biopsy in an attempt to predict the chromosomal status of the whole embryo<sup>[10,12,13]</sup>. Therefore, the information obtained is not always representative of the real chromosomal status. The adaptation of a-CGH to single cells has allowed the study of the full karyotype of blastomeres, thus identifying the true level of aneuploidy in cleavage-stage embryos, which was then reported to affect 75% of them<sup>[14-17]</sup>.

<sup>\*</sup>Corresponding author: Vy Phan, Red Rock Fertility Center, 6420 Medical Center ST, STE 100, Las Vegas, Nevada, USA  
 Tel: (1) 702 262 0079  
 Fax: (1) 702 685 6910;  
 E-mail: kxanhvyphan@yahoo.com

The aim of this study was to analyze the correlation between embryo morphology and the chromosomal status using the a-CGH technique for screening 23 chromosome pairs in a single blastomere biopsy from Day 3 embryos.

## 2. Materials and methods

### 2.1. Patient population, embryo culture and biopsy

203 patients undergoing *in vitro* fertilization (IVF) treatment and preimplantation genetic screening (PGS) with aCGH at Red Rock Fertility Center were included in this study. The study was conducted after obtaining the Institutional Review Board's approval. The average maternal age of patients was 34.7 years (range 29–41 years). Patients underwent one of the following ovarian stimulation protocols; luteal phase Lupron suppression (Leuprolide acetate; TAP Pharmaceuticals, Lake Forest, IL) with or without oral contraceptive pretreatment; gonadotropin-releasing hormone (GnRH) antagonist prevention of premature ovulation with cetrorelix (Cetrotide; EMD Derono, Rockland, MA) or ganirelix (Organon USA, Roseland NJ). In the antagonist protocol, the GnRH antagonist was added when a lead follicles measured  $\geq 14$  mm. Controlled ovarian hyperstimulation was performed with human menopausal gonadotropin (Menopur; Ferring Pharmaceuticals, Parsippany, NJ), recombinant luteinizing hormone (LH, Luveris, EMD Serono), and/or recombinant FSH (Follistim, Organon USA; Gonal-F, EMD Serono). Cycles were monitored with serum estradiol levels and transvaginal ultrasounds. When at least 2 follicles measured  $\geq 18$  mm, 5 000–10 000 units of urinary hCG (Novarel; Ferring Pharmaceuticals) were administered subcutaneously. Ultrasound-guided oocyte retrieval was performed 36 hours after hCG administration.

All mature oocytes were fertilized by intra-cytoplasmic sperm injection (ICSI) method. Embryos were cultured using Global media (LifeGlobal) with 10% Serum Substitute Supplement (SSS) (Irvine Scientific) under triple gas incubator (6.5% CO<sub>2</sub>; 5% O<sub>2</sub> and 88.5% N<sub>2</sub>).

A total of 1 257 embryos were biopsied on day 3 of embryo development and underwent aneuploidy screening with aCGH. After biopsied, the embryos were culture until day 5 or day 6 of development. Euploid embryos were either transferred to the uterus or frozen for future use.

### 2.2. Embryo scoring

Oocytes were checked for the presence of pronuclei and polar bodies 16–18 hours after ICSI. Fertilized oocytes were cultured and scored 66–68 hours after insemination for: cell

number; degree of fragmentation (without fragmentation, less than 5%, 6% to 15 %; 16% to 30% and more than 30% fragmentation); localization of fragments (local or dispersed); equally or unequally sized blastomeres).

Biopsy procedures were carried out on day 3 (66–68 hours after insemination). One blastomere was gently aspirated with the use of a biopsy pipette. After blastomere biopsy, embryos were thoroughly rinsed and transferred to a new dish of Global media with 10% SSS and cultured to day 5 and day 6. The biopsied blastomere was transferred to the tubes and sent to a Genetics laboratory for chromosome evaluation by a-CGH. Embryos marked as euploid were chosen for transfer or frozen.

### 2.3. Statistical analysis

Data was analyzed by *chi*-square analysis and relative risk test (*RR* test).

## 3. Results

1 257 cleavage-stage embryos were biopsied and analyzed for aneuploidy. A total of 783/1257 (62.3%) were aneuploidy and 474/1257 (37.7%) were euploid. All embryos were observed until the end of day 6 for developmental progress. 572 blastocysts developed from biopsied embryos, of which 257 (32.8%) were developed from aneuploidy embryos and 315 (66.5%) were developed from euploidy embryos. The competence to develop to blastocyst stage was decrease 2 times in aneuploidy embryos compare to euploidy embryos (32.8% vs. 66.5%); *RR* = 2; *P* < 0.001 (Table 1).

**Table 1**

Development of biopsied embryos to blastocyst stage.

Development of D3 embryos	Aneuploid	Euploid	<i>RR</i>
Slow/ Arrested	526(67.2%)	159(33.5%) <sup>a</sup>	1
Blastocysts	257(32.8%)	315(66.5%) <sup>b</sup>	2
Total	783	474	-

a vs. b *p* < 0.001

Figure 1 shows the results of 1 257 embryos which underwent tested for aneuploidy. The results were analyzed for each cellular stage in details, the lowest incidence of chromosomal abnormalities was found in embryo with 8 cells (53.9%) and the highest was found in embryos with 4–5 cells (87.2%). In human *in-vitro* fertilization, embryos usually develop to 8 cell stage at 54–72 hours after fertilization. In our study, we assessed the embryos at 66–68 hours after fertilization. Our results show that slow developed embryos that have 4–6 cells have significantly higher aneuploidy rate of 83.1%, nearly 1.5 times higher than embryos which have

Download English Version:

<https://daneshyari.com/en/article/3453726>

Download Persian Version:

<https://daneshyari.com/article/3453726>

[Daneshyari.com](https://daneshyari.com)