



HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: <http://ees.elsevier.com/apjtm>Original research <http://dx.doi.org/10.1016/j.apjtm.2016.01.004>Effect of lentivirus-mediated integrin $\alpha V\beta 3$ -shRNA on tumor growth of mice with lung cancer xenograftsTao Liang¹, Yong-Fu Ma², Jian Chu¹, Dao-Xi Wang¹, Yang Liu^{2*}¹Department of Thoracic Surgery, Second Artillery General Hospital of Chinese PLA, Beijing 100088, China²Department of Thoracic Surgery, Chinese PLA General Hospital, Beijing 100853, China

ARTICLE INFO

Article history:

Received 15 Nov 2015

Received in revised form 20 Dec 2015

Accepted 30 Dec 2015

Available online 13 Jan 2016

Keywords:

Lung cancer
Integrin $\alpha V\beta 3$
Xenografts
Angiogenesis
Apoptosis
Invasion

ABSTRACT

Objective: To study the effect of lentivirus-mediated integrin $\alpha V\beta 3$ -shRNA on tumor growth of mice with lung cancer xenograft.**Methods:** Lung cancer tissue, paracancer tissue and normal tissue were collected and integrin $\alpha V\beta 3$ expression was detected; BALB/c nude mice were selected, divided into integrin $\alpha V\beta 3$ knockdown group (KD group) and negative control group (NC group), and inoculated with cells stably infected by integrin $\alpha V\beta 3$ -shRNA lentivirus and cells stably infected by negative control-shRNA lentivirus, respectively, the growth of tumor tissue was continuously observed, and the number of apoptosis cells as well as the expression of angiogenesis, apoptosis and invasion genes in tumor tissue were detected.**Results:** mRNA content and protein content of integrin $\alpha V\beta 3$ in lung cancer tissue were significantly higher than those in paracancer tissue and normal tissue; increasing trend of tumor tissue volume of KD group was weaker than that of NC group, and tumor volume at various points in time of KD group was lower than that of NC group; mRNA contents and protein contents of VEGF, FGF, EGF, Bcl-2, MMP-9, MMP-12 and MMP-13 in tumor tissue of KD group were lower than those of NC group, and apoptosis index as well as mRNA content and protein content of Bax were higher than those of NC group.**Conclusions:** The expression of integrin $\alpha V\beta 3$ increases in lung cancer tissue, and lentivirus-mediated integrin $\alpha V\beta 3$ -shRNA can inhibit tumor growth of mice with lung cancer xenografts.

1. Introduction

Lung cancer is one of the most common malignant tumors worldwide and death cases caused by it rank first in cancer-related deaths in China. Non-small cell lung cancer is the most common pathological type of lung cancer, and the cancer cells have strong ability of proliferation and invasion, which directly causes high recurrence rate and distant metastasis rate in non-small cell lung cancer patients after surgical resection or chemotherapy, thus leading to poor long-term prognosis of the disease [1,2]. At present, regulatory mechanisms of lung cancer cell proliferation and invasion haven't been fully elucidated. Integrins are a family of molecules discovered in recent years

that are closely related to cell adhesion, proliferation, invasion and angiogenesis, are located in cell membrane and play a role in the form of membrane receptors. Integrin $\alpha V\beta 3$ has been confirmed to participate in the clinical and pathological process of osteosarcoma, colon cancer and other malignant tumors [3,4], but whether the molecule is involved in the development of lung cancer has not been reported. In the following research, the effect of lentivirus-mediated integrin $\alpha V\beta 3$ -shRNA on tumor growth of mice with lung cancer xenograft was analyzed.

2. Materials and methods

2.1. Materials

2.1.1. Clinical and experimental subjects

Clinical specimens were lung cancer tissue, paracancer tissue and normal tissue, and tissue was from 100 cases of patients who received radical resection of lung cancer in our hospital from May 2012 to April 2015. Lung cancer cell lines A549 were

*Corresponding author: Yang Liu, Thoracic Surgery Department, Chinese PLA General Hospital, No. 28, Fuxing Road, Haidian District, Beijing 100853, China.

Tel: +86 13501009331

E-mail: 27886870@qq.com

Peer review under responsibility of Hainan Medical College.

Foundation project: It is supported by Surface Project of National Natural Science Foundation of China (No 81573026).

purchased from the cell bank of Chinese Academy of Sciences, and BALB/c nude mice were purchased from Shanghai Slac Laboratory Animal Company.

2.1.2. Reagents and consumables

Serum and media for cell culture were from Gibco Company and consumables for culture were from Nest Company; RNA extraction and PCR amplification kits were from Beijing Tiangen Company, immunoblot kits and related antibodies were from US Sigma Company, and TUNEL kits were from US Roche Company.

2.2. Experimental methods

2.2.1. Cell culturing and transfecting methods

Lung cancer A549 cell lines were conventionally recovered and cultured, when cell density reached about 70–80%, 0.125% trypsin was used for digestion and sub-culture, and sub-cultured cells were inoculated in Petri dishes and infected with integrin $\alpha V\beta 3$ -shRNA lentivirus and negative control-shRNA lentivirus, respectively; four days after infection, 2 $\mu\text{g}/\text{mL}$ puromycin was added for pressure screening, then media were replaced every 3 d to expand the culture, and cells stably infected by integrin $\alpha V\beta 3$ -shRNA lentivirus and negative control-shRNA lentivirus were obtained, respectively.

2.2.2. Establishment of mouse models with lung cancer xenografts

Sixty BALB/c nude mice were divided into integrin $\alpha V\beta 3$ knockdown group (KD group) and negative control group (NC group), each group with 30, and 200 μL tumor cell suspension with density of $1 \times 10^8/\text{mL}$ was subcutaneously injected in necks of two groups. KD group were inoculated with cells stably infected by integrin $\alpha V\beta 3$ -shRNA lentivirus, and NC group were inoculated with cells stably infected by negative control-shRNA lentivirus.

2.2.3. Detection of tumor growth

8 d, 12 d, 16 d, 20 d and 24 d after tumor cell inoculation, diameters of xenografts were measured in three directions and marked as a, b and c, respectively, and tumor tissue volume $V = \pi abc/6$.

2.2.4. Tumor tissue collecting and processing methods

Twenty-four days after tumor cell inoculation, mice were killed and anatomized, obtained tumor tissue was washed with normal saline and equally divided into two, one was frozen in liquid nitrogen and then preserved at -80°C , and the other was fixed in formalin.

2.2.5. Detecting methods of gene expression in tumor tissue

Clinical lung tissue specimens and mouse tumor tissue preserved at -80°C were taken, RNA extraction kits and protein extraction kits were used to extract RNA specimens and protein specimens, respectively, then PCR reaction and immunoblot test were performed, respectively, and mRNA contents and protein contents of related genes were detected.

2.2.6. Detecting methods of cell apoptosis index

Tumor tissue fixed in formalin was taken and made into paraffin slices, then TUNEL kits were used for staining, all

procedures were conducted in accordance with the kit instructions, finally DAB was used for color development, 200 cells were observed under high power lens, cells with tan nuclei were taken as positive cells, the number of positive cells was counted, and the percentage of positive cells was calculated and used as apoptosis index.

2.3. Statistical methods

SPSS21.0 was used for data processing, comparison of measurement data between two groups was by *t* test, and differences were considered to be statistically significant at a level of $P < 0.05$.

3. Results

3.1. Integrin $\alpha V\beta 3$ expression in lung cancer tissue and paracancer normal tissue

mRNA content and protein content of integrin $\alpha V\beta 3$ in lung cancer tissue were significantly higher than those in paracancer tissue and normal tissue, and differences of pair wise comparison were statistically significant ($P < 0.05$) (Table 1).

3.2. Dynamic change of tumor tissue volume

After tumor cell inoculation, tumor tissue volume of both groups showed increasing trend, the increasing trend of tumor tissue volume of KD group was weaker than that of NC group, tumor tissue volume at various points in time of KD group was lower than that of NC group, and differences between two groups were statistically significant ($P < 0.05$) (Table 2).

3.3. Expression of angiogenesis molecules in tumor tissue

mRNA contents and protein contents of VEGF, FGF and EGF in tumor tissue of KD group were significantly lower than those of NC group, and differences between two groups were statistically significant ($P < 0.05$) (Table 3).

3.4. Cell apoptosis in tumor tissue

Apoptosis index in tumor tissue of KD group was higher than that of NC group, mRNA content and protein content of pro-apoptosis gene *Bax* were higher than those of NC group, and mRNA content and protein content of anti-apoptosis gene *Bcl-2* were lower than those of NC group ($P < 0.05$) (Table 4).

Table 1

Comparison of integrin $\alpha V\beta 3$ expression in lung cancer tissue, paracancer tissue and normal tissue.

Indexes	Specimen no.	Integrin $\alpha V\beta 3$ mRNA content	Integrin $\alpha V\beta 3$ protein content
Lung cancer tissue	100	278.75 \pm 32.38	341.22 \pm 39.58
Paracancer tissue	100	156.55 \pm 18.48	176.64 \pm 20.32
Normal tissue	100	100.00 \pm 14.96	100.00 \pm 11.74
<i>F</i>		17.686	23.692
<i>P</i>		<0.05	<0.05

Download English Version:

<https://daneshyari.com/en/article/3455248>

Download Persian Version:

<https://daneshyari.com/article/3455248>

[Daneshyari.com](https://daneshyari.com)