



HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: <http://ees.elsevier.com/apjtm>Original research <http://dx.doi.org/10.1016/j.apjtm.2016.01.007>

Therapeutic effect of oridonin on mice with prostate cancer

Ming Ming^{1,2,#}, Feng-Yin Sun^{1,#}, Wen-Tong Zhang¹, Ji-Ke Liu^{1,3*}¹Department of Pediatric Surgery, Qilu Hospital of Shandong University, Jinan, Shandong, 250000, China²Department of Pediatric Surgery, Taian City Central Hospital, Taian, Shandong, 271000, China³Department of Pediatric Surgery, Liaocheng Hospital, Liaocheng, Shandong, 252000, China

ARTICLE INFO

Article history:

Received 15 Nov 2015

Received in revised form 20 Dec 2015

Accepted 30 Dec 2015

Available online 11 Jan 2016

Keywords:

Oridonin

Prostate cancer

RM-1 cells

Caspase-3

ABSTRACT

Objective: To investigate the therapeutic effect and the related mechanism of oridonin on mice with prostate cancer.**Methods:** Sixty BALB/C male nude mice were selected. A model of RM-1 cell transplantation tumor of prostate cancer was built by the subcutaneous inoculation of RM-1 cells. After that, those 60 experimental mice were randomly divided into groups A, B and C. Each group had 20 mice. Mice in group A were treated with 0.2 mL of normal saline (0.9%) by intraperitoneal injection once a day; mice in group B received intraperitoneal injection of 1.875 mg/mL of oridonin once a day; and mice in group C received intraperitoneal injection of 7.5 mg/mL of oridonin once a day. Mice in the three groups were treated uninterruptedly for 5 weeks and were all killed. Then, tumors were excised and weighed to calculate their growth inhibitory rate, volume increment and anti-tumor rate. Thymus and spleen of mice in the three groups were collected to calculate the thymus and spleen index. Immunohistochemical staining was applied to observe the expression of caspase-3 in prostate cancer tissue of mice of the three groups.**Results:** The qualities and volume increment of tumors in groups B and C were significantly lower than those of group A ($P < 0.05$); the qualities and volume increment of tumors in groups C were evidently lower than those of group B ($P < 0.05$); the tumor volume increment and anti-tumor rate in group C were obviously higher than those of group B ($P < 0.05$); the thymus and spleen indexes of groups B and C were distinctly higher than those of group A ($P < 0.05$); comparison of the thymus and spleen indexes between group B and group C showed no statistical differences ($P > 0.05$). Immunohistochemical staining revealed that the caspase-3 protein in prostate cancer tissue of mice of group A expressed negatively with colorless or light-colored karyon; while the caspase-3 protein in prostate cancer tissue of mice of group B expressed positively with dark-colored karyon, centralized distribution and granular sensation; and the caspase-3 in prostate cancer tissue of mice of group C showed strong positive expression with big and darker colored karyon and dense distribution.**Conclusions:** Oridonin can inhibit the growth of RM-1 prostate cancer cells effectively and have great therapeutic effects on RM-1 cell transplantation tumor of prostate cancer.

1. Introduction

Prostate cancer, which is a common malignant disease in male, refers to a kind of epithelial malignant tumors occurring in prostate. Generally, it occurs frequently in males aged over 55

years and its morbidity increases with age [1–3]. According to some statistics, the morbidity of prostate cancer tops the list of male malignant cancers and its mortality ranks only second to lung cancer, which is severely threatening men's health [4]. The pathogenesis of prostate cancer is related to genetic factor, dietary habit and sex activity. Since the pathogenesis of the disease is concealed, the disease has progressed to its middle or advanced stage when it is diagnosed. The tumor cells have already transferred to other organs leading to the missing opportunity of radical operation [5–8]. Therefore, it is significant to find effective drugs to prolong the survival time and promote the life quality for patients. Oridonin is an

*Corresponding author: Ji-Ke Liu, PhD, Attending Physician, Department of Pediatric Surgery, Qilu Hospital of Shandong University, No. 107, Wenhua Xi Road, Jinan, Shandong, 250000, China.

Tel: +751 82166741

E-mail: liujuk1983@sina.com

Peer review under responsibility of Hainan Medical College.

These authors contributed equally to this work.

effective active ingredient extracted from natural *Rab-dosia rubescens*. It possesses the abilities to eliminate various cancer cells and fight against cancers and tumors effectively [9]. In this study, in order to investigate the therapeutic effect and the related mechanism of oridonin on mice with prostate cancer, 60 BALB/C male nude mice were selected to build a model of RM-1 cell transplantation tumor of prostate cancer, and treated with oridonin in different concentrations. Now the research results were reported as follows.

2. Materials and methods

2.1. Experimental animals

Sixty 6-month-old BALB/C male nude mice (clean animal) with weights of 18–20 g were chose. They were purchased from Beijing Lihua Experimental Animal Technology Co. Ltd. Those mice took food and drank freely under the room temperature of $(23 \pm 3) ^\circ\text{C}$. The disposition of the experimental animals was strictly abided by the laboratory animal administration rules.

2.2. Drugs and instruments

RM-1 prostate cancer cell strains were provided by Yanyu Biotechnology Co. Ltd. (Shanghai); oridonin was bought from Xi'an Hao Xuan Biological Technology Co. Ltd.; MTT and Trizol were from Baotaike Company; IMDM nutrient medium was from Gibco (USA); Olympus invert microscope was from Japan; Bio-rad microplate reader was from USA and Shimadzu ultraviolet spectrophotometer was from Australia. Oridonin was diluted by normal saline into two concentration gradients, 1.875 and 7.5 mg/mL respectively.

2.3. Model establishment

The revived RM-1 prostate cancer cells were inoculated in the IMDM nutrient medium and then it was cultured and generated in a 5% CO₂ humidity incubator under the temperature of 37 °C. The logarithmic growth phase cells were collected and prepared into cell suspension with the concentration of 10⁷ cells/mL.

2.4. Animal grouping

In order to establish the model of RM-1 cell transplantation tumor of prostate cancer, 0.2 mL of the prepared cell suspension of RM-1 prostate cancer cells was inoculated subcutaneously in the axilla of those experimental mice. The tumors formed successfully in all the experimental mice. After inoculation, those mice were randomly divided into groups A, B and C. Each group had 20 mice. Mice in group A were treated with 0.2 mL of normal saline (0.9%) by intraperitoneal injection once a day; mice in group B received intraperitoneal injection of 1.875 mg/mL of oridonin once a day; and mice in group C received intraperitoneal injection of 7.5 mg/mL of oridonin once a day. Mice in the three groups were treated uninterruptedly for 5 weeks.

2.5. Observation

During the first, third and fifth weeks after inoculation, vernier caliper was used to measure the tumor volume and draw the tumor growth curves. Then, those mice were all killed after

finishing the treatment. And the tumors were excised and weighed to calculate their growth inhibitory rate, volume increment and anti-tumor rate. Thymus and spleen of mice were collected to calculate the thymus and spleen index. Immunohistochemical staining was applied to observe the expression of caspase-3 in prostate cancer tissue of mice of the three groups.

2.6. Statistical arrangement

SPSS 10.0 was used to analyze research data and mean \pm SD was applied to express the tumor volume, volume increment and anti-tumor rate, etc. One-way ANOVA was used for intergroup comparison. Differences indicated statistical significances when $P < 0.05$.

3. Results

3.1. Comparison of tumor growth inhibition rates in different treatment periods of three groups

Since tumors in mice of group A grew continuously in all treatment periods, the tumor growth inhibition rate of group A was ignored. The tumor growth inhibition rates of group C for 1, 3 and 5 weeks were all higher than those of group B. Comparative differences between groups had statistical significances ($P < 0.05$) (Table 1).

3.2. Comparison of qualities, volumes, volume increments of tumors and anti-tumor rates of three groups

The quality and volume increment of tumors in groups B and C were significantly lower than those of group A, and the comparative differences between groups had statistical significances ($P < 0.05$); the qualities and volume increment of tumors in groups C were evidently lower than those of group B and the tumor volume increment and anti-tumor rate in group C were obviously higher than those of group B, and the comparative differences between groups also showed statistical significances ($P < 0.05$) (Table 2).

3.3. Comparison of thymus and spleen index of mice in three groups

The thymus and spleen indexes of groups B and C were distinctly higher than those of group A, and comparative differences between groups had statistical significances ($P < 0.05$); while comparison of the thymus and spleen indexes between group B and group C showed no statistical differences ($P > 0.05$) (Table 3).

Table 1

Comparison of tumor growth inhibition rates in different treatment periods of three groups.

Group	n	One week	Three weeks	Five weeks
A	20	–	–	–
B	20	19.10 \pm 9.48	30.09 \pm 1.19	36.47 \pm 3.30
C	20	52.16 \pm 2.05 [#]	63.07 \pm 1.71 [#]	69.60 \pm 3.40 [#]

Compared with group B, [#] $P < 0.05$.

Download English Version:

<https://daneshyari.com/en/article/3455253>

Download Persian Version:

<https://daneshyari.com/article/3455253>

[Daneshyari.com](https://daneshyari.com)