



# Urinary Tract Infections Among the Elderly in Benin City, Nigeria

Richard Omoregie<sup>1,2\*</sup>, Isaac Ohiohrenuan Igbarumah<sup>2</sup>,  
Christopher Aye Egbe<sup>2</sup>, Helen Ogefere<sup>3</sup>

<sup>1</sup>School of Medical Laboratory Sciences, University of Benin Teaching Hospital, Benin City, Edo State, Nigeria

<sup>2</sup>Department of Medical Microbiology, University of Benin Teaching Hospital, Benin City, Edo State, Nigeria

<sup>3</sup>Department of Medical Laboratory Science, School of Basic Medical Sciences, University of Benin, Benin City, Nigeria

Received: October 11, 2010 Revised: October 29, 2010 Accepted: November 18, 2010

The elderly have an increased risk of developing urinary tract infection (UTI). This study aimed to determine the prevalence of UTI among elderly ( $\geq 60$  years) patients in Benin City, Nigeria. Clean-catch midstream urine was collected from 943 (362 males and 581 females) elderly patients with signs and symptoms of UTI. Significant microbial isolates were identified in the urine specimens and an antimicrobial susceptibility test was performed on bacteria isolates using standard techniques. Male sex was a significant risk factor for acquiring UTI in the elderly (odds ratio=6.584; 95% confidence interval=4.081, 10.62;  $p<0.0001$ ). There was an inverse relationship between age and prevalence of UTI, as the prevalence of UTI significantly declined with increasing age ( $p=0.023$ ). *Escherichia coli* was the most common uropathogen (36.79%) in both sexes of elderly patients. Bacterial isolates were poorly susceptible to antibacterial agents used. An overall prevalence of 11.03% of UTI was observed among the elderly. Male sex showed an approximately 4–11-fold increased risk of acquiring UTI. The prevalence of UTI among the elderly decreases with increasing age. Rational use of antibiotics is advocated to stem the tide of high antibacterial resistance.

**Key Words:** elderly; Nigeria; prevalence; urinary tract infection

## Introduction

Urinary tract infections (UTIs) are a common condition causing individuals to seek medical care.<sup>1</sup> Among the populations at special risk for UTI are older adults.<sup>2</sup> UTI has been reported to be the second most common infection among older adults living in the community setting, as well as the leading site of infection in adults in long-term care facilities.<sup>2,3</sup>

*Escherichia coli* is reported as the most prevalent pathogen causing UTI.<sup>1,4–6</sup> However two

recent reports from our institution indicated that *Staphylococcus aureus* is the most prevalent organism.<sup>6,7</sup> There has also been an increase in the emergence of resistant uropathogens to antimicrobial agents among the elderly.<sup>2</sup> There are no reports of UTI among the elderly in our institution, and to the best of our knowledge, in Nigeria. Therefore, this study aimed to determine the prevalence of UTI among the elderly. Etiological agents and their susceptibility profiles were also determined.

\*Corresponding author. School of Medical Laboratory Sciences, University of Benin Teaching Hospital, P.M.B. 1111, Benin City, Edo State, Nigeria.  
E-mail: richyomos@yahoo.com

## Materials and Methods

### Study population

A total of 943 patients ( $\geq 60$  years) with signs and symptoms of UTI were recruited for this study. The patients were attending various clinics in the University of Benin Teaching Hospital from the 1<sup>st</sup> September, 2009 to 31<sup>st</sup> May, 2010. Verbal informed consent was obtained from all subjects prior to specimen collection. Approval for the study was given by the Ethical Committee of the University of Benin Teaching Hospital.

### Specimen collection and processing

Clean-catch midstream urine was collected from each patient using boric acid as a preservative. A loopful (0.001 mL) of well-mixed urine was streaked on blood agar and cysteine lactose electrolyte deficient medium (M6; Plasmatec Laboratories, United Kingdom). The plates were incubated aerobically for 24 hours and counts were expressed in a colony forming unit (cfu) per milliliter (mL). A count of  $\geq 10^5$  cfu/mL was considered significant to indicate UTI. Each urine sample (10 mL) was centrifuged at 2000g for 5 minutes. The supernatant was discarded and the deposit was examined microscopically at high

magnification for pus cells, red blood cells, cast, epithelial cells, crystals, yeast-like cells and *Trichomonas vaginalis*. A count of pus cells  $\geq 5$  under high power field ( $\times 40$  objective) was considered to indicate infection. UTI was diagnosed if the bacteria count, pus cells, or both were significantly high in an individual. The clinical isolates were identified by a standard microbiological method<sup>8</sup> and the paper disc antimicrobial susceptibility test for bacterial isolates was performed using the British Society for Antimicrobial Chemotherapy method.<sup>9</sup>

Statistical analysis was performed using the  $\chi^2$  test and odds ratio (OR) analysis using the statistical software INSTANT (GraphPad Software Inc., La Jolla, CA, USA).

## Results

Male sex among elderly patients was a significant risk factor for acquiring UTI ( $p < 0.0001$ ). The prevalence of UTI decreased with increasing age ( $p = 0.023$ , Table 1).

*E. coli* was the most predominant isolate in UTI among the elderly patients, including males and females. *E. coli* and *Candida albicans* were the most prevalent pathogens among females with a prevalence of 29.17% for each (Table 2).

**Table 1** Effect of gender and age on the prevalence of urinary tract infections among the elderly

| Characteristics | No. tested | No. infected (%) | OR    | 95% CI        | <i>p</i> |
|-----------------|------------|------------------|-------|---------------|----------|
| Sex             |            |                  |       |               |          |
| Male            | 362        | 80 (23.10)       | 6.584 | 4.081, 10.621 | <0.0001  |
| Female          | 581        | 24 (4.13)        | 0.152 | 0.094, 0.245  |          |
| Age (yr)        |            |                  |       |               |          |
| 60–69           | 332        | 48 (14.46)       |       |               | 0.023    |
| 70–79           | 320        | 35 (10.94)       |       |               |          |
| 80–89           | 230        | 19 (8.26)        |       |               |          |
| 90–99           | 61         | 2 (3.28)         |       |               |          |

OR: odds ratio; CI: confidence interval.

**Table 2** Prevalence of microbial agents of urinary tract infections among the elderly

| Organisms                     | Males <i>n</i> (%) | Females <i>n</i> (%) | Total <i>n</i> (%) |
|-------------------------------|--------------------|----------------------|--------------------|
| <i>Escherichia coli</i>       | 32 (39.02)         | 7 (29.17)            | 39 (36.79)         |
| <i>Klebsiella pneumoniae</i>  | 20 (24.39)         | 4 (16.67)            | 24 (22.64)         |
| <i>Enterobacter</i> species   | 1 (1.22)           | 1 (4.17)             | 2 (1.89)           |
| <i>Proteus</i> species        | 2 (2.44)           | 1 (4.17)             | 3 (2.83)           |
| <i>Providencia</i> species    | 6 (7.32)           | 0 (0.00)             | 6 (5.66)           |
| <i>Acinetobacter</i> species  | 5 (6.10)           | 0 (0.00)             | 5 (4.72)           |
| <i>Pseudomonas aeruginosa</i> | 2 (2.44)           | 0 (0.00)             | 2 (1.89)           |
| <i>Alcaligenes</i> species    | 5 (6.10)           | 0 (0.00)             | 5 (4.72)           |
| <i>Staphylococcus aureus</i>  | 8 (9.70)           | 4 (16.67)            | 12 (11.32)         |
| <i>Candida albicans</i>       | 1 (1.22)           | 7 (29.17)            | 8 (7.55)           |
| Total                         | 82 (77.36)         | 24 (22.64)           | 106 (100)          |

Download English Version:

<https://daneshyari.com/en/article/3472767>

Download Persian Version:

<https://daneshyari.com/article/3472767>

[Daneshyari.com](https://daneshyari.com)