

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Journal of Acute Disease

journal homepage: www.jadweb.orgReview article <http://dx.doi.org/10.1016/j.joad.2015.04.001>

Progress on macrophage's proinflammatory products as markers of acute endometriosis

Alicja Ziętek, Konrad Futyma^{*}, Łukasz Nowakowski, Marek Gogacz, Tomasz Rechberger

2nd Department of Gynecology, Medical University of Lublin, ul. Jaczewskiego 8, 20-954 Lublin, Poland

ARTICLE INFO

Article history:

Received 23 Mar 2015

Received in revised form 28 Mar 2015

Accepted 20 Apr 2015

Available online 10 Jul 2015

Keywords:

Endometriosis

Macrophages

Interleukins

Tumor necrosis factor

ABSTRACT

To provide the review of the macrophage activity products as pathophysiological markers of endometriosis by literature survey (PubMed, Cochrane). Immunoreactive cells and several of their synthesis products concentrations are elevated in the serum and peritoneal fluid in patients with endometriosis. The enhanced reactive proteins contributed to local inflammation and aggregation of endometriotic lesions. Immune response and immune surveillance of tissue play an important role in pathogenesis of endometriosis. Activated macrophages in peritoneal environment secrete immunoreactive cytokines which are responsible for inflammatory cascade of reactions. The immunoreactive cytokines should be a target not only as a disease marker but also as a part of therapeutic protocol.

1. Introduction

Endometriosis is characterized by hormone depending persistence and growth of endometrial tissue at ectopic sites, mostly pelvic peritoneum and organs: ovaries, rectum, urinary bladder. It affects 10%–15% of women of reproductive age and is associated with pelvic pain, dysmenorrhea, dyspareunia and infertility. Despite the fact that it is considered as one of the most frequently encountered gynecological disorders, the pathogenesis of endometriosis still remains poorly understood.

Four most popular theories were proposed to explain the pathogenesis and pathophysiology of endometriosis. However, none of them can particularly explore all the determining factors.

In 1927, Sampson postulated the retrograde menstruation theory, according to which the endometrial shed elements could pass along the fallopian tubes and reach the peritoneal cavity^[1]. Considering the gravity, the menstruation blood would thus reach, especially the ovaries, the Douglas pouch, uterosacral ligaments, rectum and urinary bladder. Another popular etiological theory of endometriosis is the celomic metaplasia. The fact of metaplasia of undifferentiated tissue could possibly explain the existence of endometriosis in distant sites. Dinelescu *et al.* came up with the hypothesis that activation of

K-ras gene may be responsible for the metaplastic process^[2]. There is also a hypothesis that the remnants of Mullerian ducts could be responsible for the etiology of the disease. Nevertheless, nowadays it is considered as highly improbable. In order to explain the distant location of endometriosis, Halban postulated the theory of lymph and blood dissemination of endometrial cells^[3]. According to this idea, the endometrial cells would spread around the organism through the lymph or blood vessel microclots. In 1987, Gleicher *et al.* postulated that endometriosis could be an autoimmune disease considering the presence of certain autoantibodies in serum^[4]. In the study of Barrier, it was not proved that there was any correlation between co-occurrence of endometriosis and immune-mediated polyarthritis, Sjogrens syndrome or systemic lupus erythematosus^[5]. Several studies accentuated the role of race, uterine obstruction, life conditions in the pathogenesis of endometriosis^[5,6]. Currently, more data support the immunological theory of endometriosis, according to which it is an inflammatory disorder recruiting cellular and humoral factors.

The aim of the present study is to review the literature for novel proinflammatory markers of endometriosis.

2. Materials and methods

Comprehensive searches in PubMed and Cochrane databases were conducted to identify studies published between 1995 and 2014 in English language with keywords: “macrophages”, “endometriosis”, “interleukins”, “tumor necrosis factor (TNF)”.

^{*}Corresponding author: Konrad Futyma, 2nd Department of Gynecology, Medical University of Lublin, ul. Jaczewskiego 8, 20-954 Lublin, Poland.
Tel: +48 81 7244268
Fax: +48 81 7244849
E-mail: futymakonrad@mp.pl
Peer review under responsibility of Hainan Medical College.

Data were extracted and initial screening of the title and abstract of all articles to exclude citations deemed irrelevant were performed.

3. Macrophages in endometriosis

Numerous studies have shown that macrophages are the most crucial inflammatory cells. The peritoneal fluid which normally appears in peritoneal cavity in the volume of 5–20 mL contains activated macrophages which determine and sustain the inflammation. Macrophages are an integral part of mononuclear phagocyte system which derives from bone marrow. After reaching the peritoneum macrophages remain in the peritoneal cavity as dendritic cells or scavenger macrophages. Among the macrophages exists a classical/inflammatory M1 population characterized with the antigen cytotoxicity, synthesizing cytokines like interleukin-12 (IL-12), and modulating ROS and nitric oxide factors. The M2 population of macrophages is responsible for enhancing the inflammatory response and also for the tissue repair. During the acute stage of endometriosis, an increasing number of activated macrophages in the peritoneal fluid and endometriotic implants is observed. Those activated macrophages are responsible for numerous growth factors while cytokines secretion responsible for proliferation, angiogenesis, inflammatory response and clinical manifestation of the disease^[7,8].

3.1. Interleukin-1

Among the secretory products of activated macrophages, interleukin-1 is an important immunomodulatory factor. IL-1 is the name standing for a group of molecules which induce the inflammation pathway and are responsible for the pro-inflammatory mechanisms. Among the IL-1 family, there are IL-1 α and IL-1 β which together with antagonist receptor (IL-1Ra) cooperate in responding to inflammation signals^[9,10]. IL-1 α and IL-1 β bind to the same receptor molecule. The third ligand of this receptor (IL-1r), the interleukin 1 receptor antagonist (IL-1Ra), acts as an inhibitor of IL-1 α and IL-1 β signaling by competing with them for binding sites of the receptor. The most popularly known initializing factors for producing IL-1 are lipopolysaccharides (the component of bacterial wall) and the C5-complex system^[9]. The immunological role of IL-1 is to stimulate macrophages for the synthesis of IL-2 with the receptor (IL-2R) and IL-6. The nature killer cells cytotoxicity is enhanced by IL-1, and lymphocyte T is enhanced for the synthesis of interferon γ . By humoral immune response, IL-1 β is stimulating B lymphocytes for producing antibodies. Increasing concentration of IL-1 in the inflammation area is a chemoattractive factor for monocytes and neutrophils. These cytokines increase the expression of adhesional factors ICAM-1, VCAM-1 and selectin E on endothelial cells, to enable transmigration of immunocompetent cells, such as phagocytes, lymphocytes and others, to sites of infection. The new colonies of monocytes are recruited due to the production of M-CSF, GM-CSF and G-CSF. Viganò *et al.* suggested that IL-1 β stimulated ICAM-1 dependent immune surveillance of shed endometrial cells in the peritoneal fluid environment. The IL-1 also affects the activity of the hypothalamus which leads to a rise in body temperature. That is why IL-1 is called an endogenous pyrogen. IL-1 also causes hyperalgesia. These are some frequent clinical symptoms

correlated with the menstrual bleeding in the patients diagnosed with endometriosis. They often present the peritonitis symptoms which can be incorrectly interpreted as appendicitis, extrauterine pregnancy, diverticulitis, etc^[5,9,11].

3.2. Interleukin-6

In endometriosis, macrophages produce significantly higher levels of Interleukin-6^[12]. This is both a pro-inflammatory cytokine and an anti-inflammatory myokine, encoded by IL-6 gene. IL-6 is secreted by T cells and macrophages, in response to IL-1 stimulation (in the positive feedback) and interferones, lipopolysaccharides or TNF. Osteoblasts secrete IL-6 to stimulate osteoclast formation. Smooth muscle cells in the tunica media of many blood vessels also produce IL-6 as a pro-inflammatory cytokine. As an anti-inflammatory cytokine, IL-6 is mediated through its inhibitory effects on TNF- α and IL-1. The biological activities of IL-6 are mediated by the IL-6 receptor-system which comprises two membrane proteins, the ligand binding α subunit (IL-6 R) and the signal transducing β subunit, gp130. IL-6 is an essential cytokine for differentiation of B lymphocytes into antibodies-producing cells and contributes to production of immune complexes in endometrial stromal tissue^[12,13]. In co-operation with IL-1, this cytokine promotes the production of T cells. It also applies to have an angiogenic potential (connected to IL-3) which affects the uncontrolled growth and invasion of adjacent tissue. This phenomenon of macrophage activation products enforces vascularization and angiogenic potential which is responsible for the new endometrial lesions and widespreading of the disease^[6,14]. Concerning endometriosis in the context of autoimmune diseases, IL-6 is a promoting factor for the epithelial and mesangial tissue growth so that it could be responsible for lacking of autoimmunology response of the organism. IL-6 is also an important mediator of fever and acute phase response. It is capable of crossing the blood–brain barrier and initiating synthesis of PGE2 in the hypothalamus. Endometrial stromal and epithelial cells produce IL-6 in response to hormonal and immunological activity. IL-1 α and β , TNF, interferon- γ stimulates the endometrial cell protein IL-6. It is suggested that estrogen stimulates the endometrial cell proliferation by inhibiting the production of IL-6 (which is considered to be epithelial cell inhibitor). Barrier proposed that endometriotic implants were resistant to IL-6 activity due to a low expression of IL-6R on the cell surface^[5]. There is no consensus between studies about the concentration of IL-6 in the peritoneal fluid.

3.3. Interleukin-8

Interleukin 8 is a chemokine and chemo attractant produced by macrophages. The oxidative stress correlated with local immunodeficiency is a triggering factor for releasing the levels of IL-8 (also named as CCX-8, chemokine 8). IL-8 not only induces the migration of neutrophils to the inflammation area, but also is responsible for phagocytosis, increase of intracellular calcium concentration and exocytosis. However, since years, many studies have postulated that IL-8 is also an angiogenic factor for the cells^[15,16]. Excessive neovascularization and angiogenesis are responsible for the recruitment of a new endometriotic lesions and immune surveillance of existing one. In the clinical practice by laparoscopic views, it is always noticeable that

Download English Version:

<https://daneshyari.com/en/article/3475175>

Download Persian Version:

<https://daneshyari.com/article/3475175>

[Daneshyari.com](https://daneshyari.com)