



Original Article

Perioperative parameter analysis of neonates and infants receiving laparoscopic surgery

Chia-Man Chou^{a,b,*}, Chou-Ming Yeh^c, Sheng-Yang Huang^{a,b}, Hou-Chuan Chen^a

^a Division of Pediatric Surgery, Department of Surgery, Taichung Veterans General Hospital, Taichung, Taiwan, ROC

^b Department of Medicine, National Yang-Ming University, Taipei, Taiwan, ROC

^c Department of Health, Taichung Hospital, Executive Yuan, Taichung, Taiwan, ROC

Received February 15, 2016; accepted May 11, 2016

Abstract

Background: The field of laparoscopic surgery in neonates or younger infants has benefitted from recent progress. This study aimed to determine the correlation between patient characteristics and perioperative parameters, and to explore the feasibility of laparoscopic surgery in neonates and infants.

Methods: We retrospectively collected and analyzed data on neonates and infants who received laparoscopic surgery at our institute between January 2007 and August 2015. Perioperative data, surgical outcomes, and related complications were analyzed using Spearman rank correlation coefficient.

Results: A total of 82 patients (42 male and 40 female) were included in this study. The median operative age and the median operative body weight were 2.2 months and 4.2 kg, respectively. The median operative time was 3.5 hours, and the median insufflation time was 2.0 hours. The mean intraoperative end-tidal carbon dioxide (EtCO₂) level was 37.6 mmHg, the median body temperature (BT) was 35.8°C, and the mean peak inspiratory pressure was 23.3 cmH₂O. The median follow-up duration was 23.4 months. The intraoperative BT was significantly influenced by the operative age ($p < 0.001$, $r_s = 0.52$) and body weight ($p < 0.001$, $r_s = 0.59$). The intraoperative EtCO₂ level was higher for longer operative time ($p = 0.01$, $r_s = 0.28$) and insufflation time ($p < 0.001$, $r_s = 0.39$); however, all values returned to normal when the CO₂ insufflation was stopped.

Conclusion: Laparoscopic surgery for neonates and infants can be safely performed by experienced surgeons. However, transient hypercarbia may rapidly ameliorate after CO₂ insufflation is stopped.

Copyright © 2016, the Chinese Medical Association. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: infants; laparoscopic surgery; neonates; perioperative parameter

1. Introduction

Minimally invasive surgery (MIS) in children, particularly in those under the age of 1 year, requires sophisticated

equipment and techniques and has been developed since the late 1990s.^{1–3} In neonates and infants, friable and delicate tissue, a small intra-abdominal or intrathoracic space, a steep learning curve, and distinct anesthetic deliberation for dissimilar physiological characteristics have resulted in MIS being utilized less frequently than other techniques.^{1,3–5} In addition to technological limitations and the need for precise surgical skills, MIS in neonates and infants is challenging because of their vulnerability to hypothermia and hypercarbia, which may lead to acidosis, reduced cerebral perfusion, and other unfavorable outcomes.^{2,4}

Conflicts of interest: The authors declare that they have no conflicts of interest related to the subject matter or materials discussed in this article.

* Corresponding author. Dr. Chia-Man Chou, Division of Pediatric Surgery, Department of Surgery, Taichung Veterans General Hospital, 1650, Section 4, Taiwan Boulevard, Taichung 407, Taiwan, ROC.

E-mail address: cmchou@vghtc.gov.tw (C.-M. Chou).

<http://dx.doi.org/10.1016/j.jcma.2016.05.005>

1726-4901/Copyright © 2016, the Chinese Medical Association. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Although heat loss caused by large incisions is averted in neonates and infants receiving MIS, extended surgical duration with prolonged exposure to the environment, the use of dry and cold carbon dioxide (CO₂) gas, and gas leaks from port sites or instruments all contribute to hypothermia.^{2,4} A small airway and large dead space may result in reduced gas exchange. Furthermore, a high peritoneal absorption surface per unit of weight, little peritoneal fat, and thin vessel walls leading to rapid CO₂ diffusion all contribute to hypercarbia in neonates and infants undergoing MIS.^{2,4,6} We retrospectively collected clinical data of neonates and infants who received laparoscopic surgery at our institute and analyzed their perioperative data, surgical outcomes, and related complications to clarify the safety and feasibility of MIS in such patients.

2. Methods

Between January 2007 and August 2015, 82 patients (42 male and 40 female) less than 1 year of age who received laparoscopic surgeries at our institute were included. All of the neonates and infants received laparoscopic-related interventions for various diseases. In a typical laparoscopic procedure, the patient was placed in the supine or lithotomy position. The first transumbilical port was introduced using the open Hasson technique. A 5-mm 30° endoscope (KARL STORZ-Endoskope, Surgimed Corporation, Taichung, Taiwan) was inserted via a transumbilical port, and the positions of the other two or three work ports (5 mm or 3 mm) were determined according to operational needs. The pneumoperitoneal space was maintained using CO₂ insufflation, and the pressure was 8–10 mmHg in neonates and 10–12 mmHg in infants.

The patients' medical charts were reviewed to obtain data on general information, diagnosis, perioperative parameters, surgical outcomes, complications, and follow-up durations. The perioperative parameters were the operative age, operative body weight, operative time, CO₂ insufflation time, intraoperative end-tidal CO₂ (EtCO₂) levels, body temperature (BT), and peak inspiratory pressure (PIP). Among these factors, EtCO₂ was monitored using a capnometer, which continuously measured the concentration of CO₂ in exhaled air via the endotracheal tube connecting to the respirometer of the anesthesia machine. The data were analyzed using the Spearman rank correlation coefficient. The procedure for enrolling all patients in this research was approved by the Institutional Committee on Human Research of Taichung Veterans General Hospital (TCVGH), according to the guidelines of the Declaration of Helsinki and the International Conference on Harmonisation for Good Clinical Practice (Institutional Review Board TCVGH No. CE15260B).

3. Results

The defined diagnoses and corresponding operative procedures of the included patients are listed in Table 1. The central measurement results of the preoperative data of these patients are shown in Table 2. The operative age ranged from 1 day to 11.4 months (median

Table 1
Diagnosis, patient number, and operative procedures of patients.

Diagnosis	Operative procedure	Patient No.
Adrenal cortical adenoma	Partial adrenalectomy	1
Anorectal malformation	Laparoscopic-assisted pull-through	3
Biliary atresia	Kasai operation	6
Choledochal cyst	Excision of choledochal cyst and roux-en-y hepaticojejunostomy	9
Congenital duodenal stenosis	Duodenoduodenostomy	1
Congenital diaphragmatic hernia	Primary repair	1
	Patch repair	1
Diaphragmatic eventration	Plication of diaphragm	2
Feeding intolerance in neurologically impaired children with/without gastroesophageal reflux disease	Nissen fundoplication and gastrostomy	1
	Gastrostomy	2
Hirschsprung disease	Swenson procedure	10
Hiatal hernia	Primary repair and Toupet fundoplication	5
Infantile hypertrophic pyloric stenosis	Pyloromyotomy	5
Ileal atresia	Segmental resection and anastomosis	3
Intra-abdominal lymphangioma	Excision of intra-abdominal tumor	1
Intussusception	Surgical reduction	5
Liver cysts	Unroofing of the cyst	2
Neonatal cholestasis	Intraoperative cholangiogram	1
Nonfunctional kidney in duplex collecting system	Partial nephrectomy	5
Ovarian cyst/teratoma	Partial/total oophorectomy	4
Undescended testis	Laparoscopic-assisted orchiopexy	6
Ureteropelvic junction stenosis	Dismembered pyeloplasty	7
Vesicoureteral reflux	Ureteral reimplantation	1

2.2 months), and the operative weight ranged from 2 kg to 1 kg (median 4.2 kg). The operative time ranged from 1.0 hour to 7.5 hours (median 3.5 hours), and CO₂ insufflation time ranged from 0.2 hour to 5.0 hours (median 2.0 hours). The mean intraoperative EtCO₂ level was 37.6 mmHg, the median BT was 35.8°C, and the mean PIP was 23.3 cmH₂O. Although the EtCO₂ level increased and BT was reduced at the start of insufflation, in most patients the EtCO₂ level was maintained below 42.1 mmHg, PIP was below 26.3 cmH₂O, and BT was above 35.1°C, as shown in Table 2. Hypercarbia-related intraoperative respiratory acidosis and transient hypothermia (BT < 35°C) were observed in three and four patients, respectively. All of these

Table 2
Measurement results of preoperative data for patients.

Parameters	Median	Q1	Q3
Operative age (mo)	2.25	0.60	7.38
Operative weight (kg)	4.25	3.42	7.44
Operative time (h)	3.50	2.00	5.00
Carbon dioxide insufflation time (h)	2.00	1.00	3.00
End-tidal carbon dioxide (mmHg)	37.60 (mean)	32.73	42.13
Body temperature (°C)	35.80	35.18	36.31
Peak inspiratory pressure level (cmH ₂ O)	23.30 (mean)	20.58	26.38
Follow-up (mo)	23.40	8.43	41.53

Download English Version:

<https://daneshyari.com/en/article/3475699>

Download Persian Version:

<https://daneshyari.com/article/3475699>

[Daneshyari.com](https://daneshyari.com)