

Case Report

Anesthetic management of comprehensive dental restoration in a child with glutaric aciduria type 1 using volatile sevoflurane

Wei-Nung Teng^{a,b}, Su-Man Lin^{a,b}, Dau-Ming Niu^{b,c}, Yi-Min Kuo^{a,b}, Kwok-Hon Chan^{a,b},
Chun-Sung Sung^{a,b,*}

^a Department of Anesthesiology, Taipei Veterans General Hospital, Taipei, Taiwan, ROC

^b National Yang-Ming University School of Medicine, Taipei, Taiwan, ROC

^c Department of Pediatrics, Taipei Veterans General Hospital, Taipei, Taiwan, ROC

Received November 29, 2012; accepted May 16, 2013

Abstract

Glutaric aciduria type 1 (GA1) is a rare, inherited mitochondrial disorder that results from deficiency of mitochondrial glutaryl-CoA dehydrogenase. Most patients develop neurological dysfunction early in life, which leads to severe disabilities. We present a 37-month-old girl with GA1 manifested as macrocephaly and hypotonia who received comprehensive dental restoration surgery under general anesthesia with sevoflurane. She was placed on specialized fluid management during a preoperative fasting period and anesthesia was administered without complications. All the physiological parameters, including glucose and lactate blood levels and arterial blood gas were carefully monitored and maintained within normal range perioperatively. Strategies for anesthetic management should include prevention of pulmonary aspiration, dehydration, hyperthermia and catabolic state, adequate analgesia to minimize surgical stress, and avoidance of prolonged neuromuscular blockade. We administered general anesthesia with sevoflurane uneventfully, which was well tolerated by our patient with GA1. Additionally, communication with a pediatric geneticist and surgeons should be undertaken to formulate a comprehensive anesthetic strategy in these patients. Copyright © 2014 Elsevier Taiwan LLC and the Chinese Medical Association. All rights reserved.

Keywords: dental restoration; fluid therapy; general anesthesia; glutaryl-CoA dehydrogenase; inhalation anesthetics

1. Introduction

Glutaric aciduria type 1 (GA1) is a rare autosomal recessive disorder with an estimated prevalence of 1:100,000 newborns in a Taiwanese study.¹ This disorder involves a deficiency of mitochondrial glutaryl-CoA dehydrogenase (GCDH), which results in an inborn error of lysine, hydroxyllysine, and tryptophan metabolism.² Excessive organic acids

such as glutaric acid, 3-hydroxyglutaric acid, and glutaconic acid accumulate in the brain and lead to neuronal damage, lymphocyte infiltration, elevated concentrations of inflammatory cytokines and nitric oxide, glial proliferation, atrophy of striatal neurons, and neurologic dysfunction.³ Though newborn screening and early treatment can prevent such neurological damage in most patients with GA1, routine treatment does not protect against the encephalopathic crisis that can be precipitated by fever, gastroenteritis (GE), and surgery.⁴ Therefore, intensified emergency treatment targeting prevention or reversal of encephalopathic crisis has been recommended to commence without delay and performed aggressively, especially during surgical intervention.⁵ However, there are only a few reports regarding general anesthesia conducted under either total intravenous anesthesia (TIVA)

Conflicts of interest: The authors declare that there are no conflicts of interest related to the subject matter or materials discussed in this article.

* Corresponding author. Dr. Chun-Sung Sung, Department of Anesthesiology, Taipei Veterans General Hospital, 201, Section 2, Shih-Pai Road, Taipei 112, Taiwan, ROC.

E-mail address: cssung@vghtpe.gov.tw (C.-S. Sung).

with propofol for ventriculoperitoneal shunting or inhalational anesthesia with sevoflurane for reduction of hip dislocation in patients with GA1.^{6,7} We report here the anesthetic management with sevoflurane for a 37-month-old girl with GA1 undergoing comprehensive dental restoration surgery.

1.1. Patient consent statement

Written informed consent was signed and obtained from the patient's parent. The authors assured the patient's parent that the information will be published without the patient's name attached and every attempt made to maintain anonymity.

2. Case report

A 37-month-old girl with GA1 who weighed 19 kg and who presented with macrocephaly and hypotonia was scheduled for comprehensive dental restoration surgery for treatment of multiple dental caries. Because glutaric aciduria was suspected during neonatal genetic screening, she was referred to our hospital for diagnosis and treatment on postnatal Day 10. The diagnosis was confirmed by the large amount of urinary glutaric acid along with elevated serum glutarylcarnitine. The patient was treated with oral carnitine supplementation and protein restriction with special formula milk (XLYS, LOW TRY Analog). Magnetic resonance imaging of the brain performed under sedation anesthesia with intravenous midazolam and thiopental at 3 months of age showed prominent extra-axial cerebrospinal fluid space, marked widening of Sylvian fissures and middle cranial fossa, and relative loss of bilateral frontotemporal volume. The patient had a history of repeated upper respiratory tract infection (URI) and GE at her 5-, 9-, and 13-month check-ups. No acute neuromotor deterioration was present, and psychomotor development assessed by the age of 7 months showed no psychological delay except mild motor delay. On preanesthetic clinical examination, the patient was asymptomatic except for macrocephaly, with an occipitofrontal circumference of 53 cm (>97th percentile for age), and hypotonia.

2.1. Anesthetic management

During the preoperative fasting period, the patient received intravenous fluid therapy with 10% dextrose in 0.33% saline with potassium chloride (KCl, at a concentration of 6 mEq/500 mL) infused at 60 mL/hour and 20% intralipid infused at 6 mL/hour. After routine monitors and preoxygenation were applied, anesthesia was induced with atropine (0.01 mg/kg), thiamylal (5 mg/kg), fentanyl (2 µg/kg), and cisatracurium (0.2 mg/kg) under parental accompaniment, and manual ventilation was applied using the Sellick maneuver. The patient underwent nasal intubation intubated with a 5-mm cuffed endotracheal tube, and anesthesia was maintained with sevoflurane (1–1.5 minimum alveolar concentration) in 50% oxygen and cisatracurium 0.03 mg/kg every 60 minutes. A 24-gauge arterial catheter was inserted

into the left radial artery. Mechanical ventilation was targeted at maintaining an end-tidal carbon dioxide level of 35–40 mmHg. The patient's body temperature was maintained between 36°C and 37°C with a warming blanket and intravenous fluid therapy was continued until oral intake resumed. All the physiological parameters, blood levels of glucose and lactate, and arterial blood gas were closely monitored and maintained within normal range perioperatively. Ketorolac (0.5 mg/kg) was intravenously administered 30 minutes before the end of the surgical procedure, which was 5 hours in duration. Neuromuscular blockade was reversed with neostigmine and the patient was extubated without difficulty in the operating room. The patient resumed food intake 8 hours later and was discharged uneventfully the next day.

3. Discussion

GA1 is a rare genetic disorder caused by GCDH deficiency, with organic acid accumulation in the brain and neurodegeneration. Secondary carnitine deficiency will occur as the accumulated organic acids are detoxified by carnitine. Patients with GA1 often present with macrocephaly at birth or dystonia shortly afterward. Brain imaging performed shortly after birth usually shows frontoparietal atrophy with widening of Sylvian fissures and arachnoid cysts, and the brain is vulnerable to head trauma that can lead to acute subdural or retinal hemorrhage.^{3,7} Brain atrophy and neuronal loss following injury to the basal ganglia may be accompanied by symptoms such as dystonia, spasticity, rigidity, posture impairment, and complications that impair oral feeding and communication.^{3,7} Patients usually present with an episode of acute infection accompanied by fever and dehydration, and may become hypotonic, lose head control, and have choreoathetosis similar to seizures. Hypotonia slowly improves over weeks but is then replaced with rigidity and dystonia; therefore, patients may develop severe disability and remain wheelchair-bound. These children develop decreased coordination of swallowing movements and are prone to pulmonary aspiration. Some patients may even require nasogastric tube feeding and later require permanent gastrotomy.⁴ Although these patients have difficulty taking or performing tasks, cognition is usually normal. Acute neurological deterioration or encephalopathic crisis, marked by spasm, dystonia, and lethargy, is caused by glutaric acid passage through the blood–brain barrier.^{7,8} Patients with GA1 will develop neurological damage with complications and reduced life expectancy following encephalopathic crisis, and both pneumonia and encephalopathic crisis are the most common causes of death in these patients.^{3,4,8}

With neonatal screening, early routine treatment with carnitine supplement special restriction diet, and neuroprotective emergency treatment, most patients with GA1 may have protection against encephalopathic crisis and possibly lead a normal life.^{7,9–11} Screening identifies elevated glutarylcarnitine in blood spots, and newborns with positive screening results are then confirmed by urinary

Download English Version:

<https://daneshyari.com/en/article/3475993>

Download Persian Version:

<https://daneshyari.com/article/3475993>

[Daneshyari.com](https://daneshyari.com)