



Successful treatment of refractory functional dyspepsia with osteopathic manipulative treatment

Nichole J. Mirocha, DO, MHS,^a Jonathan D. Parker, DO, MS, PGY-1^b

From the ^aAOA/ACGME accredited Family Medicine Residency Program, Department of Family & Community Medicine, Southern Illinois University School of Medicine, Springfield, IL; and

^bKirkville College of Osteopathic Medicine, A.T. Still University of Health Sciences, Kirkville, MO

KEYWORDS:

Dyspepsia;
GERD;
Viscero-somatic;
Osteopathic
manipulative treatment

Functional gastrointestinal disorders (FGIDs) describe patients who present with symptoms that arise from the gastrointestinal (GI) tract yet have failed to be explained by a standard GI work-up. Somatic findings at midthoracic levels correspond to localized visceral dysfunction such as decreased peristalsis and vasoconstriction. The somatovisceral concept suggests that osteopathic manipulative treatment (OMT) provides relief by normalizing the autonomic tone. There are few reports in the literature that provide examples of successful treatment of FGID by OMT. This case describes a patient with postprandial nausea and vomiting who was successfully treated with OMT.

© 2012 Elsevier Inc. All rights reserved.

Case presentation

A 37-year-old female patient presented to her family physician with a four-year history of intermittent postprandial nausea and vomiting, which the patient described as “regurgitation.” These symptoms were not related to a particular food, time of day, or the onset of any new medications. She reported abdominal pain only at the time of vomiting and she denied hematemesis, dysphagia, diarrhea, constipation, hematochezia, melena, or weight loss. Upon review of systems, she denied fever, flank pain, vaginal discharge, pelvic pain, dysuria, hematuria, urinary hesitancy, or incontinence. She had been seen four years previously by a gastroenterologist and had an esophagogastroduodenoscopy, upper gastrointestinal (GI) series, and trial of several medications including H-2 blockers, proton-pump inhibitors, and metaclopramide. At that time she was given a diagnosis of gastroesophageal reflux disease (GERD) and denied

seeking additional testing or treatment for her GI symptoms from other providers in the intervening years.

Other than GERD, her past medical history included hypertension, hypothyroidism, ovarian cysts, major depressive disorder, transient ischemic attack, and endometriosis for which hysterectomy was performed. She was a G2P2 with a history of two normal spontaneous vaginal deliveries.

Her medications were sertraline, levothyroxine, metoprolol, amlodipine, and pantoprazole. She denied taking any over-the-counter medications or supplements.

Her relevant family history revealed her father had died at age 74 from a cancer in the small intestine of which she did not know further details. Her maternal uncle had liver cancer of which she could also not provide further details. Her social history was negative for tobacco, alcohol, and recreational drug use.

On presentation, her blood pressure was 126/94 and pulse was 84 beats per minute. Physical examination showed a healthy-appearing female weighing 165 pounds. Her abdomen was soft, nontender, nondistended, and with normoactive bowel sounds. Her osteopathic examination revealed tissue congestion over the superior

Corresponding author: Jonathan D. Parker, DO, MS, PGY-1, Tuscaloosa Family Medicine Residency, University of Alabama, Tuscaloosa, AL 35401.

E-mail address: jdparker@cchs.ua.edu.

mesenteric ganglion, a flexed T5 rotated right and sidebent right, and a group curve at T7-9 flexed rotated right and sidebent left. The remainder of her physical examination was normal.

Diagnosis and topic discussion

Given this patient's prior diagnosis of GERD by her gastroenterologist after extensive diagnostic studies, and the chronic nature of symptoms over four years without worrisome progression or serious signs such as weight loss, and coupled with the current history and physical examination, our differential diagnosis included GERD, gastroparesis, bulimia nervosa, food allergy, somato-visceral and viscerosomatic reflex, and medication side effects. She was given a working diagnosis of postprandial nausea and vomiting with somatic dysfunction of the thoracic spine.

Given that the patient had been worked up in the past by gastroenterology, wide-ranging workup was not pursued. However, further investigation by her primary care provider for dyspepsia/postprandial nausea and vomiting at this presentation would have included further questioning about over the counter medications including: NSAIDs, Aspirin, and less common causes such as Garlic, Gingko, Saw Palmetto, Feverfew, Iron, and Chaste tree berry. Further labwork to be considered would include TSH/Free T4 if the patient would have exhibited symptoms of dyspepsia along with symptoms of thyroid disease and/or fasting blood glucose/HbA1C if the patient had not been screened or had exhibited symptoms of diabetes.

Additionally, one may consider imaging studies including barium esophagography with small bowel follow-through, ultrasound to investigate the liver, gallbladder, and pancreas, and direct visualization of her upper GI tract with esophagogastroduodenoscopy.

One special consideration for evaluation for dyspepsia is *H. Pylori* testing by stool antigen or ¹³C-urea testing. The utility of *H. Pylori* testing in patients with dyspepsia without known PUD is debatable and beyond the scope of this paper—though, the decision could be made to perform this test given clinical presentation.

Functional gastrointestinal disorders (FGIDs) are a group of abnormalities that afflict patients with symptoms that are assumed to arise from the gastrointestinal tract yet have failed to be explained by a standard work-up of endoscopy, laboratory studies, and radiological evaluation.¹ The Rome Foundation process has sought to classify FGIDs since 1990 and has evolved its classifications in three separate consensuses: ROME I, ROME II, and, most recently, ROME III. The Rome I consensus divided FGIDs into six divisions based on the anatomic region in which the symptoms are thought to originate.² Functional dyspepsia (FD) is the disorder thought to encompass the gastroduodenum. The ROME III consensus categorized FD into two major presenta-

tions: (a) Postprandial distress syndrome (PDS), which is characterized by early satiety, postprandial fullness, and nausea (and/or vomiting); and (b) epigastric pain syndrome (EPS), characterized by meal-independent epigastric pain and burning.² The difficulty with the overlap of GERD and EPS, with its frequent simultaneous presentation in patients with FD, has been recognized by ROME III; however, uniform guidelines for classification and characterization have not been established.³ Nonetheless, this probably causes more of an academic problem for research stratification and less of an impedance for patient care. This patient appeared to have PDS given her four-year history of postprandial nausea and vomiting without a specific organic, systemic, or metabolic explanation despite prior workup by a gastroenterologist.

The incidence and prevalence of FD is difficult to accurately describe because of the nature and overlap of the symptoms. However, it has been reported that as many as 20% to 30% of people have chronic or recurrent dyspepsia and the annual incidence of first-time sufferers is about 1%; more than 50% of these patients lack an organic case despite diagnostic testing.³

The parasympathetic nerve supply to the upper gastrointestinal tract is from the anterior and posterior vagal trunks derived from the vagus nerve (CN X). Increased parasympathetic activity results in increased hydrochloric acid production in the stomach and a higher rate of peristalsis.⁴ The sympathetic nerve supply originates from the T5-T9 segments of the spinal cord, forming the greater splanchnic nerve with synapsis in the celiac plexus. Increased sympathetic activity results in decreased peristalsis and vasoconstriction.⁴

The anatomy and physiology of the assumed origin of the symptoms in a patient with FD becomes of value when considering the somato-visceral connection. Since the early work of Louisa Burns, DO, the somato-visceral concept has been widely accepted and used in osteopathic medicine,⁵⁻⁹ suggesting that somatic dysfunction in T5-T9 could result in corresponding changes in acid production, gut motility, and vascular tone (and consequently oxygen/nutrient delivery). A review of the literature does not reveal any specific studies or case reports on the use of OMT in the treatment of patients with FGIDs. There are, however, case reports on the successful adjuvant use of OMT in the treatment of peptic ulcer disease (PUD)¹⁰ and GERD.¹¹ Both the report on PUD, by Morris and Dickey, and the report on GERD, by Branyon, suggest that the mechanism by which OMT provides relief is the effect the manipulation has on normalizing the autonomic tone, which is consistent with the historical understanding of the somato-visceral concept.¹² They also provide thorough discussions on the osteopathic literature supporting this mechanism.

In this case, the chief complaint was postprandial nausea and vomiting, which differs from the most common presentation of either GERD or PUD. Recent studies have indicated another possible mechanism of OMT on

Download English Version:

<https://daneshyari.com/en/article/3813695>

Download Persian Version:

<https://daneshyari.com/article/3813695>

[Daneshyari.com](https://daneshyari.com)