



## Original Article

# Recognition of the three deduced probable human leukocyte antigen haplotypes in association with HLA-A\*31:30 (A\*31:30-B\*15-DRB1\*14) and HLA-B\*40:55 (A\*02:07-B\*40:55-DRB1\*04:05 and A\*26:01-B\*40:55-DRB1\*09:01) in a Taiwanese population

Kuo-Liang Yang<sup>a,b,\*</sup>, Py-Yu Lin<sup>a</sup><sup>a</sup>Laboratory of Immunogenetics, Tzu Chi Cord Blood Bank and Buddhist Tzu Chi Marrow Donor Registry, Buddhist Tzu Chi Stem Cells Center, Buddhist Tzu Chi General Hospital, Hualien, Taiwan<sup>b</sup>Department of Laboratory Medicine, Tzu Chi University, Hualien, Taiwan

## ARTICLE INFO

## Article history:

Received 5 August 2013

Received in revised form

5 September 2013

Accepted 11 September 2013

## Keywords:

Haplotypes

HLA

Sequence-based typing

Stem cell

Transplantation

## ABSTRACT

**Objectives:** HLA-A\*31:30 and HLA-B\*40:55 are two rarely observed alleles in the HLA-A locus and HLA-B locus, respectively. The objective of this study is to report three deduced probable human leukocyte antigen (HLA) haplotypes in association with HLA-A\*31:30 and HLA-B\*40:55 in unrelated bone marrow hematopoietic stem cell donors.

**Materials and methods:** A sequence-based typing method was used to confirm the two low-incidence alleles observed. A polymerase chain reaction was performed to amplify exons 2, 3, and 4 of the HLA-A, -B, and -C loci and exon 2 of the HLA-DRB1 locus with group-specific primer sets. Amplicons were sequenced using the BigDye Terminator Cycle Sequencing Ready Reaction Kit in both directions according to the manufacturer's protocols.

**Results:** The DNA sequence of A\*31:30 is identical to A\*31:01:02 in exons 2, 3, and 4, except for a nucleotide substitution at residue 539 (T→G) resulting in an amino acid replacement at position 156 (Leu→Trp). We deduced the probable HLA haplotype in association with A\*31:30 as A\*31:30-B\*15-DRB1\*14. The DNA sequence of B\*40:55 is identical to B\*40:01:01 in exons 2, 3, and 4 except for a nucleotide exchange at residue 814 (G→A) resulting in an amino acid substitution at position 248 (Val→Met). Two probable HLA haplotypes associated with B\*40:55 may be deduced as A\*02:07-B\*40:55-DRB1\*04:05 and A\*26:01-B\*40:55-DRB1\*09:01.

**Conclusion:** Information about the deduced HLA haplotypes associated with the rare A\*31:30 and B\*40:55 alleles that we reported here is valuable for HLA tissue typing laboratories for reference purposes and for stem cell transplantation donor search coordinators to determine the likelihood of finding compatible donors in unrelated bone marrow donor registries for patients bearing these two uncommon HLA alleles. Because A\*31:30 and B\*40:55 have been found in Taiwanese population so far, we think the haplotypes that we reported here are most likely conserved in their population.

Copyright © 2013, Buddhist Compassion Relief Tzu Chi Foundation. Published by Elsevier Taiwan LLC. All rights reserved.

## 1. Introduction

The major histocompatibility complex (MHC) in humans consists of several loci of genes located on the short arm of chromosome 6 at 6p21.3. These loci are classified into Class I, II, and III of

the MHC and the genes of human leukocyte antigen (HLA) alleles are situated in the MHC Class I and II regions. The HLA genes are characterized by their extreme allelic polymorphism and their variations and diversity among different ethnic groups and racial populations. As the HLA molecule similarity between donors and recipients is being used as a prediction factor for graft survival and graft-versus-host disease, it is imperative to characterize precisely any new allele encountered during routine HLA typing procedures. To facilitate successful and comprehensive unrelated bone marrow donor searches for patients in need of hematopoietic stem cell transplantation, we are persistently working on resolving

Conflicts of interest: none.

\* Corresponding author. Buddhist Tzu Chi Stem Cells Center, Buddhist Tzu Chi General Hospital, 707, Section 3, Chung-Yang Road, Hualien, Taiwan. Tel.: +886 38561825x3373; fax: +886 38567851.

E-mail address: edward@tzuchi.com.tw (K.-L. Yang).

cDNA	80	90	100	110	120	130	140	150	160	170
A*31:01:02	GCTCCA	CTCCATGAGG	TATTTTCA	CATCCGTGTC	CCGGCCCCGGC	CGCGGGGAGC	CCCGCTTCAT	CGCGTGGGC	TACGTGGACG	ACACGCAGTT
A*31:30	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
cDNA	180	190	200	210	220	230	240	250	260	270
A*31:01:02	CGTGC	GACAGCGAGC	CCGCGAGCCA	GAGGATGGAG	CCGCGGGGCGC	CGTGGATAGA	GCAGGAGAGG	CCTGAGTATT	GGGACCAGGA	GACACCGGAAT
A*31:30	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
cDNA	280	290	300	310	320	330	340	350	360	370
A*31:01:02	GTGAAGGCC	ACTCACAGAT	TGACCGAGTG	GACCTGGGGA	CCCTGCGCGG	CTACTACAAC	CAGAGCGAGG	CCG GTTCTCA	CACCATCCAG	ATGATGTATG
A*31:30	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
cDNA	380	390	400	410	420	430	440	450	460	470
A*31:01:02	GCTGCGACGT	GGGTCGGAC	GGGCGCTTCC	TCCGCGGGTA	CCAGCAGGAC	GCCTACGAGC	GCAAGGATTA	CATCGCCTTG	AACGAGGACC	TGCGCTCTTG
A*31:30	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
cDNA	480	490	500	510	520	530	540	550	560	570
A*31:01:02	GACCGCGGCG	GACATGGCGG	CTCAGATCAC	CCAGCGCAAG	TGGGAGGCGG	CCCGTGTGGC	GGAGCAGTTG	AGAGCCTACC	TGGAGGGCAC	GTGCGTGGAG
A*31:30	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
cDNA	580	590	600	610						
A*31:01:02	TGGCTCCGCA	GATACCTGGA	GAACGGGAAG	GAGACGCTGC	AGCGCACGG					
A*31:30	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

**Fig. 1.** Comparison of DNA sequences between HLA-A\*31:30 and -A\*31:01:02 in exons 2 and 3. The DNA sequence of A\*31:30 is identical to A\*31:01:02 in exons 2 and 3, except at residue 539 where the T of A\*31:01:02 is replaced by G (shaded). cDNA = complementary DNA.

ambiguous or unidentified alleles that we find to offer better services for HLA matching and donor selection.

HLA-A\*31:30 and HLA-B\*40:55 were first reported to the immunogenetics (IMGT)/HLA database in 2010 and 2004, respectively [1], without an indication of probable HLA-associated haplotypes. Here we report the deduced probable HLA haplotypes in association with A\*31:30 and B\*40:55. We further postulate that the deduced plausible HLA haplotype in association with A\*31:30 is most likely conserved in the Taiwanese population, based on its low frequency in the general population and the fact that it has so far been reported only in a Taiwanese population [1]. Similarly, the deduced probable B\*40:55-associated HLA haplotype A\*02:07-B\*40:55-DRB1\*04:05 is most likely conserved in the Taiwanese population, whereas the other deduced probable B\*40:55-associated haplotype, A\*26:01-B\*40:55-DRB1\*09:01, is conserved in Japanese and Taiwanese populations.

## 2. Materials and methods

Peripheral whole blood samples from three unrelated bone marrow stem cell donors with Taiwanese ethnicity were collected in acid citrate dextrose (ACD) anticoagulant. Formal written consents were signed by the donors before blood collection. The ACD whole blood was stored at  $-80^{\circ}\text{C}$  until use. Genomic DNA was extracted using the QIAamp DNA Blood Mini Kit according to the manufacturer's instructions (Qiagen, Hilden, Germany). Genomic DNA typing of HLA-A, -B, and -DRB1 loci was first performed using Dynal Reli-sequence-specific oligonucleotide probe (SSO) HLA-A, -B, and -DRB1 Typing Kits (Dynal Biotech, Bromborough, Wirral, UK), followed by the sequence-specific primer (SSP) typing method (AllSet Gold SSP HLA high-resolution kits; Dynal Biotech, Invitrogen, Brown Deer, WI, USA) to reach high-resolution allelic subtypes. The sequence-based typing method [2–6] was used to confirm the low-incidence alleles observed, and in cases of anomalous results and typing ambiguities from the SSO and SSP typing

protocols. Polymerase chain reaction was carried out to amplify exons 2, 3, and 4 of the HLA-C locus and exon 2 of the DQB1 locus with group-specific primer sets as previously described [7]. Amplicons were sequenced by the BigDye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems, Foster City, CA, USA) in both directions. In this study, A\*31:30 from one blood donor and B\*40:55 from another two blood donors were sequenced and analyzed.

## 3. Results

We confirmed that the DNA sequence of A\*31:30 is identical to A\*31:01:02 in exons 2, 3, and 4, except for a nucleotide substitution at residue 539 (T  $\rightarrow$  G) [1] (Fig. 1). The nucleotide substitution caused an amino acid replacement at residue 156 (Leu  $\rightarrow$  Trp; Fig. 2). The extended HLA typing of our donor carrying A\*31:30 was A\*02:07, A\*31:30, B\*15:01, B\*46:01, DRB1\*09:01, and DRB1\*14:05. Together with the HLA typing of the cell (328573) with A\*31:30 reported to the IMGT/HLA database by Yang et al (A\*11:01, A\*31:30, B\*15, DRB1\*14, and DRB1\*16) [1], the probable HLA haplotype in association with A\*31:30 may be deduced as A\*31:30-B\*15-DRB1\*14.

Historically, we detected three unrelated bone marrow stem cell donors bearing the B\*40:55 allele in our bone marrow donor registry. We confirmed that the DNA sequence of B\*40:55 is identical to B\*40:01:01 in exons 2, 3, and 4 except for a nucleotide substitution at position 814 (G  $\rightarrow$  A; Fig. 3). The nucleotide replacement resulted in an amino acid substitution at position 248 (Val  $\rightarrow$  Met; Fig. 4). The extended HLA typing of the three bone marrow donors was as follows: A\*02:07, A\*33:03, B\*40:55, B\*46:01, DRB1\*04:05, DRB1\*09:01; A\*02:07, B\*40:55, B\*51:01, DRB1\*04:05, DRB1\*11:01; and A\*24:02, A\*26:01, B\*40:55, B\*46:01, DRB1\*04:05, DRB1\*09:01. Together with the extended HLA typing of the Japanese donor (TBC 46239) bearing B\*40:55 reported to the IMGT/HLA database (A\*24:02, A\*26:01, B\*40:01, B\*40:55, DRB1\*09:01, DRB1\*15:01) [1], the following two plausible HLA haplotypes associated with

AA Pos.	10	20	30	40	50	60	70	80	90	100
A*31:01:02	GSHSMRYFTT	SVSRPGRGEP	RFlAVGYVDD	TQFVRFDSDA	ASQRMEPRAP	WIEQERPEYD	DQETRNKVAH	SQIDRVDLGT	LRGYYNQSEA	GSHTIQMMYG
A*31:30	*-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
AA Pos.	110	120	130	140	150	160	170	180		
A*31:01:02	CDVGS	DGRFL	RGYQDAYDG	KDYIALNEDL	RSWTAADMAA	QITQRKWEAA	RVAEQLRAYL	EGTCVEWLR	YLENGKETLQ	RT
A*31:30	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

**Fig. 2.** Comparison of amino acid sequences between HLA-A\*31:30 and -A\*31:01:02 in exons 2 and 3. The amino acid sequence of A\*31:30 is identical to A\*31:01:02 in exons 2 and 3, except at residue 156 where the L of A\*31:01:02 is replaced by W (shaded). HLA = human leukocyte antigen.

Download English Version:

<https://daneshyari.com/en/article/3841930>

Download Persian Version:

<https://daneshyari.com/article/3841930>

[Daneshyari.com](https://daneshyari.com)