

Slightly lower incidence of ectopic pregnancies in frozen embryo transfer cycles versus fresh in vitro fertilization-embryo transfer cycles: a retrospective cohort study

Wim Decler, M.D.,^a Kaan Osmanagaoglu, Ph.D.,^a Geertrui Meganck, M.D.,^b and Paul Devroey, Ph.D.^c

^a IVF Centrum, AZ Jan Palfijn Hospital, Gent; ^b Onze-Lieve-Vrouwziekenhuis, Aalst; and ^c Center for Reproductive Medicine, Dutch-speaking Free University Brussels, Brussels and AZ Jan Palfijn Hospital, Gent, Belgium

Objective: To analyze the incidence of ectopic pregnancies (EPs) in fresh and frozen/thawed cycles.

Design: A retrospective cohort study on the incidence of EPs in all fresh IVF cycles from January 2002 until December 2012. This was compared with the incidence of tubal pregnancies in patients undergoing transfer of frozen/thawed embryos during the same time period.

Setting: The IVF program at Fertility Center, AZ Jan Palfijn, Gent, Belgium.

Patient(s): A total of 11,831 patients undergoing IVF (i.e., the entire population of the IVF Center) were retrospectively analyzed.

Intervention(s): The IVF cycles, fresh IVF-ET, frozen/thawed ET. Laparoscopy for treatment of EP.

Main Outcome Measure(s): Primary end point: incidence of EPs in both groups. Secondary end points: clinical pregnancy rate (PR), rate of EPs per clinical pregnancy.

Result(s): In the fresh IVF cycle group, 10,046 patients underwent oocyte retrieval; 9,174 of them had an ET; 2,243 of these patients had a clinical pregnancy. Of these, 43 (0.47%) appeared to have an ectopic localization of their pregnancy. In the group of the patients undergoing frozen/thawed ET (1,785 patients) there were 467 pregnancies and 6 ectopic implants (0.34%). The incidence of the EPs per established clinical pregnancy was 1.92% for the fresh vs. 1.28% for the frozen/thawed cycles.

Conclusion(s): No significant difference could be demonstrated on the incidence of EP in fresh IVF cycles vs. frozen/thawed cycles in a large cohort of patients. (Fertil Steril® 2014;101:162–5. ©2014 by American Society for Reproductive Medicine.)

Key Words: Ectopic pregnancy, IVF, frozen/thawed cycle, relative risk

Discuss: You can discuss this article with its authors and with other ASRM members at <http://fertilityforum.com/declerwe-ectopic-pregnancy-frozen-cycle-embryo-transfer-ivf/>



Use your smartphone to scan this QR code and connect to the discussion forum for this article now.*

* Download a free QR code scanner by searching for "QR scanner" in your smartphone's app store or app marketplace.

Infertility patients have always been associated with a higher risk of tubal pregnancy (1). Tubal surgery, including microsurgical operations and laparoscopic interventions such as salpingectomy or tubal reanastomosis, led to an astonishing number (15%–20%) of ectopic pregnancies (EPs) (2).

Even after IVF where the tubal passage was bypassed and the embryos were transferred directly in the uterine cavity, still a significant number of ectopic localizations of the gravidity were found (3). The reasons are numerous, such as tubal disease (4), increased uterine contractions due to ovarian

stimulation (5), dysfunction of the uterine musculature due to high P levels (6), and side effects of the medication (7).

Recently it has been reported (8) that the EP rate is significantly reduced after the replacement of frozen/thawed embryos. To corroborate a retrospective analysis of the data of the IVF Center of the Hospital Jan Palfijn in Gent (Belgium) was performed from January 1, 2002 to December 31, 2012.

Received April 24, 2013; revised September 16, 2013; accepted October 3, 2013; published online November 12, 2013.

W.D. has nothing to disclose. K.O. has nothing to disclose. G.M. has nothing to disclose. P.D. has nothing to disclose.

Reprint requests: Wim Decler, M.D., AZ Jan Palfijn, Henri Dunantlaan 5, 9000 Gent, Belgium (E-mail: dokter@fertility-belgium.be).

Fertility and Sterility® Vol. 101, No. 1, January 2014 0015-0282/\$36.00

Copyright ©2014 American Society for Reproductive Medicine, Published by Elsevier Inc. <http://dx.doi.org/10.1016/j.fertnstert.2013.10.002>

MATERIALS AND METHODS

All patients who underwent a fresh IVF treatment and had an ET between

January 1, 2002 and December 31, 2012, were included (group A). They were compared with all patients who had a transfer with a frozen/thawed cycle during the same period of time (11 years) (group B). The treatment procedures and the comparison of data were approved by the local ethical committee as being part of the standard quality control measurements.

Group A patients were stimulated with FSH (recombinant FSH; Puregon, MSD and Gonal F, Serono), sometimes in combination with urinary extracted FSH (hMG; Menopur, Ferring). The stimulation was performed with pituitary suppression and the administration of GnRH antagonists (Cetrotide 0.25 mg, Serono or Orgalutran 0.25 mg/0.5 mL, MSD) from day 6 onward until triggering. Final oocyte maturation was achieved by administration of 5,000–10,000 IU of hCG (Pregnyl, MSD) as soon as three follicles reached 18 mm in diameter at the time of ultrasound evaluation of the stimulation. Historically, 10,000 IU hCG was administered until 2008 to trigger ovulation. From 2009 onward the dose of hCG was reduced to 5,000 IU only to minimize the risk for ovarian hyperstimulation syndrome (OHSS). In less than 10% of patients—selected indications such as endometriosis or poor embryo quality in a previous stimulation cycle—down regulation of the cycle with GnRH agonists before ovarian stimulation (long protocol stimulation) was chosen for controlled stimulation. In these patients the triggering was performed by the administration of hCG. The oocyte retrieval took place by an ultrasound-guided transvaginal approach 36 hours after triggering. Fertilization of the oocytes took place either by IVF or intracytoplasmic sperm injection (ICSI), according to the sperm quality. An indication for ICSI was found if the total number of fast progressive motility sperm was $\leq 1.10^6$ and/or normal morphology was $\leq 3\%$ (Krüger criteria) or if a poor fertilization was seen in previous cycles. The embryos were cultured in either G1 Plus (Vitrolife) or ISM1 (Origio). Embryo transfer was generally performed on day 3 after fertilization. Only in case of repeated failure of implantation and if sufficient embryos of good quality were available, blastocyst culture was chosen with ET on day 5 after pick-up. If only one or two embryos were available for transfer, it took place on the second day after fertilization ($<5\%$). The number of transferred embryos varied from one to three, according to the Belgian legislation (Table 1). Supernumerary embryos of good quality (9) A or B type, less than 10% fragmentation, seven to eight-cell stage

on day 3, or blastocyst stage on day 5 were frozen (slow freezing with cryogenesis).

In group B patients the frozen/thawed embryos were transferred after optimization of the cycle with clomiphene citrate (CC) (100 mg/d; Clomid, Sanofi-Aventis) from day 5 until day 9 of the cycle, and induction of ovulation with 5,000 IU of hCG as soon as a follicle of 20 mm was visualized by ultrasound examination. For the patients without ovulation, E_2 valerate (Progynova, Bayer) was given vaginally in increasing doses (from 2 mg/d on day 1 to 6 mg/d on day 9). Progesterone (600 mg) (Utrogestan, Piette-Besins) was administered to mimic the luteal phase as soon as a 9-mm thick endometrium was visualized. Embryo transfer took place 5–7 days after hCG administration. In the anovulatory group the embryo replacement was performed 4–6 days after the initiation of micronized P (Utrogestan).

Human chorionic gonadotropin and P levels were measured 13 days (day 3 of transfer) after transfer, except for the blastocyst transfers where a pregnancy diagnosis already took place 11 days after transfer. For the patients who had a transfer on day 2 (4-cell stage), hCG level was measured on day 14 after transfer. In all patients with positive hCG measurement, the administration of micronized P vaginally was continued and a confirmation of the clinical pregnancy (echographic visualization of gestational sac) was realized by vaginal ultrasound examination approximately 10 days later. Subsequent follow-up of the pregnancy development, the evolution of the stimulated ovaries, and the presence of fetal heart activity 5 weeks after transfer were established.

The diagnosis of EP was made either by direct extrauterine visualization of the gestational sac or by finding an empty uterine cavity with increasing hCG levels more than 500 ng/mL. In all patients suspect for an ectopic localization of the pregnancy, the final confirmation was made by direct laparoscopic visualization. Laparoscopy was performed in all patients with hCG levels $>1,000$ ng/mL and with an empty cavity on ultrasound examination. Laparoscopic longitudinal incision of the tube, or salpingectomy in badly damaged tubes, was performed in all of those patients. This intervention was, without exception, performed laparoscopically. Especially in the group of tube-conserving surgery an adequate follow-up of the decreasing hCG levels postoperatively was established.

The percentages EP per ET obtained in patients undergoing fresh autologous ETs and those undergoing transfer with frozen/thawed embryos were compared. The same was done for the percentage of EPs per clinical pregnancy. The Fisher's exact test was performed one-sided at the 5% significance level. In doing so it is tested whether it is more likely to have an EP when receiving fresh ET (compared with frozen/thawed ET).

TABLE 1

Belgian legislation limits for number of transferred embryos.

Age cycleNo.	< 36 y-1 d	36 y to 40 y-1 d	40 y to 43 y-1 d	43 y to 45-1 d
1	1	2	No max	No max
2	1 or 2	2	No max	No max
3	2	3	No max	No max
4	2	3	No max	No max
5	2	3	No max	No max
6	2	3	No max	No max
Cryopreserved cycle	2	2	2	2

Decler. Relative risk of EUG in frozen IVF cycle. Fertil Steril 2014.

RESULTS

From January 1, 2002 to December 31, 2012, 9,174 fresh autologous ETs were performed. They were the result of 10,046 oocyte retrievals from the same period. This number resulted in 2,243 pregnancies. The overall clinical pregnancy

Download English Version:

<https://daneshyari.com/en/article/3938419>

Download Persian Version:

<https://daneshyari.com/article/3938419>

[Daneshyari.com](https://daneshyari.com)