#### Gynecology and Minimally Invasive Therapy 5 (2016) 112-115

Contents lists available at ScienceDirect

# **Gynecology and Minimally Invasive Therapy**

journal homepage: www.e-gmit.com

Original article

# Efficacy of short-term training for acquisition of basic laparoscopic skills

# Jun Takeda <sup>a</sup>, Iwaho Kikuchi <sup>b, \*</sup>, Ayako Kono <sup>a</sup>, Rie Ozaki <sup>a</sup>, Jun Kumakiri <sup>a</sup>, Satoru Takeda <sup>a</sup>

<sup>a</sup> Department of Obstetrics and Gynecology, Juntendo University, Faculty of Medicine, Tokyo, Japan <sup>b</sup> Department of Obstetrics and Gynecology, Juntendo University Urayasu Hospital, Chiba, Japan

#### ARTICLE INFO

Article history: Objective: To determine whether our newly developed short-term training program contributes to the Received 31 December 2014 acquisition of basic laparoscopic surgery skills. Received in revised form Design: Prospective study (Canadian Task Force Class II). 30 January 2015 Setting: University Hospital. Accepted 5 June 2015 Participants: Four obstetrics-gynecology residents who participated in our 2-month laparoscopic Available online 26 June 2015 training course. Interventions: Virtual reality laparoscopic surgery simulator-based assessment of four basic tasks: (1) Keywords: "camera manipulation"; (2) "eye-hand coordination"; (3) "two-handed maneuvers"; and (4) "cutting," laparoscopic surgery simulator before and after the course. laparoscopy training Measurements and main results: Mean times required to perform the tasks before and after training were resident teaching surgical education compared. The mean times required to perform three of the four tasks (except camera manipulation) virtual reality were significantly reduced after training. Total instrument path lengths were reduced, especially for instruments used by the dominant hand. Conclusion: Use of the virtual reality laparoscopic surgery simulator allowed us to objectively assess residents' acquisition of basic laparoscopic skills. We found that residents more readily acquired dominant-hand skills during their 2-month training. We conclude that our training system serves as an effective initial step towards the acquisition of the necessary laparoscopic surgery skills, even though residents do not actually perform surgeries during the training period. Copyright © 2015, The Asia-Pacific Association for Gynecologic Endoscopy and Minimally Invasive Therapy. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Introduction

Over the past 10 years, laparoscopic surgery has advanced rapidly, and indications have expanded. Because of the minimal invasiveness, reduced postoperative pain, shorter hospital stay, and improved wound cosmetics, patients tend to choose laparoscopic surgery over open surgery, and surgeons are thus required to learn the necessary skills. Laparoscopic surgery requires that the operator and assistants be adept at using long instruments under the limitation of a two-dimensional image. Acquisition of these

\* Corresponding author. Department of Obstetrics and Gynecology, Juntendo University Urayasu Hospital, Tomioka 2-1-1, Urayasu City, Chiba 279-0021, Japan. E-mail address: kikuchiban@hotmail.com (I. Kikuchi).

laparoscopic skills requires particular training, and the learning curve differs from that of open surgery.<sup>1</sup> Thus, we have developed our own laparoscopic training program to provide initial assistance in the development of the necessary manual dexterity. The program consists of two different courses-basic and advanced. The basic course is a 2-month course designed for obstetricsgynecology residents and aimed mainly at the management of inpatients. The advanced course is intended for more experienced surgeons, and successful completion of this course is acknowledged by the provision of a certificate of expertise in laparoscopic surgery. Approximately 100 operators and 500 assistants enroll in the advanced course annually.

During their formal training, residents may not acquire adequate skills. To address this issue, we used a virtual reality (VR) simulator to objectively assess the efficacy of our training program, which is directed towards enhancing residents' basic laparoscopic skills.









Conflicts of interest: The authors have no conflicts of interest relevant to this article.

http://dx.doi.org/10.1016/j.gmit.2015.06.001

<sup>2213-3070/</sup>Copyright © 2015, The Asia-Pacific Association for Gynecologic Endoscopy and Minimally Invasive Therapy. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

## Materials and methods

# VR simulator

For the objective assessment of residents' basic laparoscopic skills, a LAP Mentor VR simulator (Simbionix USA Corp., Cleveland, OH, USA) was used, and residents performed four basic tasks: (1) "camera manipulation"; (2) eye-hand coordination"; (3) "two-handed maneuvers"; and (4) "cutting" (Figure 1).

## Camera manipulation

For the camera manipulation task, residents were presented with a VR observation experience based on a 30° oblique camera view. The task required residents to search for a red dot displayed on the monitor, center the dot in the camera view, and then release the shutter (Figure 1A).

#### Eye-hand coordination

For the eye-hand coordination task, residents held an instrument with a red tip in one hand and an instrument with a blue tip in the other hand. The monitor displayed a bar with either red or blue dots at the top, and one of the dots flashed on and off. This task required residents to touch the flashing dot using the instrument tip of the corresponding color (Figure 1B).

#### Two-handed maneuvers

For the two-handed maneuvers task, residents held two pairs of grasping forceps, one in each hand. The monitor displayed blue jelly in which red dots appeared to be embedded. When the jelly was lifted with an instrument, the embedded red dots turned green and could be removed from the jelly with the other instrument. This task required residents to remove the green dots and place them in the designated basket (Figure 1C).

#### Cutting

J. Takeda et al. / Gynecology and Minimally Invasive Therapy 5 (2016) 112-115

For the cutting task, residents held grasping forceps and scissors forceps. The monitor displayed jelly, the bottom surface of which was connected to the "floor" with elastic threads. This task required residents to lift the jelly with the grasping forceps and then cut the connecting threads with the scissors forceps (Figure 1D).

## Participants

Four right-handed obstetrics-gynecology residents participated in the basic training course between August 2010 and December 2011. They each had 2 years of experience in general medicine but limited experience in laparoscopic surgery. The residents' performance of each task was assessed with the use of the simulator before and after 2 months of clinical training. During the clinical training period, the residents did not practice on the simulator. Also during training, residents acted as the first or second assistant, not as the operator. The main duties of the first assistant were to pull tissues with forceps and to cut surgical sutures with scissors forceps. The second assistant was responsible for holding the camera. The residents participated in approximately 30 live procedures as the first assistant and in 40 live procedures as the second assistant during the training period.

The operators were four surgeons with > 10 years of experience in laparoscopic surgery and each of the four operators had experience in over 1000 cases.

## Assessments and statistical analysis

The time required to complete each task and the instrument path lengths were assessed, both before and after the clinical training. The completion times are expressed as mean  $\pm$  standard deviation. The completion times before and after training were compared, and differences were analyzed with a paired *t* test, with



Figure 1. Screen appearance of the basic task modules. (A) Camera manipulation; (B) eye-hand coordination; (C) two-handed maneuvers; and (D) cutting.

Download English Version:

# https://daneshyari.com/en/article/3948261

Download Persian Version:

https://daneshyari.com/article/3948261

Daneshyari.com