



www.figo.org

Contents lists available at ScienceDirect

## International Journal of Gynecology and Obstetrics

journal homepage: www.elsevier.com/locate/ijgo



## EDUCATION AND TRAINING

## Laparoscopic virtual reality simulator and box trainer in gynecology

Ali Akdemir\*, Fatih Şendağ, Mehmet K. Öztekin

Department of Obstetrics and Gynecology, Ege University School of Medicine, Izmir, Turkey



## ARTICLE INFO

## Article history:

Received 14 July 2013

Received in revised form 23 October 2013

Accepted 17 January 2014

## Keywords:

Box trainer

Education

Laparoscopy

LapSim

Surgical training

Virtual reality

## ABSTRACT

**Objective:** To investigate whether a virtual reality simulator (LapSim) and traditional box trainer are effective tools for the acquisition of basic laparoscopic skills, and whether the LapSim is superior to the box trainer in surgical education. **Methods:** In a study at Ege University School of Medicine, Izmir, Turkey, between September 2008 and March 2013, 40 first- and second-year residents were randomized to train via the LapSim or box trainer for 4 weeks, and 20 senior residents were allocated to a control group. All 3 groups performed laparoscopic bilateral tubal ligation. Video records of each operation were assessed via the general rating scale of the Objective Structured Assessment of Laparoscopic Salpingectomy and by operation time in seconds. **Results:** Compared with the control group, the LapSim and box trainer groups performed significantly better in total score ( $P < 0.01$  and  $P < 0.01$ , respectively) and time ( $P = 0.03$  and  $P = 0.01$ , respectively). There were no differences between the LapSim and box trainer groups. **Conclusion:** Novice residents who trained on a LapSim or box trainer performed better live laparoscopies than residents who trained via standard clinical surgical education. Training with a virtual reality simulator or box trainer should be considered before actual laparoscopic procedures are carried out. © 2014 International Federation of Gynecology and Obstetrics. Published by Elsevier Ireland Ltd. All rights reserved.

## 1. Introduction

Laparoscopic surgery has had a fundamental role in gynecology over the past 2 decades [1]. On the one hand, it has not only reduced mortality and yielded better cosmetic results, but also given patients an opportunity for a shorter recovery time [2,3]. On the other hand, laparoscopic surgery presents some well-known technical complexities, requiring the ability to switch from 3D to 2D views, the handling of long rigid instruments that amplify any tremors or movement, adjustments for impaired tactile feedback, and familiarity with the fulcrum effect [1,4,5].

It is well known that decision making is superior to dexterity in terms of open surgery success; however, the latter is more important in laparoscopy [6]. These technical difficulties can be overcome by gaining novel and unique psychomotor skills; however, this is time consuming and requires a convenient teaching curriculum [7,8]. Moreover, patient safety and quality control mechanisms have come into prominence; when these events merge with increasing financial constraints, the efficacy of surgeons and cost-effectiveness of the procedures become more important in the operating room [9,10]. For these reasons, training in a pressure-free environment with virtual reality simulators and box trainers has become popular in the field of laparoscopic skill improvement.

The box trainer is a relatively inexpensive and multifunctional device that includes real laparoscopic instruments to give the student the option to train on animal parts and synthetic materials [11,12]. Newer virtual reality systems such as LapSim (Surgical Science,

Gothenburg, Sweden), which provides a considerable facility for both training and assessment, facilitate the replication of tasks such as cutting, grasping, and suturing [13–16]. However, these systems are relatively expensive and also require regular maintenance costs.

The primary aim of the present study was to investigate whether the LapSim virtual reality simulator and traditional box trainer are effective tools in the acquisition of laparoscopic psychomotor skills by comparing the LapSim and the box trainer with classic surgical education. A secondary aim was to test whether the LapSim virtual reality simulator is superior to the traditional box trainer in surgical education.

## 2. Materials and methods

The present prospective, randomized, blind, controlled trial was carried out between September 1, 2008, and March 31, 2013, among 60 gynecologic specialty residents at Ege University School of Medicine, Izmir, Turkey, who had no experience with the LapSim or box trainer. Approval for the study was obtained from the ethics committee of the university, and informed consent was obtained from residents and patients.

The obstetrics and gynecology specialization program takes 5 years in Turkey. Twenty of the participants were senior residents (postgraduate year 5); 40 were first- and second-year residents (postgraduate years 1–2). The senior residents comprised the control group and had previous experience with simple laparoscopic operations performed by a single hand, such as diagnostic laparoscopy and/or assisting senior colleagues (clinical standard education). The remaining 40 residents were randomly assigned to the LapSim group or the box trainer group. The training was spread out over a period of 5 weeks. During the first week, all groups received teaching in basic laparoscopy, laparoscopic bilateral tubal ligation (BTL), and the purpose of the study. During the

\* Corresponding author at: Department of Obstetrics and Gynecology, Ege University School of Medicine, Bornova TR-35100, Izmir, Turkey. Tel.: +90 505 674 89 04; fax: +90 232 343 07 11.

E-mail addresses: md.aliakdemir@gmail.com, ali.akdemir@ege.edu.tr (A. Akdemir).

subsequent 4 weeks (1 hour per week), the LapSim group and the box trainer group completed separate training sessions. The control group did not receive any further training.

After the training program, all residents performed their first laparoscopic BTL and were supervised by an experienced laparoscopic surgeon. All procedures were performed from the left side in order to standardize the comparison of the residents' performances. Patients with a severe systemic disorder, previous open abdominal surgery, a body mass index (BMI, calculated as weight in kilograms divided by the square of height in meters) of less than 18 or more than 30, or other factors that might have potentially negative effects on the surgical procedure were not included in the study.

The pneumoperitoneum was created via Veress needle through the umbilicus, and a 10-mm umbilical trocar for the optic system and 5-mm bilateral lower abdominal trocars for instrumentation were inserted. The ampullary–isthmic junction was identified with grasping forceps through the ipsilateral lower abdominal trocar, and the ampullary–isthmic junction was coagulated and cut with bipolar forceps and scissors through the contralateral lower abdominal trocar. Residents held both lower quadrant instruments and the supervising surgeon held the camera. Supervisors were not allowed to hold the lower quadrant instruments, and a researcher was responsible for observing and recording the procedure.

To evaluate technical performance and operation time, the raw video records were assessed by 2 independent observers, who were blind to the residents' identities. For technical performance, the "general skills items" of the Objective Structured Assessment of a Laparoscopic Salpingectomy (OSA-LS)," comprising a 5-item, general rating scale (economy of movements, confidence of movement: instrument handling, economy of time, respect for tissue, flow of operation/operative technique) and a 5-item, task-specific rating scale [17], was used (Table 1). The OSA-LS has been validated in a previous study [18]. The total score for each operation was determined by averaging the scores given by both independent observers. The operation time began when the resident held both instruments and ended when both lower abdominal trocars were taken out.

The LapSim virtual reality simulator is a PC-based system that includes a 19-inch monitor and a laparoscopic interface module with 2 instruments and a foot switch. The software is run on a dual-processor Pentium D 3-GHz computer with 1 GB of RAM and a GeForce 6800 graphics card using Windows XP Professional (Microsoft, Redmond, WA, USA) (Fig. 1).

LapSim provided training in 8 tasks (Fig. 2). Task 1, camera navigation: center randomly appearing balls in the camera view. Task 2, instrument navigation: indicate emerging balls by both instruments sequentially. Task 3, coordination: touch randomly appearing virtual spheres (1 hand holds the camera, and the other hand manipulates the instrument). Task 4, grasping: grasp, stretch, and remove virtual blood vessels. Task 5, cutting: 1 instrument grasps a virtual vessel to identify the target zone, while the other instrument cuts the vessel through the target zone with an ultrasonic dissector. Task 6, lifting and grasping: 1 instrument lifts a virtual object to visualize a hidden needle, while the other instrument grasps and carries the needle to a target zone. Task 7, suturing: suture the tissue-like target zone and tie the knot. Task 8, dissection: 1 instrument grasps and stretches an object to allow the other instrument to dissect a small vessel-like tissue by using monopolar cautery.

The box model trainer was constructed from dark plastic in the shape of a rectangular prism (45 × 30 × 25 cm). Five holes were cut out for the camera and trocars. The right side of the box trainer was left open for placement of the training tools (Fig. 3). The box trainer provided training in 7 tasks (Fig. 4). Task 1, excise a drawn circle with a 4-cm diameter from a thin sponge media directly on a line. Task 2, move pegs on a board. Task 3, cut the outer balloon of 2 balloons without rupturing the inner balloon, which is filled with ultrasound gel for mimicking ovarian cyst enucleation. Task 4, grasp and throw beans

**Table 1**  
Assessment chart for general skills of OSA-LS.

Item	OSA-LS score	1	2	3	4	5
Economy of movements						
Confidence of movement/instrument handling		Many unnecessary moves Repeatedly makes tentative or awkward moves with instruments	Efficient motion, but some unnecessary moves Competent use of instruments, although occasionally appears stiff or awkward	Efficient motion, but some unnecessary moves Competent use of instruments, although occasionally appears stiff or awkward	Efficient motion, but some unnecessary moves Competent use of instruments, although occasionally appears stiff or awkward	Maximum economy of movement Fluid moves with instruments and no awkward
Economy of time		Too much time taken to perform sufficiently	Intermediate time taken to perform sufficiently	Intermediate time taken to perform sufficiently	Intermediate time taken to perform sufficiently	Minimal time used to perform sufficiently
Respect for tissue		Frequently uses unnecessary force on tissue, risk of damage by inappropriate use of instruments, or instruments often out of sight	Careful handling of tissue but occasionally risk of (minimal) damage, or instruments out of sight	Careful handling of tissue but occasionally risk of (minimal) damage, or instruments out of sight	Careful handling of tissue but occasionally risk of (minimal) damage, or instruments out of sight	Consistent handling of tissues appropriately with no risk of damage, instruments always in sight
Flow of operation/operative technique		Imprecise, wrong technique in approaching the operative interventions, or constant supervisor corrections	Careful technique in approaching the operative interventions, or constant supervisor corrections	Careful technique with occasional errors or little supervisor correction	Careful technique with occasional errors or little supervisor correction	Fluent, secure, and correct technique in all stages of operative procedure, no supervisor corrections
Total score						

Abbreviation: OSA-LS, Objective Structured Assessment of a Laparoscopic Salpingectomy.

Download English Version:

<https://daneshyari.com/en/article/3950952>

Download Persian Version:

<https://daneshyari.com/article/3950952>

[Daneshyari.com](https://daneshyari.com)