

Accepted Manuscript

Four cases of postoperative pneumothorax among 2814 consecutive laparoscopic gynecologic surgeries: A possible correlation between postoperative pneumothorax and endometriosis

Tetsuya Hirata, M.D., Ph.D., Akari Nakazawa, M.D., Shinya Fukuda, M.D., Yasushi Hirota, M.D., Ph.D., Gentaro Izumi, M.D., Ph.D., Masashi Takamura, M.D., Ph.D., Miyuki Harada, M.D., Ph.D., Kaori Koga, M.D., Ph.D., Osamu Wada-Hiraike, M.D., Ph.D., Tomoyuki Fujii, M.D., Ph.D., Yutaka Osuga, M.D., Ph.D.

PII: S1553-4650(15)00349-0

DOI: [10.1016/j.jmig.2015.04.023](https://doi.org/10.1016/j.jmig.2015.04.023)

Reference: JMIG 2560

To appear in: *The Journal of Minimally Invasive Gynecology*


Received Date: 4 March 2015

Revised Date: 14 April 2015

Accepted Date: 15 April 2015

Please cite this article as: Hirata T, Nakazawa A, Fukuda S, Hirota Y, Izumi G, Takamura M, Harada M, Koga K, Wada-Hiraike O, Fujii T, Osuga Y, Four cases of postoperative pneumothorax among 2814 consecutive laparoscopic gynecologic surgeries: A possible correlation between postoperative pneumothorax and endometriosis, *The Journal of Minimally Invasive Gynecology* (2015), doi: 10.1016/j.jmig.2015.04.023.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Four cases of postoperative pneumothorax among 2814 consecutive laparoscopic gynecologic surgeries: A possible correlation between postoperative pneumothorax and endometriosis

Tetsuya Hirata, M.D., Ph.D., Akari Nakazawa, M.D., Shinya Fukuda M.D., Yasushi Hirota M.D., Ph.D., Gentaro Izumi, M.D., Ph.D., Masashi Takamura, M.D., Ph.D., Miyuki Harada, M.D., Ph.D., Kaori Koga, M.D., Ph.D., Osamu Wada-Hiraike, M.D., Ph.D., Tomoyuki Fujii, M.D., Ph.D., Yutaka Osuga, M.D., Ph.D.

Department of Obstetrics and Gynecology, Faculty of Medicine, University of Tokyo, Tokyo, Japan

Corresponding author: Tetsuya Hirata, M.D., Ph.D.

Department of Obstetrics and Gynecology, Faculty of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, Japan 113-8655

Tel: +81-3-3815-5411 Ext. 33407. Fax: +81-3-3816-2017

E-mail address: thira-tky@umin.ac.jp

Precis: Postoperative pneumothorax incidence in patients who underwent laparoscopic surgery for benign gynecologic disease was 0.14%. Postoperative pneumothorax frequency was significantly higher among cases with endometrioma than among those without.

Keywords: postoperative pneumothorax, laparoscopic surgery, endometriosis

Download English Version:

<https://daneshyari.com/en/article/3957125>

Download Persian Version:

<https://daneshyari.com/article/3957125>

[Daneshyari.com](https://daneshyari.com)