


Accepted Manuscript

The incidence of Interstitial pregnancy after IVF/ET and the outcome of a consecutive series of 38 cases managed by Laparoscopic cornuostomy or cornual repair

Jieyu Wang, Dong Huang, Xiaona Lin, Sotiris H. Saravelos, Jianmin Chen, Xiao Zhang, TinChiu Li, Songying Zhang



PII: S1553-4650(16)00134-5

DOI: [10.1016/j.jmig.2016.02.022](https://doi.org/10.1016/j.jmig.2016.02.022)

Reference: JMIG 2805

To appear in: *The Journal of Minimally Invasive Gynecology*

Received Date: 20 January 2016

Revised Date: 23 February 2016

Accepted Date: 27 February 2016

Please cite this article as: Wang J, Huang D, Lin X, Saravelos SH, Chen J, Zhang X, Li T, Zhang S, The incidence of Interstitial pregnancy after IVF/ET and the outcome of a consecutive series of 38 cases managed by Laparoscopic cornuostomy or cornual repair, *The Journal of Minimally Invasive Gynecology* (2016), doi: 10.1016/j.jmig.2016.02.022.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The incidence of Interstitial pregnancy after IVF/ET and the outcome of a consecutive series of 38 cases managed by Laparoscopic cornuostomy or cornual repair.

Jieyu Wang ¹, Dong Huang ¹, Xiaona Lin ¹, Sotiris H Saravelos ², Jianmin Chen ¹,
Xiao Zhang ¹, TinChiu Li ², Songying Zhang ^{1,3}

1, Department of Obstetric & Gynecology, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou, China

2, Department of Obstetric & Gynecology, Prince of Wales Hospital, The Chinese University of Hong Kong

3, Corresponding author. E-mail address: zhangsongying@126.com

We report incidence and factors of interstitial pregnancy after IVF/ET and suggest laparoscopic cornuostomy or cornual repair. should be a first choice for both intact and ruptured interstitial pregnancy according to our experience.

Download English Version:

<https://daneshyari.com/en/article/3961579>

Download Persian Version:

<https://daneshyari.com/article/3961579>

[Daneshyari.com](https://daneshyari.com)