


Accepted Manuscript

Effect of undiagnosed deep adenomyosis after failed NovaSure endometrial ablation

Bianca B. Mengerink, MD Anneke A.M. van der Wurff, MD, PhD Johanne F. ter Haar, MD Ilse A. van Rooij, MD, PhD Johanna M.A. Pijnenborg, MD, PhD



PII: S1553-4650(14)01442-3

DOI: [10.1016/j.jmig.2014.10.006](https://doi.org/10.1016/j.jmig.2014.10.006)

Reference: JMIG 2400

To appear in: *The Journal of Minimally Invasive Gynecology*

Received Date: 21 August 2014

Revised Date: 4 October 2014

Accepted Date: 7 October 2014

Please cite this article as: Mengerink BB, van der Wurff AAM, ter Haar JF, van Rooij IA, Pijnenborg JMA, Effect of undiagnosed deep adenomyosis after failed NovaSure endometrial ablation, *The Journal of Minimally Invasive Gynecology* (2014), doi: 10.1016/j.jmig.2014.10.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title

Effect of undiagnosed deep adenomyosis after failed NovaSure endometrial ablation

Authors: Bianca B. Mengerink, MD^{a b}

Anneke A.M. van der Wurff, MD, PhD^c

Johanne F. ter Haar, MD^d

Ilse A. van Rooij, MD, PhD^d

Johanna M.A. Pijnenborg, MD, PhD^d

^a Rijnstate Hospital, Department of Obstetrics & Gynecology, Arnhem, The Netherlands

^b Radboud University Medical Centre, Department of Obstetrics & Gynecology, Nijmegen,

The Netherlands

^c St Elisabeth Hospital, Department of Pathology, Tilburg, The Netherlands

^d TweeSteden Hospital, Department of Obstetrics & Gynecology, Tilburg, The Netherlands

Address and correspondence:

Bianca B. Mengerink, MD

Department of Obstetrics & Gynecology, Rijnstate Hospital

PO Box 9555, 6800 TA Arnhem, The Netherlands

Tel: +31-88005888 / Fax: +31-880057878

Email: BiancaMengerink@radboudumc.nl

Synopsis : Deep adenomyosis was present in 40.9% of patients after failed NovaSure endometrial ablation versus 21.4% of control patients undergoing hysterectomy.

Type of article: Clinical article

Download English Version:

<https://daneshyari.com/en/article/3961811>

Download Persian Version:

<https://daneshyari.com/article/3961811>

[Daneshyari.com](https://daneshyari.com)