

Contents lists available at ScienceDirect

# Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com



## Case Report

# Prenatal diagnosis of parapagus diprosopus dibrachius dipus twins with spina bifida in the first trimester using two- and three-dimensional ultrasound



Pei-Yin Yang <sup>a</sup>, Ching-Hua Wu <sup>b</sup>, Guang-Perng Yeh <sup>c</sup>, Charles Tsung-Che Hsieh <sup>c,\*</sup>

- <sup>a</sup> Department of Obstetrics and Gynecology, Chiayi Christian Hospital, Chiayi, Taiwan
- <sup>b</sup> Obstetrics and Gynecology Ultrasound Lab, Changhua Christian Hospital, Changhua, Taiwan
- <sup>c</sup> Department of Obstetrics and Gynecology, Changhua Christian Hospital, Changhua, Taiwan

#### ARTICLE INFO

Article history Accepted 6 March 2014

Keywords: conjoined twins first trimester prenatal diagnosis spina bifida three-dimensional ultrasound

#### ABSTRACT

Objective: Here, we report a case of parapagus diprosopus twins with spina bifida diagnosed in the first trimester of pregnancy using two-dimensional (2D) and three-dimensional (3D) ultrasound.

Case report: A 28-year-old Taiwanese woman, gravid 1, para 0, visited our hospital due to an abnormal fetal head shape discovered by 2D ultrasound at 11-weeks gestation. Parapagus diprosopus twins with spina bifida were diagnosed after ultrasound examination. The characteristics of parapagus diprosopus twins are more illustrative in 3D ultrasound than in 2D ultrasound. After counseling, termination of pregnancy was chosen by the couple. Although necropsy was declined, the gross appearance and radiograph of the abortus confirmed our diagnosis.

Conclusion: With the help of 3D ultrasound, we made an early and definitive diagnosis of conjoined

Copyright © 2015, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

#### Introduction

Conjoined twins are very uncommon, with an incidence of ~1:50,000 to ~1:100,000 [1-3], occurring in 1% of monochorioic twins [1]. The logical theory of the origin of conjoined twins proposed by an expert would be the "spherical theory", which describes the secondary union of two originally separate embryonic discs in one of two areas: the dorsal neural tube or the ventral yolk sac after the 13<sup>th</sup> day of conception [4]. Due to significant advances in ultrasound technology, prenatal diagnosis of conjoined twins in the first trimester is feasible [5]. The first case of prenatal diagnosis of conjoined twins in the first trimester was thoracophalipagus at 11-weeks gestation [6].

Early diagnosis and detailed assessments of the shared vital organs are necessary for optimal management of the pregnancy and birth. With the use of three-dimensional (3D) imaging technology, the space orientation and anatomical details are more

E-mail address: 40129@cch.org.tw (C.T.-C. Hsieh).

clearly demonstrative than with traditional two-dimensional (2D) ultrasonography [2,5]. Here, we report a rare case of parapagus diprosopus twins associated with spina bifida diagnosed at 12weeks gestation by using 2D and 3D ultrasound.

### Case presentation

A 28-year-old female, gravida 1, para 0, visited our hospital at 12 weeks of pregnancy. She had received an ultrasound exam in Vietnam, during which an abnormal fetal head shape was recognized. She returned to Taiwan for the subsequent prenatal exam. The couple are not consanguineous. Neither the woman nor her husband had a family member with a history of twinning or congenital anomalies.

We used a high-resolution ultrasound machine (Voluson 730 Expert, General Electric, Zipf, Austria), an abdominal transducer (AB 2-7 convex probe, 2-7 MHz, General Electric), and a 3D transabdominal transducer (RAB 2-5 L4D probe, 2-5 MHz, General Electric). The crown-rump length was 55 mm, compatible with the gestational age. With 2D ultrasonography, we identified a normalappearing trunk and extremities, one stomach, one heart, and one bladder. However, we also noted widening of the skull like a

<sup>\*</sup> Corresponding author. Department of Obstetrics and Gynecology, Changhua Christian Hospital, Number 135 Nan-hsiao Street, Changhua 500, Taiwan.

helmet. Midline of the brain and bilateral widened central plexuses were traced in the 2D ultrasound (Figure 1). We performed a 3D ultrasound, which indicated a normal body appearance, but a widened thoracic cage and four normal limbs (Figure 2). We then performed a 3D rendering with skeletal mode. Widened cervicothoracic spines were recognized and partial conjunction of spines was suspected (Figure 3). Two posterior fontanelles could be observed with 3D surface rendering. The fusion of the faces was much clearer in the 3D surface rendering. We also used 3D tomographic ultrasound imaging (TUI) to survey the fetal spine and confirmed our suspicion of spina bifida due to an overlying skin defect of the spine (Figure 4). Three orbits were also indicated by the 3D TUI technique (Figure 5). We were not able to define the intracranial structures due to the early gestational age and fetal position.

After discussion with the family, the couples decided to terminate this pregnancy. Although necropsy was declined, the appearance of the abortus was compatible with the results of the 2D and 3D ultrasonography (Figures 6 and 7). There were four limbs and a normal trunk, but a wide thoracic cage, a rounded and wide shared cranial bone, three eyes due to inner-side fusion in two eyes, two noses, and two sets of lips (diprosopus, parapagus, dibrachius, and dipus). The postabortal radiography showed two joined cranial vaults and cervicothoracic spines (Figure 8). All these findings were compatible with our ultrasonographic findings.

#### Discussion

Different types of fusion exist in the cases of conjoined twins. Spencer [4] proposed that conjoined twins should be classified by the proposed site of the union and be divided into two groups: ventral and dorsal [4]. Ventral unions, such as rostral, caudal, and lateral unions, occur most commonly. Dorsal unions, such as craniopagus, rachipagus, and pygopagus occur less frequently [4,7]. Our case was a ventral lateral union, meaning parapagus. Diprosopus was the subtype of the parapagus, meaning two faces.

The rarest type of conjoined twins is the diprosopus, representing only ~0.4% [8]. This type of conjoined twin has two faces, one head, and one body. In 1992, Fontanarosa et al [9] reported a diprosopus twin diagnosed in the first trimester [9]. Initially, their impression was of a fetus with a large encephalocele. They found this fetus with a bifid head, a broad thorax, a single heart, and four normal limbs using 2D ultrasonography. After termination, the abortus had two faces symmetrically and medially fused, craniorachischisis, and left gastroschisis.



Figure 1. Two-dimensional ultrasound of the brain section of our case at 12-weeks gestation.



Figure 2. Three-dimensional ultrasound of our case at 12-weeks gestation using the surface mode



**Figure 3.** Three-dimensional ultrasound of our case at 12-weeks gestation in the skeletal mode. The black arrows indicate the duplication of cervical spines.

Ülker et al [10] presented a case of diprosopus without gastroschisis. Due to the rarity of this condition, the authors were not initially recognized and they had to repeat the exam to ensure an accurate diagnosis. In our case, the diagnosis was easy to make with 3D ultrasound because of its excellent space orientation. We reviewed all of the cases diagnosed as parapagus diprosopus in the first trimester (Table 1). The earliest gestational age at diagnosis was 8-weeks gestation [11]. In that case, obvious bifid cranium poles were noted in early pregnancy, enabling them to make the diagnosis at a very early gestational age. In our case, the gestational age at presenting was 12-weeks gestation, however, our case did not have a bifid cranium. The only clue from the 2D sonography that aided the diagnosis was the abnormal fetal head shape. Our case illustrates the importance of incorporation of fetal skull shape into fetal anatomy screening.

The experts proposed that most cases of parapagus diprosopus have duplicated cervical and upper thoracic spines, and varying

## Download English Version:

# https://daneshyari.com/en/article/3975255

Download Persian Version:

https://daneshyari.com/article/3975255

<u>Daneshyari.com</u>