



## Original Article

# Decreased endometrial vascularity and receptivity in unexplained recurrent miscarriage patients during midluteal and early pregnancy phases



Shu-Yin Tan <sup>a</sup>, Fu Hang <sup>b,1</sup>, Gowreesunkur Purvarshi <sup>b</sup>, Min-Qing Li <sup>b</sup>, Da-Hua Meng <sup>a</sup>, Ling-Ling Huang <sup>b,\*</sup>

<sup>a</sup> Department of Obstetrics and Gynecology, GuangXi Women and Children Hospital, GuangXi, China

<sup>b</sup> Department of Obstetrics and Gynecology, The First Affiliated Hospital of GuangXi Medical University, GuangXi, China

## ARTICLE INFO

## Article history:

Accepted 8 October 2014

## Keywords:

endometrial vascularity  
endometrial volume  
placental volume  
recurrent miscarriage  
three-dimensional power Doppler  
ultrasound

## ABSTRACT

**Objective:** To evaluate the predictive value of three-dimensional (3D)-power Doppler sonography on recurrent miscarriage.

**Materials and methods:** The study patients were divided into a recurrent miscarriage group (30 cases) and a normal pregnancy group (21 cases). Measurement of endometrial thickness was performed using two-dimensional transvaginal ultrasound in the midluteal phase. The endometrial volume, vascularization index (VI), flow index (FI), and vascularization-flow index (VFI) in midluteal and placenta volume, as well as the VI, FI, and VFI of early pregnancy were measured using Virtual Organ Computer-aided Analysis of 3D-power Doppler ultrasound.

**Results:** Endometrial thickness, endometrial volume, endometrial vascular data, VI, FI, and VFI of the midluteal phase were lower in the recurrent miscarriage group compared with the normal pregnancy group ( $p < 0.05$ ). Placental volume, VI, and VFI during early pregnancy were lower in the miscarriage group compared with the normal pregnancy group ( $p < 0.05$ ). There was no significant change in FI between the recurrent miscarriage and control groups during early pregnancy ( $p > 0.05$ ). The predictive accuracy of endometrial thickness, endometrial volume, VI, FI, and VFI in the midluteal phase, and placenta volume, VI, FI, and VFI in early pregnancy as measured by the receiver operating characteristic curve to predict miscarriage before 12 gestational weeks in participants was 0.681, 0.876, 0.770, 0.720, 0.879, 0.771, 0.907, 0.592, respectively.

**Conclusion:** The 3D-power Doppler ultrasound is a more comprehensive and sensitive method for evaluating endometrial receptivity. Endometrial volume, VI, FI, and VFI in the midluteal phase, as well as VI in early pregnancy, can be considered as predictive factors for recurrent miscarriage.

Copyright © 2015, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

## Introduction

Recurrent miscarriage (RM) affects approximately 3% of women of reproductive age [1]. Genetic, endocrine, infectious, and anatomical factors, as well as autoantibodies, abnormal prothrombotic state, and other aspects are involved in the etiology

[2–5]. Endometrial receptivity can also be explained as an endometrial state allowing embryo adhesion, invasion, and implantation [6]. Endometrial receptivity correlates with endometrial angiogenesis and the degree of vascularization [7]. Endometrial angiogenesis and vascularization form the basis for the formation of the decidual capillary network in the placenta and the foundation of embryonic growth and development. Endometrial and subendometrial hypoxia caused by inadequate blood flow induces low receptivity, which decreases embryo implantation and increases spontaneous abortion [8].

In recent years with the development of ultrasonic technology, an effective nontraumatic detection technique has been provided

\* Corresponding author. Department of Obstetrics and Gynecology, The First Affiliated Hospital of GuangXi Medical University, Number 6 Shuang Yong Road, NanNing, 530021, GuangXi, China.

E-mail address: [huanglingling00@aliyun.com](mailto:huanglingling00@aliyun.com) (L.-L. Huang).

<sup>1</sup> Co-first author.

for evaluating endometrial receptivity. Several ultrasonic evaluation parameters for endometrial receptivity have been put forward, including endometrial thickness, endometrium types, endometrial volume, and endometrial and subendometrial blood flow. Three-dimensional (3D) ultrasonography using power Doppler angiography (3D-CPA) accurately measures endometrial volume and vascular parameters, including: (1) vascularization index (VI), which represents the number of blood vessels in the measured region; (2) blood flow index (FI), which represents the blood flow strength sensed over 3D-scanning time in the measured region; and (3) vascularization flow index (VFI), which represents the sum of blood flow and vascularization in the measured region. These three parameters reflect endometrial and placental blood supply sensitivity. Hafner et al [9] found that placental clearance flow could be detected through vaginal 3D-CPA at 5-weeks gestation, and that the VI and VFI positively correlated with crown-rump length and embryonic bud size, which provided the basis for the study of early pregnancy placental blood flow parameters using ultrasonic monitoring.

Most research focuses on the measurement of midluteal phase endometrial volume and vascular parameters to predict assisted reproductive success rate. Few investigations have explored predicting the risk of recurrent miscarriage using 3D ultrasound and no study has used early pregnancy placental volume and vascular parameters to evaluate the risk of recurrent miscarriage. The present study investigates endometrial volume and vascular parameters in the midluteal phase of recurrent miscarriage patients and follows patients until pregnancy, measuring first-trimester placental volume and blood flow parameters and comparing these data to normal fertile women in order to clarify the predictive value of 3D-power Doppler ultrasound for recurrent miscarriage.

## Materials and methods

### Patients

The experiment was conducted in the prenatal diagnosis center of The First Affiliated Hospital of GuangXi Medical University and GuangXi Women and Children Hospital (GuangXi, China). Signed informed consent was obtained from all participants and the study was approved by the Research Ethics Board of the First Affiliated Hospital of GuangXi Medical University.

From January 2013 to January 2014, 122 women were enrolled in the study and 51 women remained until study completion. The experimental design was a case control study. The women were divided into two groups: the recurrent miscarriage (study) group and the normal pregnancy (control) group. Inclusion criteria for the study group were: (1) the pregnancy was ending before 12 gestational weeks; (2) patients had experienced two or more spontaneous abortions; (3) parents both had no chromosomal abnormalities; (4) patients had no abnormality of the uterus; (5) patients had no history of toxic material exposure; and (6) patients had a healthy medical history outside of miscarriage. Inclusion criteria for the control group were: (1) the pregnancy lasted until 12 gestational weeks; (2) patients had no history of spontaneous abortion; (3) patients had no abnormality of the uterus; (4) patients had no history of toxic material exposure; and (5) patients had a healthy medical history. Case baseline assessment included age.

### Ultrasound data acquisition

Using a Voluson E8 Ultrasound (GE Medical Systems Kretztechnik GmbH, Zipf, Austria), a ~5–9-MHz transvaginal volumetric

probe, and Virtual Organ Computer-aided Analysis (VOCAL) software (GE Medical Systems Kretztechnik GmbH), all patients received a transvaginal ultrasound scan performed by a single investigator unaware of participant status.

The entire endometrial section was displayed in uterine longitudinal sections using two-dimensional (2D) Doppler ultrasonography. Taking the maximum distance of the interface between the diameter of anterior and posterior uterine myometrium, then perpendicular across the midline of the endometrium as endometrial thickness, three continuous measurements were taken and averaged.

Using the optional multiplatform model and a volume angle of 120°, the volume of the sampling frame size was adjusted to ensure that the endometrium was completely oriented within the sampling frame. Volume scanning was then started and three planar volumetric measurements obtained and stored for analysis. The best reference plane was selected to display the endometrial cavity from the uterus fundus to the cervical. The VOCAL software application was used to manually sketch and set the volume of data extraction for every aspect angle within 30°, then trace the outline of the endometrium in each section. The system automatically calculated and displayed the volumetric results after completion of outlining, then applied 3D histogram-analysis software to calculate the energy in order to obtain endometrial volume flow parameters, VI, FI, and VFI.

Intrauterine early pregnancy can be diagnosed using B-scan ultrasonography, showing a “double ring sign” gestational sac or yolk sac. At the beginning of 6–8 weeks in early pregnancy, the placenta can be identified using B-scan ultrasonography. Placenta essence can be explored using a stronger uniform light spot than echogenicity of the surrounding muscle tissue, which is mostly “crescent-shaped.” All patients received an early pregnancy ultrasound examination using the methods described above and the placental volume, endometrial volume flow parameters VI, FI, and VFI were measured.

Recurrent miscarriage patients who met the inclusion criteria were recruited. Their ovulation was monitored using transvaginal ultrasound for the first 10 days of menstruation until ovulation was confirmed. At 5–7 days after ovulation, midluteal-phase (embryo in the womb, endometrial implantation window) endometrial thickness was measured using 2D Doppler ultrasonography, and 3D-power Doppler ultrasound was used to measure the endometrial volume, VI, FI, and VFI. At 14–16 days after ovulation (pregnant 4 weeks), patients who confirmed pregnancy with serum  $\beta$ -Human chorionic gonadotropin ( $\beta$ -HCG)  $\geq 100$  mIU/mL and progesterone  $\geq 20$  ng/mL had their cycles continuously monitored before measuring placental volume and VI, FI, and VFI at 6–8 weeks of pregnancy. These patients were included in the study group if spontaneous abortion occurred before 12 weeks of pregnancy.

The women who met the inclusion criteria for the control group underwent the same monitoring protocol as the study group. These women were followed up to 12 weeks and then included in the control group if spontaneous abortion had not occurred before 12 gestational weeks.

### Statistical analysis

Using SPSS version 18.0 software (SPSS, Inc., Armonk, NY, USA) for statistical analysis, measurement data were expressed as mean  $\pm$  standard deviation. The independent samples *t*-test was applied to compare data. A *p* value  $< 0.05$  was considered statistically significant. Receiver operating characteristic (ROC) curves were applied to compare the predicted performance of different indicators.

Download English Version:

<https://daneshyari.com/en/article/3975471>

Download Persian Version:

<https://daneshyari.com/article/3975471>

[Daneshyari.com](https://daneshyari.com)