



Original Article

# The Outcomes of Anterior Spinal Fusion for Cervical Compressive Myelopathy—A Retrospective Review

## 壓迫性脊髓型頸椎病之前路手術治療效果的回顧分析

Suen Tsz-King\*, Wong Kwok-Ho, Ho Yuen-Fong

Department of Orthopaedics and Traumatology, Caritas Medical Centre, Kowloon, Hong Kong

### ARTICLE INFO

*Article history:*

Accepted September 2010

*Keywords:*

anterior spinal fusion  
cervical myelopathy  
prognostic factors

### ABSTRACT

**Background:** This study examined the prognostic factors that affect the surgical outcome of anterior spinal fusion in patients with cervical compressive myelopathy (CCM).

**Methods:** A retrospective review of the result of surgical treatment in 49 cases of CCM using anterior spinal fusion surgical techniques. The study period was from June 1990 to June 2008, with a mean follow-up of 6.57 years. Seven patients died during the follow-up period because of other medical diseases. The Japanese Orthopaedic Association score for cervical myelopathy and recovery rate were used to assess the functional results of the remaining 42 cases.

**Results:** A mean Japanese Orthopaedic Association score improvement from 10.21 to 14.23 points after the operation was observed. The overall recovery rate was 65.38%, with excellent or good results of 80%. After a year of follow-up, we did not find significant improvement or deterioration on the neurological status on further follow-up. Acute complications were not common. One patient underwent re-operation because of slipping of the graft during resuscitation after early post-operative acute myocardial infarction. This patient showed post-operative neurological deterioration. Two patients had subsidence of the graft, three had pseudo-arthritis, and two had graft resorption. There were also three patients with degenerative changes of the adjacent vertebra. No re-operation was done for patients with these late complications in this review.

**Conclusion:** Anterior cervical decompression with bone fusion is a viable surgical option for patients with one level of anterior cervical cord compression, especially for patients with kyphosis or straight canal spine. For patients with two- to three-level involvement, anterior cervical decompression with bone fusion provides good functional result in proper selection of cases. We also identified some prognostic factors (male sex, symptoms less than 1 year, and age less than 70 years) in predicting a favourable outcome of anterior spinal fusion for CCM.

### 中文摘要

**背景:** 本研究探索經前路手術(ASF)處理頸椎壓迫性脊髓病 (Cervical Compressive Myelopathy) 的一些影響手術結果之預後因素

**方法:** 本研究回被包括1990-2008年間曾採用盆骨移植前路手術(ASF)處理頸椎壓迫性脊髓病 (Cervical Compressive Myelopathy) 病例共49宗, 平均隨訪時間約6年半。當中七名病人在研究期間因其他內科疾病去世。其餘42名病人在手術前後採用日本骨科學會的頸椎脊髓病評分方法(JOA score)評估功能復原結果。

**結果:** 術後的平均JOA score評分由10.21改善到14.23分, 平均復原率為65.38%, 百分之八十病人有良好或以上功能之復原。手術一年後隨訪, 病人的神經的功能已經沒有顯著改變。手術後初期引致急性的併發症並不常見。只有一名病人因急性心肌梗塞需要心肺復甦術而引致在術後盆骨移植移位而要重做手術, 她是唯一術後功能比手術前差的個案。在隨訪期間, 骨融合不完全而引至假關節及附近頸椎有提早退化跡象各有三例, 另外有骨移植吸收及下陷各有兩例, 但沒有病人因以上延遲的併發症需要再做手術。

**總結:** ASF適用於一節段頸椎前路脊髓受壓的病人, 尤其是患者有頸椎後凸畸形或直的頸椎管, 是其中一種可行的手術。患者有兩或三節段頸椎脊髓受壓時, 術後亦有良好的功能性復原, 但需要經過適當地選擇個案。我們亦發現一些影響手術結果之預後因素, 如男性、70歲以下及病徵1年以內的病人會有較佳的復原。

\* Corresponding author. E-mail: aarsuen@gmail.com.

## Introduction

Cervical compressive myelopathy (CCM) is a spinal cord dysfunction as a result of a slow and gradual mechanical compression and vascular compromise. Patients with underlying spinal stenosis may deteriorate rapidly after acute neck injury. In general, operation should be considered when the patients have marked disturbance of activities of daily living and walking gait. The recommended treatment was cervical spinal cord decompression preserving spinal stability and mobility. The choices of operations include laminoplasty and anterior cervical decompression with bone fusion (ASF) depending on the causes, extension, direction of compression, presence of neck pain, and surgeons' experience.<sup>1,2</sup>

The effectiveness of using laminoplasty in the treatment of multilevel cervical stenosis is well described in the literature.<sup>3,4</sup> ASF was suggested for patients with more localised cervical disc degeneration and for patients with spinal kyphosis. Some surgeons used spinal plating at the same operations to achieve immediate stability and to improve the fusion rate. However, this additional procedure may increase the operation time and complications of the implants.

In the literature, some<sup>5,6</sup> reported less favourable outcomes in elderly patients with history of preceding trauma and long duration of prominent neurological signs.<sup>7</sup> Other unfavourable factors identified in the patients included lower radiological Pavlov ratio (PR), lower cord compression ratio (CCR), and T2 cord signal changes in magnetic resonance imaging (MRI).<sup>8</sup>

The exact incidence of CCM is unknown in Hong Kong. There were very few reports related to the surgical outcomes of the local Chinese patients. The objective of this study was to review the operative results of the patients in our hospital and to evaluate the prognostic factors that are applicable to our local community.

## Materials and Methods

From 1990 to 2008, there were 49 consecutive patients with CCM who had ASF operations done in our department. Seven patients died from unrelated causes during follow-up. Thus, a total of 42 patients were recruited for the study. ASF was limited to those with three or less levels of involvement. We did not use internal fixation in all cases.

### Surgical technique

The Smith-Robinson approach<sup>9</sup> was used. The right neck surgical approach was used in all the cases. The exact spinal segments were determined using intra-operative radiographs. Disectomy with partial corpectomy was done for single-level disc disease. Subtotal corpectomy was performed when two adjacent degenerative discs had to be tackled. Excision of posterior longitudinal ligament was not routinely performed. ASF was done using a tri-cortical bone graft taken from the iliac crest. Internal fixation using plating was not done in our series.

Philadelphia neck collar was applied after the operation for single-level operation. For cases with two or three levels of bone fusions done, a halo-vest was used for 2–4 weeks before changing to a semi-rigid neck collar. Normally, the neck collar will be worn till bone fusion occurred at 8–12 weeks.

### Assessment

The functional results of the operation were assessed using the CCM scoring system proposed by the Japanese Orthopaedics

Association (JOA).<sup>10</sup> The functional improvement was expressed by the recovery rate percentage (RR%) as adopted by the JOA, and the final results were rated as excellent, good, fair, unchanged, and worsen, shown as follows:

1.  $RR\% = \frac{\text{post-score} - \text{pre-score}}{\text{pre-score}} \times 100\%$
2. Functional result basing on RR%:  
Excellent, from 75% to 100%  
Good, from 50% to 74%  
Fair, from 25% to 49%  
Unchanged, from 0% to 24%  
Worsen, less than 0%.

The patient sex, age, trauma history, symptom duration, and severity of the neurological signs were analysed. All the abnormal neurological signs detected clinically were classified into early or late signs using the criteria shown in Table 1.

The pre- and post-operative PR, the CCR, and signal change in the T2 image of MRI were also recorded. All the results were analysed using the Pearson correlation and  $\chi^2$  test.

## Results

From July 1990 to June 2008, 42 patients with ASF operation done were included in the study, and their results are shown in Table 2. The mean follow-up was 6.57 years (range, 10 months to 18 years). The average patient age was  $56.45 \pm 15.11$  years (range, 28–86 years). The sex ratio of male to female was 30:12.

The causes of CCM were disc prolapse (15), spondylosis (26), and ossification of posterior longitudinal ligament (1). The levels were affected in the frequency of C5/6 (31), C4/5 (24), C3/4 (15), C6/7 (10), and C2/3 (2) spinal segments.

No patient required combined approach. The ASF was of one level in nine cases, two levels in 26 cases, and three levels in seven cases. The average level of involvement was  $1.95 \pm 0.62$ .

Concomitant spinal disorders and other joint diseases were not uncommon; eight patients had associated lumbar spinal stenosis, and among them, three had an operation done sometime earlier. There were three patients with history of stroke and five with prostatism.

### JOA score

The mean JOA score improved from  $10.21 \pm 3.49$  points to  $14.23 \pm 2.83$  points. The overall RR% was  $65.38 \pm 26.36\%$ . The JOA functional rating showed 14 cases being excellent, 20 good, 6 fair, 1 unchanged, and 1 worsened. Therefore, 80% of the patients had good or excellent improvement of their symptoms. Their relevant data are illustrated in Table 2.

### Prognostic factors

The statistical significances of the different prognostic factors with good to excellent functional results are shown in Table 3. The

**Table 1**  
Summary of early and late myelopathic signs

| Early myelopathic signs     | Late myelopathic signs       |
|-----------------------------|------------------------------|
| Didiachokinesia             | Gross spastic                |
| Fine motor deficits         | Gross motor deficits         |
| Mild increase in jerk       | Marked increase in jerk      |
| Mild or unsustained clonus  | Sustained clonus             |
| Decreased proprioception    | Gross difficulty in balance  |
| Difficulty with tandem gait | Difficulty with routine gait |
| FES Grade 1–2               | FES Grade 3–4                |

Download English Version:

<https://daneshyari.com/en/article/4072709>

Download Persian Version:

<https://daneshyari.com/article/4072709>

[Daneshyari.com](https://daneshyari.com)