




Contents lists available at ScienceDirect

## Trauma Case Reports

journal homepage: [www.elsevier.com/locate/tcr](http://www.elsevier.com/locate/tcr)



### Case Report

# A novel technique for reimplanting extruded bone fragments in open fractures

Sameer Rathore<sup>\*</sup>, Indukuri Viswanatha Reddy<sup>1</sup>, A.H. Ashwin Kumar<sup>2</sup>

Department of Orthopaedics, Block 3, 3rd Floor, Krishna Institute of Medical Sciences, Secunderabad 500003, India

#### ARTICLE INFO

##### Article history:

Accepted 28 May 2016

Available online 25 June 2016

##### Keywords:

Open fractures

Bone loss

Extruded bone fragments

Reimplantation of extruded bone fragments

Management of bone loss

#### ABSTRACT

Extruded bone fragments are a rare complication of high-energy open fractures. Generally, management is thorough debridement and managing the bone defect. In the literature, there are only a few case reports where successful retention of the free bone fragment has been done. Disinfection of bone fragment is done by autoclaving or use of antiseptic/antibiotic solution. Autoclaving leads to complete loss of viable cells and antiseptic/antibiotic solutions do not disinfect completely. In this case report, authors present an innovative technique of disinfecting the bone fragment effectively with minimum compromise on biology.

A 38-year-old male with compound grade III B comminuted fracture of distal femur with 2 extruding bone pieces was managed by thorough debridement, external fixator and antibiotic cement spacer. The extruded bone fragments were rinsed in saline and diluted betadine and implanted in subfascial plane in healthy soft tissues in the thigh along with a few antibiotic beads for assuring disinfection. After 1 week, when no clinical signs of infection were found, the site was opened, cement spacer removed, free fragments positioned anatomically and rigid internal fixation was done. Fracture united at 6 months with good functional outcome. At last follow-up at 1 year, the patient was mobilising freely and there were no signs of low grade infection.

The key points of this procedure are:

- 1) Viability of bone fragment maintained while achieving disinfection.
- 2) Traumatized soft tissues healed and prepared for accepting the free bone fragment.

<sup>\*</sup> Corresponding author. Tel.: +91 8989457001, +91 9573007744; fax: +91 040 27840980.

E-mail addresses: [dr.sameer.rathore@gmail.com](mailto:dr.sameer.rathore@gmail.com) (S. Rathore), [vishortho@yahoo.com](mailto:vishortho@yahoo.com) (I.V. Reddy), [ashwin\\_ortho@yahoo.com](mailto:ashwin_ortho@yahoo.com) (A.H. Ashwin Kumar).

<sup>1</sup> Tel.: +91 9701007892; fax: +91 040 27840980.

<sup>2</sup> Tel.: +91 7 680 993 355; fax: +91 040 27 840 980.

- 3) Use of antibiotic cement counters any remaining chances of infection after thorough debridement.
- 4) Faster union with maintenance of bone length and alignment with use of anatomic fragments.

Extensive search of literature was done and this procedure was found to be novel. A larger case series can help in determining the utility of this technique in compound fractures.

© 2016 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

---

## Introduction

Extruded bone fragments are a rare complication seen in high-energy open fractures. But with increase in road traffic accidents, the incidence of such high-energy trauma is on the rise. Generally, management of such open fractures is through debridement, including taking out of the free fragments of bone and then management of the bone gap [1]. The management of the bone defect can be done by bone transport, fibula strut grafting, cancellous bone grafting or Masquelet technique. In a situation where the extruded segment of bone is available for reimplantation, many benefits include maintenance of skeletal and soft tissue length, averting the morbidity associated with autograft harvest and obviating the need for allograft bone or prolonged bone transport procedures.

In the literature, there have only been a few case reports where successful reimplantation of extruded bone fragments has been done. In all the published case reports, disinfection of the bone has been done by autoclaving, antibiotic or antiseptic treatment [2–6]. A few studies have also been done on the treatment of contaminated bone by these modalities and they show loss of osteogenic potential in the treated bone either completely or to a significant extent. Results of these studies show that autoclaving and chlorhexidine treatment disinfects the bone satisfactorily but leaves no viable cells in the graft [4,7,8]. Treatment with povidone-iodine does not disinfect the bone in all instances [4,7]. Reimplantation of bone segment with no viable cells is equivalent to using an allograft with no osteogenic potential. Incomplete disinfection attaches high risk to the procedure and can lead to devastating complication of infection.

In this paper, the authors wish to present a case report of successful reimplantation of an extruded cortico-cancellous fragment of distal femur by a novel technique which preserves the viability of the bone fragment as well as counteracting infection in an effective manner.

## Case report

A 38-year-old male suffered a RTA and had a compound grade IIIb fracture of the distal femur with intercondylar comminution and extruded bone fragments. One fragment was cortico-cancellous, measuring roughly 6 cm × 3 cm and another was cortical measuring roughly 3 cm × 2 cm. Both bone fragments had no soft tissue attachments. The fracture was thoroughly debrided and extruded bone pieces were rinsed in saline and diluted povidone-iodine solution. A knee spanning external fixator was applied maintaining fracture alignment and length. The bone defect was filled with antibiotic cement spacer. The spacer was composed of 40 g of PMMA mixed with 4 g of vancomycin and 500 mg of gentamycin. The free fragments were then placed in the subfascial plane in the thigh through an incision that was extended for debridement of the wound. A few antibiotic cement beads on an SS wire were placed alongside the fragments of the bone. The wound was then closed primarily.

Intravenous antibiotics were started from the time of presentation. Injectable Cefuroxime 1.5 g i.v. BD was given for 2 weeks, injectable Amikacin 500 mg i.v. BD and injectable Metrogyl 500 mg i.v. BD was given for 2 weeks. No oral antibiotics were given further.

Patient was kept on external fixator for 1 week. In this 1 week, the wound did not show any signs of infection. So patient was planned for re-exploration and definitive fixation of the fracture. On opening the wound in the operation theatre, it was found to be healthy with improvement in gross texture of tissue.

Download English Version:

<https://daneshyari.com/en/article/4100508>

Download Persian Version:

<https://daneshyari.com/article/4100508>

[Daneshyari.com](https://daneshyari.com)